



**HAL**  
open science

# Numerical aspect of uniform ensemble controllability for linear systems

Jérôme Lohéac, Martin Lazar

► **To cite this version:**

Jérôme Lohéac, Martin Lazar. Numerical aspect of uniform ensemble controllability for linear systems. 2024. hal-04850750

**HAL Id: hal-04850750**

**<https://hal.science/hal-04850750v1>**

Preprint submitted on 20 Dec 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Numerical aspect of uniform ensemble controllability for linear systems

Martin Lazar<sup>1</sup>

Jérôme Lohéac<sup>2</sup>

December 20, 2024

## Abstract

We consider the ensemble controllability problem for a linear time-invariant system  $\dot{x}(t, \theta) = A(\theta)x(t, \theta) + B(\theta)u(t)$ , where  $A$  and  $B$  are continuous matrices with respect to the parameter  $\theta$ , which belongs to some compact set  $\Theta \subset \mathbb{R}$ . Given any continuous initial state datum  $\theta \mapsto x^0(\theta)$  and any continuous target state  $\theta \mapsto x^1(\theta)$ , we investigate the numerical computation of a  $\theta$ -independent open loop control  $u$  such that  $x^0$  is steered, in a given time  $T > 0$ , at a distance  $\varepsilon > 0$  of  $x^1$  in the uniform norm (with respect to the parameter).

We approach the problem both theoretically and numerically. Using the Fenchel-Rockafellar duality, we first prove the existence and uniqueness of the ensemble control of a minimal  $L^2$  norm. The numerical recovery of the optimal control is obtained by solving the dual problem, which consists in the unconstrained minimization of a non-differentiable functional in the space of Radon measures.

**Keywords:** parameter dependent systems, ensemble controllability, linear control system, Fenchel-Rockafellar duality, numerical algorithms.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries</b>	<b>3</b>
<b>3</b>	<b>Minimal <math>L^2</math>-norm controls</b>	<b>5</b>
3.1	Optimality conditions	5
3.2	Continuity properties of the minimizers	10
<b>4</b>	<b>Systems with a finite number of parameter values</b>	<b>14</b>
<b>5</b>	<b>Discretized problem and convergence</b>	<b>18</b>
<b>6</b>	<b>Numerical strategies</b>	<b>22</b>
6.1	Greedy coordinate descent	22
6.2	Reduced minimization problem	26
6.3	Proposed numerical strategies	26
<b>7</b>	<b>Numerical examples</b>	<b>27</b>
<b>A</b>	<b>Proof of Proposition 6.1</b>	<b>32</b>
<b>B</b>	<b>Some unreachable states</b>	<b>35</b>

---

<sup>1</sup>Faculty of Electrical Engineering and Applied Computing, University of Dubrovnik, Croatia

<sup>2</sup>Université de Lorraine, CNRS, CRAN UMR 7039 Nancy, France

# 1 Introduction

The concept of ensemble controllability [1, 8, 16, 21, 23] needs no more publicity nowadays with regard to its practical importance. Its aim is to control systems whose natural dynamic depends on some unknown or uncertain parameters in a robust and computationally efficient manner. In particular, the goal is to find a parameter invariant control that performs well for each particular realization of the parameters. This kind of problem arises for instance in the transport of quantum particles, where the goal is to develop external excitation that can simultaneously steer the ensemble of systems with variation in their internal parameters from an initial state to a desired final state [20]. One can also see [3] where it is used to study the controllability of the Bloch equation, for an ensemble of non-interacting half-spins, in a static magnetic field, with dispersion in the Larmor frequency. Other real life practical example can be found, as for instance in non-holonomic systems theory where ensemble controllability is used to derive an approximate steering algorithm for a non-holonomic unicycle in the presence of model perturbation [4]. Nowadays, however, the most widely application is related to machine learning, in particular to training processes in supervised learning. By using the dynamic perspective of neural networks, the problem of their training can be recast as a control problem in which the network coefficients (weights and biases) stand for data(parameter)-independent controls [13].

From the generic notion of ensemble controllability, many other notions emerged as the one of uniform ensemble controllability [16, 20, 21], uniform ensemble reachability,  $L^q$ -ensemble reachability [11]. Even if this topic is studied actively, it is far from being well understood. The necessary and sufficient condition of ensemble controllability can be obtained as a special form of the generalization of the Kalman rank theorem to infinite dimensional Banach spaces (e.g. [28, Theorem 3.1.1]). However, such a characterization is primarily theoretical and not suitable for checking the controllability properties of a particular system. In the above-mentioned non-exhaustive references, the authors have derived necessary and/or sufficient conditions for linear time invariant systems to be uniformly ensemble controllable in some special cases. From these references, especially [8, 10, 11], it could be seen that the topology of  $\Theta$  plays a crucial role in the controllability issue. Roughly speaking, for regular enough parameter dependent systems, uniform ensemble controllability cannot hold if the compact set  $\Theta$  is homeomorphic to some compact subset of  $\mathbb{R}^d$  (with  $d \geq 2$ ) with non-empty interior.

Most papers on the topic analyze conditions under which a system is ensemble controllable, but do not consider the associated optimization problem of finding a feasible control of minimal norm, nor do they deal with numerical recovery of the solution. An approximate construction of a feasible control is provided only in some special cases (e.g. [3, 4]). In [15] the authors consider an associated penalization problem in which the constraint of reaching the prescribed target with some a priori given precision is replaced by an additional term in the cost functional that penalizes deviation of the final state from the target. Consequently, such a problem consists in the unconstrained minimization of a smooth functional and is numerically easier to handle. In [25], the uniform ensemble control problem is replaced by a simultaneous one (consisting of hitting a reachable target). The author discusses the strategies by which the latter can be approximated by its discretized version (which is equivalent to an exact control problem for a large, finite-dimensional system). Let us also mention [26], where an ensemble control is designed using polynomial approximations. However, the results of [26] seem only applicable for single input systems.

In [18] we have characterized the optimal ensemble control in the discrete case (i.e., for a finite number of parameters). Then, assuming Lipschitz regularity, we showed how to obtain a sub-optimal control for the continuous case. In the present article, we directly tackle the case of a continuous set  $\Theta$ , and derive optimality conditions. Based on them we propose several numerical strategies for recovering the minimal norm control. More precisely, we prove the existence and uniqueness of a minimal norm ensemble control, and express it through the solution of the corre-

50 sponding dual problem, which consists in the unconstrained minimization of a non-differentiable  
 51 functional in the space of Radon measures. The numerical strategies that we propose in this paper  
 52 are obtained using the optimality condition on the adjoint problem.

53 The paper is organized as follows. In the next section, we recall the notion and some basic  
 54 results on uniform ensemble controllability. In Section 3, we introduce the optimization problem of  
 55 finding a uniform ensemble control of a minimal  $L^2$  norm and derive the associated dual problem.  
 56 We demonstrate the existence of a unique solution for both problems and list their properties. In  
 57 particular, the continuity of solutions with respect to the problem entries is obtained. Section 4  
 58 deals with a special but important case of discretized parameters (obtaining only a finite number of  
 59 values), where we also discuss the sparsity properties of the optimal measure. The convergence of  
 60 solutions when passing from discrete to continuous parameter setting is demonstrated in Section 5.  
 61 Numerical iterative algorithms are presented in Section 6, while their performances are checked  
 62 and discussed on several particular examples in Section 7. Appendices A and B contain some  
 63 technical and related results.

64 **Notation.** The following notations are used in the paper.  
 65  $\mathbb{N} = \{0, 1, 2, \dots\}$  stands for the set of natural numbers including zero, while  $\mathbb{N}^*$  denotes the set of  
 66 positive integers.  $\mathbb{R}_+$  is the set of nonnegative real numbers, and we set  $\mathbb{R}_+^* = \mathbb{R}_+ \setminus \{0\}$ . For a  
 67 matrix  $M \in \mathbb{R}^{n \times m}$ ,  $M^* \in \mathbb{R}^{m \times n}$  is the transpose matrix of  $M$ .  $|\cdot|$  stands for the Euclidean norm  
 68 in  $\mathbb{R}^n$ , and  $\langle \cdot, \cdot \rangle$  is the scalar product on  $\mathbb{R}^n$ . The open (respectively closed) ball of  $\mathbb{R}^n$  centered on  
 69 some  $x$  and of radius  $\varepsilon > 0$  is denoted by  $B(x, \varepsilon)$  (respectively  $\bar{B}(x, \varepsilon)$ ).  
 70 For  $X$  a Banach space and  $f : X \rightarrow \mathbb{R} \cup \{\infty\}$ , we define  $\text{dom } f = \{x \in X \mid f(x) < \infty\}$ , and  $\text{cont } f$   
 71 is the set of points  $x \in \text{dom } f$  where  $f$  is continuous. For  $S \subset X$ ,  $\text{int } S$  is the interior of  $S$ .  
 72 Given  $n \in \mathbb{N}^*$ , by  $\mathcal{C}(\Theta)^n$  we denote the space of continuous complex vector functions on  $\Theta \subseteq \mathbb{R}$   
 73 equipped with  $L^\infty$  norm, i.e.,  $\mathcal{C}(\Theta)^n = \mathcal{C}(\Theta; \mathbb{R}^n)$  and  $\|f\|_{\mathcal{C}(\Theta)^n} = \sup_\theta |f(\theta)|$ .  $\mathcal{B}_\infty(y, \varepsilon)$  is the closed  
 74 ball in the space  $\mathcal{C}(\Theta)^n$  centered in  $y$  and of radius  $\varepsilon$ .  
 75  $\mathcal{M}(\Theta)$  stands for the set of Radon measures supported on  $\Theta$ , while by  $\mathcal{M}(\Theta)^n$  we  
 76 denote the dual space to  $\mathcal{C}(\Theta)^n$  with the dual product given by  $\langle \mu, f \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} =$   
 77  $\sum_{i=1}^n \int_\Theta f_i(\theta) d\mu_i(\theta) = \int_\Theta f(\theta)^* d\mu(\theta)$ . In addition, for  $F \in \mathcal{C}(\Theta)^{n \times m}$ , we define  $F^* \flat \mu \in \mathcal{M}(\Theta)^m$   
 78 by  $\langle F^* \flat \mu, f \rangle_{\mathcal{M}(\Theta)^m, \mathcal{C}(\Theta)^m} = \langle \mu, Ff \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} = \int_\Theta (f(\theta)^* F(\theta)^*) d\mu(\theta)$ .

## 79 2 Preliminaries

80 Let  $\Theta$  be a compact subset of  $\mathbb{R}$  and consider for every constant  $\theta \in \Theta$  the system described by

$$\dot{x}(t, \theta) = A(\theta)x(t, \theta) + B(\theta)u(t) \quad (t > 0, \theta \in \Theta). \quad (2.1)$$

81 In (2.1), we assume that  $A \in \mathcal{C}(\Theta)^{n \times n}$  and  $B \in \mathcal{C}(\Theta)^{n \times m}$ . The input  $u$  is independent of the  
 82 parameter  $\theta$ , and  $\dot{x}$  is the derivative of  $x$  with respect to the time variable  $t$ . Given an initial state  
 83  $x^0 \in \mathcal{C}(\Theta)^n$  and an input  $u \in L^2(0, T)^m$ , the solution of (2.1) at time  $t$ , is denoted by  $x(t, \theta; x^0, u)$ .  
 84 More precisely, we have,

$$x(t, \theta; x^0, u) = e^{tA(\theta)}x^0(\theta) + \int_0^t e^{(t-\tau)A(\theta)}B(\theta)u(\tau) d\tau \quad (t \geq 0, \theta \in \Theta). \quad (2.2)$$

85 **Definition 2.1** (Uniform Ensemble Controllability). The system (2.1) is said to be *uniformly*  
 86 *ensemble controllable* if for any  $x^0, x^1 \in \mathcal{C}(\Theta)^n$  and any  $\varepsilon > 0$ , there exist a time  $T > 0$  and a  
 87 control  $u \in L^2(0, T)^m$ , such that

$$\sup_{\theta \in \Theta} |x(T, \theta; x^0, u) - x^1(\theta)| \leq \varepsilon.$$

88 The terminology *uniform* in the above definition is used to emphasize the fact that the set of  
 89 output functions is the continuous one endowed with the uniform norm. This definition actually  
 90 makes sense since for any continuous initial state datum  $x^0$  and any  $\theta$ -independent input  $u \in$   
 91  $L^2(0, T)^m$ , the output trajectory is continuous with respect to the parameter  $\theta$ .

92 **Remark 2.2.** The systems considered in the present paper focus on the case where  $A(\theta)$  and  $B(\theta)$   
 93 are real matrices. However, all the obtained results can be applied to complex matrix, having in  
 94 mind that if  $A(\theta) \in \mathbb{C}^{n \times n}$  and  $B(\theta) \in \mathbb{C}^{n \times m}$  are complex matrices, then the complex system can  
 95 be identified to a real system with matrices  $\begin{pmatrix} \Re A & -\Im A \\ \Im A & \Re A \end{pmatrix} \in \mathbb{R}^{2n \times 2n}$  and  $\begin{pmatrix} \Re B & -\Im B \\ \Im B & \Re B \end{pmatrix} \in \mathbb{R}^{2n \times 2m}$ ,  
 96 where  $\Re$  and  $\Im$  denote the real and imaginary part respectively.

97 Let us also mention that it is classical that the set of parameter  $\Theta$  is assumed to be a subset of  $\mathbb{C}$ .  
 98 In this paper we only consider the case where  $\Theta \subset \mathbb{R}$ . This assumption is made, firstly, because  
 99 it avoids technical difficulties and, secondly, because, as said in the introduction, from [8, 10, 11]  
 100 typical sets  $\Theta \in \mathbb{C}$  for which uniform ensemble controllability can be expected are one dimensional  
 101 objects.

102 The following facts can be found in [11].

103 **Proposition 2.3.** *If (2.1) is uniformly ensemble controllable, then it is uniformly ensemble con-*  
 104 *trollable in any time  $T > 0$ , i.e., for any  $x^0, x^1 \in \mathcal{C}(\Theta)^n$ , any  $T > 0$  and any  $\varepsilon > 0$ , there exists a*  
 105 *control  $u \in L^2(0, T)^m$ , such that  $\sup_{\theta \in \Theta} |x(T, \theta; x^0, u) - x^1(\theta)| \leq \varepsilon$ .*

106 **Proposition 2.4.** *If (2.1) is uniformly ensemble controllable, then for every  $\theta \in \Theta$ , the pair*  
 107  *$(A(\theta), B(\theta))$  is controllable.*

108 **Remark 2.5.** The direct consequence of the last result is that the control operator can not vanish  
 109 in any point, i.e.  $B(\theta) \neq 0$  for every parameter  $\theta$ .

110 More generally, we have the following exact controllability property.

111 **Proposition 2.6.** *Assume (2.1) is uniformly ensemble controllable. For every  $K \in \mathbb{N}^*$  and every*  
 112  *$\theta_1, \dots, \theta_K \in \Theta$ , two by two distinct, the pair  $(\mathbb{A}, \mathbb{B})$  is controllable, where*

$$\mathbb{A} = \begin{pmatrix} A(\theta_1) & & \\ & \ddots & \\ & & A(\theta_K) \end{pmatrix} \in \mathbb{R}^{Kn \times Kn}, \quad \text{and} \quad \mathbb{B} = \begin{pmatrix} B(\theta_1) \\ \vdots \\ B(\theta_K) \end{pmatrix} \in \mathbb{R}^{Kn \times m}.$$

113 Let us also mention the unique continuation property for the adjoint system (cf. [10]). To this  
 114 end, we denote by  $\mathcal{M}(\Theta)$  the set of Radon measures supported in  $\Theta$ , and we define the *control to*  
 115 *final state operator*  $\Phi_T \in \mathcal{L}(L^2(0, T)^m, \mathcal{C}(\Theta)^n)$  by

$$(\Phi_T u)(\theta) = \int_0^T e^{(T-t)A(\theta)} B(\theta) u(t) dt \quad (\theta \in \Theta, u \in L^2(0, T)^m), \quad (2.3)$$

116 i.e.,  $(\Phi_T u)(\theta) = x(T, \theta, 0, u)$  with  $x$  given by (2.2). When there is no fear of ambiguity, we shall  
 117 use abbreviation  $x^T = \Phi_T u$ .

Note that the adjoint of  $\Phi_T$  is  $\Phi_T^* \in \mathcal{L}(\mathcal{M}(\Theta)^n, L^2(0, T)^m)$  defined by

$$(\Phi_T^* \mu)(t) = \int_{\Theta} B(\theta)^* e^{(T-t)A(\theta)^*} d\mu(\theta) = \left\langle e^{(T-t)A^*} \flat \mu, B \right\rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^{n \times m}} \quad (t \in (0, T), \mu \in \mathcal{M}(\Theta)^n). \quad (2.4)$$

118 **Proposition 2.7.** *The system (2.1) is uniformly ensemble controllable in time  $T > 0$  if and only if*

$$\left\{ \mu \in \mathcal{M}(\Theta)^n \mid \forall t \in [0, T], \int_{\Theta} B(\theta)^* e^{(T-t)A(\theta)^*} d\mu(\theta) = 0 \right\} = \{0\}. \quad (2.5)$$

119 By the last proposition the uniform ensemble controllability implies the unique continuation  
120 property of the adjoint operator, i.e.,

$$\Phi_T^* \mu = 0 \implies \mu = 0. \quad (2.6)$$

121 In particular, the Gramian operator defined as  $\Lambda_T := \Phi_T \Phi_T^* \in \mathcal{L}(\mathcal{M}(\Theta)^n; \mathcal{C}(\Theta)^n)$  is injective.

## 122 3 Minimal $L^2$ -norm controls

### 123 3.1 Optimality conditions

124 The aim of this section is to characterize the minimal  $L^2$ -norm control steering the system (2.1)  
125 from  $x^0$  to a distance  $\varepsilon > 0$  of  $x^1$  in a given time  $T > 0$ . More precisely, given  $T > 0$ ,  $\varepsilon > 0$  and  
126  $x^0, x^1 \in \mathcal{C}(\Theta)^n$ , we aim to find a minimizer of

$$\min \left\{ \frac{1}{2} \|u\|_{L^2(0,T)^m}^2, \right. \\ \left. u \in L^2(0,T)^m \quad \text{and} \quad \sup_{\theta \in \Theta} \left| y(\theta) - \int_0^T e^{(T-t)A(\theta)} B(\theta) u(t) dt \right| \leq \varepsilon. \right. \quad (P)$$

where  $y \in \mathcal{C}(\Theta)^n$  is the final target adjusted by the solution of the homogeneous equation, i.e.,

$$y(\theta) = x^1(\theta) - e^{TA(\theta)} x^0(\theta) \quad (\theta \in \Theta).$$

127 When there is no fear of ambiguity, we shall also refer to  $y$  as (adjusted) target.

We rewrite the problem (P) by introducing functions  $f \in \mathcal{C}(L^2(0,T)^m)$  and  $g : \mathcal{C}(\Theta)^n \rightarrow \mathbb{R}_+ \cup \{\infty\}$  defined as,

$$f(u) = \frac{1}{2} \|u\|_{L^2(0,T)^m}^2 \quad (u \in L^2(0,T)^m), \\ g(\xi) = \begin{cases} 0 & \text{if } \|\xi - y\|_{L^\infty(\Theta)^n} \leq \varepsilon, \\ \infty & \text{otherwise} \end{cases} \quad (\xi \in \mathcal{C}(\Theta)^n).$$

128 In particular,  $g$  is the indicator function of the ball  $\mathcal{B}_\infty(y, \varepsilon) \subset \mathcal{C}(\Theta)^n$ .

Hence, (P) can be expressed as

$$\min_{u \in L^2(0,T)^m} f(u) + g(\Phi_T u).$$

Note that  $f$  and  $g$  are convex functions, and our aim is to approach the optimization problem (P) by the Fenchel-Rockafellar duality. To this end, we consider the Fenchel conjugates of  $f$  and  $g$  defined as

$$f^*(v) = \sup_{u \in L^2(0,T)^m} \left( \langle v, u \rangle_{L^2(0,T)^m} - f(u) \right) = \frac{1}{2} \|v\|_{L^2(0,T)^m}^2 \quad (v \in L^2(0,T)^m), \\ g^*(\mu) = \sup_{\xi \in \mathcal{C}(\Theta)^n} \left( \langle \mu, \xi \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - g(\xi) \right) = \sup_{\substack{\xi \in \mathcal{C}(\Theta)^n \\ \|\xi - y\|_{L^\infty(\Theta)^n} \leq \varepsilon}} \langle \mu, \xi \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \\ = \langle \mu, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n} \quad (\mu \in \mathcal{M}(\Theta)^n).$$

129 Here, and in the rest of this paper, for  $\mu = (\mu_1, \dots, \mu_m) \in \mathcal{M}(\Theta)^n$ , we set

$$\|\mu\|_{\mathcal{M}(\Theta)^n} = \sup_{\substack{\xi \in \mathcal{C}(\Theta)^n \\ \|\xi\|_{L^\infty(\Theta)^n} \leq 1}} \langle \mu, \xi \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} = \sup_{\substack{\xi_1, \dots, \xi_n \in \mathcal{C}(\Theta) \\ \sup_{\Theta} \sum_{i=1}^n |\xi_i|^2 \leq 1}} \sum_{i=1}^n \langle \mu_i, \xi_i \rangle_{\mathcal{M}(\Theta), \mathcal{C}(\Theta)}.$$

130 Specially, if the components of  $\mu$  are written in the form  $\mu_i = f_i \flat \nu$  for some  $i$ -independent scalar  
 131 and positive measure  $\nu$ , then  $\|\mu\|_{\mathcal{M}(\Theta)^n} = \left\langle \left( \sqrt{\sum_i f_i^2} \right) \flat \nu, 1 \right\rangle_{\mathcal{M}(\Theta), \mathcal{C}(\Theta)} = \int_{\Theta} |f(\theta)| \, d\nu(\theta)$ . In the  
 132 special case of  $\mu$  being a combination of disjoint Dirac masses  $\mu = \sum_k m_k \delta_{\theta_k}$  for  $m_k \in \mathbb{R}^n$ , its  
 133 norm is given by  $\sum_k |m_k|$ .

134 **Theorem 3.1.** *Assume (2.1) is uniformly ensemble controllable. Then for every  $y \in \mathcal{C}(\Theta)^n$  and*  
 135 *every  $\varepsilon > 0$ , the minimization problem (P) admits a unique minimizer  $u_{opt} \in L^2(0, T)^m$  given by*  
 136 *the expression*

$$u_{opt} = \Phi_T^* \mu_{opt},$$

137 where  $\mu_{opt} \in \mathcal{M}(\Theta)^n$  minimizes the dual functional

$$J(\mu) = \frac{1}{2} \|\Phi_T^* \mu\|_{L^2(0, T)^m}^2 - \langle \mu, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n} \quad (\mu \in \mathcal{M}(\Theta)^n). \quad (3.1)$$

138 Furthermore, the minimizer  $\mu_{opt}$  is unique, and satisfies

$$\|u_{opt}\|_{L^2(0, T)^m}^2 = \langle \mu_{opt}, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon \|\mu_{opt}\|_{\mathcal{M}(\Theta)^n}, \quad (3.2)$$

139

$$\text{supp } \mu_{opt} \subset \{\theta \in \Theta \mid |(\Phi_T u_{opt})(\theta) - y(\theta)| = \varepsilon\} \quad (3.3)$$

140 and, for every test function  $v \in \mathcal{C}(\Theta)^n$  such that  $\Phi_T u_{opt} + v \in \mathcal{B}_\infty(y, \varepsilon)$ ,

$$\langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \geq 0. \quad (3.4)$$

141 Finally,  $\mu_{opt}$  is the zero measure if and only if  $y \leq \varepsilon$ .

142 **Remark 3.2.** As  $|(\Phi_T u_{opt})(\theta) - y(\theta)|$  is the distance of the optimal final state from the target  $x^1$ ,  
 143 the characterization (3.3) restricts the support of the optimal measure to the parameters  $\theta \in \Theta$   
 144 for which the optimal final state lies exactly on the boundary of the target ball  $\mathcal{B}_\infty(x^1, \varepsilon)$ .

*Proof.* We first observe that  $f$  is strictly convex, and  $\text{dom } g$  is not empty. Furthermore, we observe that  $\{u \in L^2(0, T)^m \mid \Phi_T u \in \text{dom } g\}$  is a convex set and  $g$  is lower semi-continuous. These facts ensure the existence and uniqueness of a minimizer of (P) (see e.g. [24, Theorem 2.19]). Note also that  $\text{cont } g = \text{int dom } g$  and  $\text{dom } f = L^2(0, T)^m$ , hence, the uniform ensemble controllability assumption guaranties that  $\Phi_T \text{dom } f \cap \text{cont } g \neq \emptyset$ . Thus, according to Fenchel duality (see e.g. [5, Theorem 4.4.3], [24, Theorem 3.51] or [12]), we have

$$\begin{aligned} \min_{u \in L^2(0, T)^m} f(u) + g(\Phi_T u) &= \max_{\mu \in \mathcal{M}(\Theta)^n} -f^*(\Phi_T^* \mu) - g^*(-\mu) = - \min_{\mu \in \mathcal{M}(\Theta)^n} f^*(\Phi_T^* \mu) + g^*(-\mu) \\ &= - \min_{\mu \in \mathcal{M}(\Theta)^n} \frac{1}{2} \|\Phi_T^* \mu\|_{L^2(0, T)^m}^2 - \langle \mu, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n}. \end{aligned}$$

145 In addition (3.1) admits a minimizer  $\mu_{opt} \in \mathcal{M}(\Theta)^n$ , and we have

$$f(u_{opt}) + g(\Phi_T u_{opt}) = -f^*(\Phi_T^* \mu_{opt}) - g^*(-\mu_{opt}) \quad (3.5)$$

146 and

$$\Phi_T^* \mu_{opt} \in \partial f(u_{opt}) \quad \text{and} \quad -\mu_{opt} \in \partial g(\Phi_T u_{opt}),$$

147 where  $\partial$  stands for the sub-differential. As  $f$  is a smooth function, its sub-differential is a singleton,  
 148 and from the first of the above inclusions we get that  $u_{opt} = \Phi_T^* \mu_{opt}$ .

149 Noticing that  $f^* = f$ ,  $g(\Phi_T u_{opt}) = 0$ , we get from (3.5),  $2f(u_{opt}) = -g^*(-\mu_{opt})$ , which leads  
 150 to (3.2).

151 We also have that  $x_{opt}^T := \Phi_T u_{opt} \in \mathcal{C}(\Theta)^n$  is such that  $\|x_{opt}^T - y\|_{L^\infty(\Theta)^n} \leq \varepsilon$ , i.e.,  
 152  $g(x_{opt}^T) = 0$ . Then,  $-\mu_{opt} \in \partial g(x_{opt}^T)$  implies that for every  $v \in \mathcal{C}(\Theta)^n$ ,  $g(x_{opt}^T + v) =$   
 153  $g(x_{opt}^T + v) - g(x_{opt}^T) \geq \langle -\mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ . In particular, if  $\|x_{opt}^T + v - y\|_{L^\infty(\Theta)^n} \leq \varepsilon$ , we  
 154 have  $0 \leq \langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ , thus proving (3.4).

155 From here we obtain the characterization (3.3) of the support of  $\mu_{opt}$ . Indeed, for every  
 156  $v \in \mathcal{C}(\Theta)^n$  such that  $\text{supp } v \subset \{\theta \in \Theta \mid |x_{opt}^T(\theta) - y(\theta)| < \varepsilon\}$ , there exists  $\alpha > 0$  such  
 157 that  $\|x_{opt}^T + \alpha v - y\|_{L^\infty(\Theta)^n} < \varepsilon$  and  $\|x_{opt}^T - \alpha v - y\|_{L^\infty(\Theta)^n} < \varepsilon$ . Hence, we shall have  $0 \leq$   
 158  $\langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$  and  $0 \leq -\langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ , leading to  $0 = \langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ .

159 Dealing with the uniqueness of  $\mu_{opt}$ , assume by contradiction that  $\mu_1$  and  $\mu_2$  are minimizer  
 160 of  $J$ . By the uniqueness of  $u_{opt}$ , we have  $u_{opt} = \Phi_T^* \mu_1 = \Phi_T^* \mu_2$ . According to Proposition 2.7, we  
 161 conclude that  $\mu_1 = \mu_2$ .

Finally, if  $\|y\|_{\mathcal{C}(\Theta)^n} \leq \varepsilon$  (meaning that the free dynamics brings the state within the target  
 ball) by the definition (3.1) of the functional  $J$  and the Cauchy-Schwarz inequality, we have that  
 for every measure  $\mu \in \mathcal{M}(\Theta)^n$

$$J(\mu) \geq -\|\mu\|_{\mathcal{M}(\Theta)^n} \|y\|_{\mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n} \geq 0 = J(0),$$

162 i.e., the functional  $J$  is nonnegative and  $\mu_{opt} = 0$  is its unique minimizer.

163 The reverse implication holds trivially, as if  $\|y\|_{\mathcal{C}(\Theta)^n} > \varepsilon$  then a nonzero control is required in  
 164 order to satisfy the constraint (of reaching the target ball). Using classical optimization techniques  
 165 arguments, it can be even shown that in that case  $\|\mu_{opt}\|_{\mathcal{M}(\Theta)^n}$  is bounded from below by a positive  
 166 constant depending linearly on the difference  $\|y\|_{\mathcal{C}(\Theta)^n} - \varepsilon$  (e.g. [17, Proposition 2.3]).  $\square$

167 **Proposition 3.3.** *Using the assumptions and notations introduced in Theorem 3.1, the optimal*  
 168 *measure  $\mu_{opt}$  is of the form*

$$d\mu_{opt}(\theta) = \alpha(\theta) (y(\theta) - x_{opt}^T(\theta)) d|\mu_{opt}|(\theta), \quad (3.6)$$

169 where  $|\mu_{opt}| = \sum_{i=1}^n |\mu_{opt,i}| \in \mathcal{M}(\Theta)$  is the sum of total variations of components of the vector  
 170 measure  $\mu_{opt}$ , while  $\alpha : \Theta \rightarrow \mathbb{R}_+$  is an integrable (with respect to  $|\mu_{opt}|$ ) nonnegative scalar function.

171 *Proof. Step 1.* We first show that

$$\langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \geq 0, \quad (3.7)$$

172 for every test function  $v \in \mathcal{C}(\Theta)^n$  such that  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle \geq 0$  whenever  $|(x_{opt}^T(\theta) - y(\theta))| =$   
 173  $\varepsilon$ ,  $\theta \in \Theta$ .

174 Suppose first that  $v$  satisfies the last inequality strictly, i.e., that  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle > 0$   
 175 for every  $\theta \in \Theta$  satisfying  $|x_{opt}^T(\theta) - y(\theta)| = \varepsilon$ . Then we show there exists a  $\theta$  independent  
 176 scalar  $\alpha > 0$  small enough such that  $\|x_{opt}^T + \alpha v - y\|_{L^\infty(\Theta)^n} \leq \varepsilon$ . Indeed, let us set  $\Theta_{opt} =$   
 177  $\{\theta \in \Theta \mid |x_{opt}^T(\theta) - y(\theta)| = \varepsilon\}$ , since  $\theta \mapsto |x_{opt}^T(\theta) - y(\theta)|$  is continuous, we have that  $\Theta_{opt}$  is  
 178 compact. On this set, we have  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle > 0$ , using the uniform continuity on  $\Theta$  of this  
 179 function, we get the existence of  $\delta > 0$  such that  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle > 2\delta$  for every  $\theta \in \Theta_{opt}$ ,  
 180 and there also exists  $\eta > 0$  such that  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle > \delta$  for every  $\theta \in \Theta_{opt} + B(0, \eta)$ . Using  
 181 the fact that  $\Theta \setminus (\Theta_{opt} + B(0, \eta))$  is compact, we also obtain the existence of  $\tilde{\varepsilon} < \varepsilon$  such that  
 182  $|x_{opt}^T(\theta) - y(\theta)| \leq \tilde{\varepsilon}$  for every  $\theta \in \Theta \setminus (\Theta_{opt} + B(0, \eta))$ . All in all, we have for every  $\alpha > 0$ ,



- for  $\theta \in \Theta_{opt} + B(0, \eta)$ ,

$$\begin{aligned} |x_{opt}(\theta) + \alpha v(\theta) - y(\theta)|^2 &= |x_{opt}(\theta) - y(\theta)|^2 - 2\alpha \langle v(\theta), y(\theta) - x_{opt}(\theta) \rangle + \alpha^2 |v(\theta)|^2 \\ &\leq \varepsilon^2 - 2\alpha\delta + \alpha^2 \|v\|_{L^\infty(\Theta)}^2. \end{aligned}$$

- for  $\theta \in \Theta \setminus (\Theta_{opt} + B(0, \eta))$ ,

$$|x_{opt}^T(\theta) + \alpha v(\theta) - y(\theta)|^2 \leq \tilde{\varepsilon}^2 + 2\alpha\tilde{\varepsilon} \|v\|_{L^\infty(\Theta)} + \alpha^2 \|v\|_{L^\infty(\Theta)}^2.$$

183 Hence, taking  $\alpha > 0$  small enough, we have  $|x_{opt}^T(\theta) + \alpha v(\theta) - y(\theta)| \leq \varepsilon$  for every  $\theta \in \Theta$ . By (3.4)  
184 this ensures that  $\langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \geq 0$  for every  $v \in \mathcal{C}(\Theta)^n$  such that  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle > 0$   
185 for every  $\theta \in \Theta_{opt}$ .

186 It remains to prove that (3.7) holds true, even if  $\langle v(\theta), y(\theta) - x_{opt}^T(\theta) \rangle \geq 0$  for every  $\theta \in \Theta_{opt}$ . In  
187 fact, if  $v$  satisfies the last condition, then  $\langle v(\theta) + \alpha(y(\theta) - x_{opt}^T(\theta)), y(\theta) - x_{opt}^T(\theta) \rangle > 0$  for every  
188  $\theta \in \Theta_{opt}$  and every  $\alpha > 0$ . This ensures that  $0 \leq \langle \mu_{opt}, v(\theta) + \alpha(y(\theta) - x_{opt}^T(\theta)) \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ , and  
189 hence, taking the limit  $\alpha \rightarrow 0$ , we obtain the desired result.

190 **Step 2.** Now we prove the relation (3.6)

191 First, let us note that  $\mu_{opt}$  is absolutely continuous with respect to the measure  $|\mu_{opt}|$ . By the  
192 Lebesgue-Radon-Nikodym theorem (see e.g. [14]) there exists an integrable (with respect to  $|\mu_{opt}|$ )  
193 vector function  $\varphi$  such that  $d\mu_{opt}(\theta) = \varphi(\theta) d|\mu_{opt}|(\theta)$ .

194 It remains to prove that  $\varphi = \alpha(y - x_{opt}^T)$  for some  $\alpha : \Theta \rightarrow \mathbb{R}_+$  which is integrable (with respect  
195 to  $|\mu_{opt}|$ ). To this end, without loosing generality, let us write  $\varphi$  as

$$\varphi(\theta) = \alpha(\theta)(y(\theta) - x_{opt}^T(\theta)) + w(\theta), \quad (3.8)$$

196 where  $w(\theta) \in \{y(\theta) - x_{opt}^T(\theta)\}^\perp$ , while  $\alpha$  is an arbitrary (not necessarily non-negative) scalar  
197 function. Without loss of generality, we can assume that  $\text{supp } \varphi \subset \text{supp } \mu_{opt}$ , i.e., we can assume  
198 that  $\alpha$  and  $w$  vanish on  $\Theta \setminus \text{supp } \mu_{opt}$ . Taking into account that  $y - x_{opt}^T$  is continuous on  $\Theta$  and that  
199  $|y - x_{opt}^T| = \varepsilon$  on  $\text{supp } \mu_{opt}$ , we get that  $\alpha(\theta) = \langle y(\theta) - x_{opt}^T(\theta), \varphi(\theta) \rangle / \varepsilon^2$  for every  $\theta \in \text{supp } \mu_{opt}$ .  
200 This ensures that  $\alpha$  is  $|\mu_{opt}|$ -measurable, and then  $w = \varphi - \alpha(y - x_{opt}^T)$  is also  $|\mu_{opt}|$ -measurable.

Define  $v = \psi(y - x_{opt}^T)$ , where  $\psi \in \mathcal{C}(\Theta)$  is an arbitrary non-negative, scalar test function.  $v$  is therefore an eligible test function in (3.7), implying

$$\begin{aligned} 0 \leq \langle \mu_{opt}, v \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} &= \int_{\Theta} \langle (\alpha(\theta)(y(\theta) - x_{opt}^T(\theta)) + w(\theta)), \psi(\theta)(y(\theta) - x_{opt}^T(\theta)) \rangle d|\mu_{opt}|(\theta) \\ &= \int_{\Theta} \psi(\theta) |y(\theta) - x_{opt}^T(\theta)|^2 \alpha(\theta) d|\mu_{opt}|(\theta) = \left\langle \alpha \mathbb{1}_{\text{supp } \mu_{opt}}, \psi |y - x_{opt}^T|^2 \right\rangle_{\mathcal{M}(\Theta), \mathcal{C}(\Theta)}. \end{aligned}$$

201 As  $|\mu_{opt}|$  is a non-negative measure, by varying  $\psi$  and taking again into account that  $|y - x_{opt}^T| = \varepsilon$   
202 on  $\text{supp } \mu_{opt}$ , it follows  $\alpha \geq 0$  for  $|\mu_{opt}|$ -a.e.  $\theta$ . Taking  $\psi = 1$ , we also get that  $\alpha$  is  $|\mu_{opt}|$ -integrable.

Furthermore, (3.2) together with  $\|u_{opt}\|_{L^2(0,T)^m}^2 = \|\Phi_T^* \mu_{opt}\|_{L^2(0,T)^m}^2 = \langle \mu_{opt}, x_{opt}^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$   
leads to

$$\varepsilon \|\mu_{opt}\|_{\mathcal{M}(\Theta)^n} = \langle \mu_{opt}, y - x_{opt}^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}.$$

At this stage we exploit the form (3.8) and put  $\mu_{opt} = (\alpha(y - x_{opt}^T) + w) \llcorner |\mu_{opt}|$  in the last  
expression. By taking into account that  $w(\theta)$  is orthogonal to  $y(\theta) - x_{opt}^T(\theta)$ , and recalling that

$|y - x_{opt}^T| = \varepsilon$  on  $\text{supp } \mu_{opt}$ , we get

$$\begin{aligned} \varepsilon \int_{\Theta} |\alpha(\theta)(y(\theta) - x_{opt}^T(\theta)) + w(\theta)| \, d|\mu_{opt}|(\theta) &= \int_{\Theta} \alpha(\theta) |(y(\theta) - x_{opt}^T(\theta))|^2 \, d|\mu_{opt}|(\theta) \\ &= \varepsilon \int_{\Theta} \alpha(\theta) |(y(\theta) - x_{opt}^T(\theta))| \, d|\mu_{opt}|(\theta) \end{aligned} \quad (3.9)$$

203 As  $\alpha$  is a non-negative function and  $|\alpha(y - x_{opt}^T) + w| \geq \alpha |y - x_{opt}^T|$ , the equality (3.9) implies  
 204  $|\alpha(y - x_{opt}^T) + w| = \alpha |y - x_{opt}^T|$  for  $|\mu_{opt}|$ -a.e.  $\theta$ , that is to say that  $w = 0$  for  $|\mu_{opt}|$  (and conse-  
 205 quently for  $\mu_{opt}$ ) a.e.  $\theta$ .  $\square$

206 **Remark 3.4.** The last result covers the well-known behavior of the optimal control in the deter-  
 207 ministic case, i.e., when  $\Theta$  is a singleton. In that case, unless the optimal control is trivial, the  
 208 solution of the dual problem is a vector having the same direction as  $y - x_{opt}^T$  (see e.g. [19]).  
 209 In general, when a functional of the form (3.1) is defined on a Hilbert space, its minimizer  $h_{opt}$  is  
 210 either trivial or a solution to the Euler-Lagrange equation which formally reads as:

$$y - x_{opt}^T = \varepsilon \frac{h_{opt}}{\|h_{opt}\|}.$$

211 In that case it follows directly that the minimizer has the same direction as  $y - x_{opt}^T$ . This kind of  
 212 result is stronger than the one obtained in the last proposition, and it corresponds to a constant  
 213 function  $\alpha$  in (3.6). The reason why such approach is not possible in our setting is that the norm  
 214 term  $\|\mu\|_{\mathcal{M}(\Theta)^n}$  entering the functional  $J$  is not differentiable.

215 As a consequence of the last proposition, we obtain the following characterization of the optimal  
 216 measure in the scalar case.

217 **Corollary 3.5.** *Using the assumptions and notations introduced in Theorem 3.1, let us suppose*  
 218  $n = m = 1$ .

219 *Then the control operator does not change sign, i.e., there exists  $\epsilon \in \{-1, 1\}$  and  $\beta \in \mathcal{C}(\Theta)$  such*  
 220 *that  $\beta(\theta) > 0$  and  $B(\theta) = \epsilon\beta(\theta)$  for every  $\theta \in \Theta$ .*

221 *Set  $\Theta_+ = \{\theta \in \Theta \mid y(\theta) - x_{opt}^T(\theta) = \varepsilon\}$  and  $\Theta_- = \{\theta \in \Theta \mid y(\theta) - x_{opt}^T(\theta) = -\varepsilon\}$ . Then*  
 222  *$\text{supp } \mu_{opt} \subset \Theta_- \cup \Theta_+$  and  $\mu_{opt}$  is nonnegative (respectively, nonpositive) on  $\Theta_+$  (respectively  $\Theta_-$ ).*

223 *In addition, if  $\Theta_- = \emptyset$  (respectively  $\Theta_+ = \emptyset$ ), then either  $u_{opt} \equiv 0$  or  $\text{sign}(u_{opt}(t)) = \epsilon$*   
 224 *(respectively  $\text{sign}(u_{opt}(t)) = -\epsilon$ ) for every  $t \in [0, T]$ .*

225 *Proof.* The fact that  $B \in \mathcal{C}(\Theta)$  does not vanish directly follows from Proposition 2.4. Being a  
 226 continuous function, its sign is constant.

227 The support and sign conditions of  $\mu_{opt}$  directly follow from (3.3) and Proposition 3.3, respectively.  
 228 For the sign of  $u_{opt}$ , let us assume that  $\Theta_- = \emptyset$  (the proof for other case is identical). We have for  
 229 every  $t \in [0, T]$ ,

$$u_{opt}(t) = \int_{\Theta} e^{(T-t)A^*(\Theta)} B(\theta) \, d\mu_{opt}(\theta) = \epsilon \int_{\Theta_+} e^{(T-t)A^*(\Theta)} \beta(\theta) \, d\mu_{opt}(\theta),$$

230 We conclude by noticing that, for every  $\theta \in \Theta$ ,  $e^{(T-t)A^*(\Theta)} \beta(\theta) > 0$  and  $\mu_{opt}$  is nonnegative. In  
 231 particular,  $u_{opt} \equiv 0$  if and only if  $\mu_{opt} = 0$  and, according to Theorem 3.1, this holds if and only if  
 232  $\|y\|_{\mathcal{C}(\Theta)^n} \leq \varepsilon$ . Otherwise, the optimal control has a constant sign equal to  $\epsilon$  for every  $t \in [0, T]$ .  $\square$

233 The rest of this subsection provides results by which one can construct a family of optimization  
 234 problem (P) for which the minimizers of (3.1) can be explicitly calculated. This will allow us to

235 compare the results obtained by the numerical procedure to be presented in the Section 6 with the  
 236 exact ones and to verify efficiency and correctness of the proposed algorithms (see Section 7).

237 The idea is to invert the problem: we specify a positive measure  $\tilde{\mu}$  and calculate  $\tilde{x}^T = \Phi_T \Phi_T^* \tilde{\mu}$ ,  
 238 the final state of the system run by the control  $\tilde{u} = \Phi_T^* \tilde{\mu}$ . The adjusted target  $y$  is then carefully  
 239 chosen as a center of a square whose one vertex is  $\tilde{x}^T$  (note that the geometrical representation of  
 240 a ball in  $\mathcal{C}(\Theta)^n$  is actually a square). In particular, in accordance to Proposition 3.3, the difference  
 241  $y - \tilde{x}^T$  has the same direction as  $\tilde{\mu}$  for every parameter value  $\theta$ .

242 **Proposition 3.6.** *Using the assumptions and notations introduced in Theorem 3.1 take  $\varepsilon > 0$ ,*  
 243  *$\eta \in \mathcal{M}(\Theta)$  a nonnegative scalar measure, and  $f \in \mathcal{C}(\Theta)^n$  such that  $|f|$  does not vanish. Define*  
 244  *$\tilde{\mu} = f \flat \eta$ ,  $\tilde{x}^T = \Phi_T \Phi_T^* \tilde{\mu}$  and  $y = \tilde{x}^T + \varepsilon \frac{f}{|f|}$ . Then  $\tilde{\mu}$  is the minimizer of  $J$  defined by (3.1).*

*Proof.* Let us first observe that for every  $\mu \in \mathcal{M}(\Theta)^n$ , we have

$$\begin{aligned} J(\mu) &= \frac{1}{2} \|\Phi_T^* \mu\|_{L^2(0,T)^m}^2 - \langle \mu, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n} \\ &= \frac{1}{2} \|\Phi_T^* \mu\|_{L^2(0,T)^m}^2 - \langle \Phi_T^* \mu, \Phi_T^* \tilde{\mu} \rangle_{L^2(0,T)^m} + \varepsilon \left( \|\mu\|_{\mathcal{M}(\Theta)^n} - \left\langle \mu, \frac{f}{|f|} \right\rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \right) \end{aligned}$$

and

$$J(\tilde{\mu}) = -\frac{1}{2} \|\Phi_T^* \tilde{\mu}\|_{L^2(0,T)^m}^2 + \varepsilon \left( \int_{\Theta} |f| \, d\eta - \int_{\Theta} \frac{\langle f, f \rangle}{|f|} \, d\eta \right) = -\frac{1}{2} \|\Phi_T^* \tilde{\mu}\|_{L^2(0,T)^m}^2.$$

245 Let  $\mu_{opt} \in \mathcal{M}(\Theta)^n$  be the minimizer of  $J$ , we have  $J(\mu_{opt}) \leq J(\tilde{\mu})$ , that is to say that

$$\frac{1}{2} \|\Phi_T^* (\mu_{opt} - \tilde{\mu})\|_{L^2(0,T)^m}^2 \leq -\varepsilon \left( \|\mu_{opt}\|_{\mathcal{M}(\Theta)^n} - \left\langle \mu_{opt}, \frac{f}{|f|} \right\rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \right).$$

246 Obviously, we have  $\|\mu_{opt}\|_{\mathcal{M}(\Theta)^n} \geq \left\langle \mu_{opt}, \frac{f}{|f|} \right\rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ , from which we can conclude that we  
 247 necessarily have  $\Phi_T^* (\mu_{opt} - \tilde{\mu}) = 0$ . We conclude that  $\mu_{opt} = \tilde{\mu}$  using the unique continuation  
 248 property (2.6) of the adjoint.  $\square$

249 As a direct corollary, we have the following result.

250 **Corollary 3.7.** *Using the assumptions and notations introduced in Theorem 3.1. Suppose  $n = 1$ ,*  
 251 *let  $\varepsilon > 0$ ,  $\epsilon \in \{-1, 1\}$ ,  $\eta \in \mathcal{M}(\Theta)$  a nonnegative measure,  $y = \epsilon(\Phi_T \Phi_T^* \eta + \varepsilon)$ . The minimizer*  
 252 *of (3.1) is  $\epsilon \eta$ .*

253 *Proof.* This is Proposition 3.6 with  $f = \epsilon$ .  $\square$

254 From Proposition 3.6 and Corollary 3.7, we see that there is no reason that the optimal mea-  
 255 sure  $\mu_{opt}$  is sparse, i.e., the Lebesgue measure of its support is in general not zero. This might look  
 256 surprisingly at the first glance, as  $\mu_{opt}$  is a minimizer of the functional containing the  $\|\cdot\|_{\mathcal{M}(\Theta)^n}$   
 257 term. This non-sparsity pitfall will be discussed in Section 4 in the case where the parameter set  
 258 is of a finite cardinality.

## 259 3.2 Continuity properties of the minimizers

260 The characterization of the solution through the adjoint problem, that is obtained in the previous  
 261 subsection, allows us also to demonstrate the continuity of solutions to optimization problem (P)  
 262 with respect to the tolerance  $\varepsilon$ , the time  $T$  and the adjusted target  $y$ . More precisely, we consider  
 263 a sequence of functionals

$$J_k(\mu) = \frac{1}{2} \|\Phi_{T_k}^* \mu\|_{L^2(0,T)^m}^2 - \langle \mu, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon_k \|\mu\|_{\mathcal{M}(\Theta)^n} \quad (\mu \in \mathcal{M}(\Theta)^n). \quad (3.10)$$

264 where:

- 265 (i)  $y_k \rightarrow y$  strongly in  $\mathcal{C}(\Theta)^n$ ,  
 266 (ii)  $T_k \rightarrow T$  in  $\mathbb{R}_+^*$   
 267 (iii)  $\varepsilon_k \rightarrow \varepsilon$  in  $\mathbb{R}_+^*$ .

268 By Theorem 3.1, for each  $k \in \mathbb{N}$ , the functional  $J_k$  admits the unique minimizer  $\mu_k$  satisfying  
 269 the relation

$$\|\Phi_{T_k}^* \mu_k\|_{L^2(0, T_k)^m}^2 = \langle \mu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon_k \|\mu_k\|_{\mathcal{M}(\Theta)^n} \quad (3.11)$$

270 and the control  $u_k = \Phi_{T_k}^* \mu_k$  is the solution of the optimal ensemble control problem

$$\min_{u \in L^2(0, T)^m} \frac{1}{2} \|u\|_{L^2(0, T)^m}^2, \quad \text{and} \quad \|y_k - \Phi_{T_k} u\|_{L^\infty(\Theta)^n} \leq \varepsilon_k. \quad (P_k)$$

271 We are going to prove that  $(\mu_k)_k$  and  $(u_k)_k$  converge respectively vaguely and strongly (in the  
 272 appropriate spaces) to the limit points  $\mu$  and  $u$ , respectively, where  $\mu$  is the minimizer of the limit  
 273 functional (3.1) and  $u$  is the solution of the ensemble optimal control problem (P).

274 Note that the convergence of time horizons  $T_k$  implies strong convergence of the operators  $\Phi_{T_k}$   
 275 and  $\Phi_{T_k}^*$ . The following technical result characterizes strong convergence of arbitrary operators on  
 276 Banach spaces.

277 **Lemma 3.8.** *Let  $(L_k)_k$  be a sequence of linear operators in  $\mathcal{L}(X, Y)$ , with  $X$  and  $Y$  being Banach  
 278 spaces. The following statements are equivalent.*

279 (a)  $L_k \rightarrow L$  strongly in  $\mathcal{L}(X, Y)$ , i.e.,

$$(\forall u \in X) \quad L_k u \rightarrow Lu \quad \text{in } Y.$$

280 (b) For every weakly  $*$  converging sequence  $(v_k)_k$  in  $Y'$  it holds

$$v_k \xrightarrow{*} v \implies L_k^* v_k \xrightarrow{*} L^* v \quad \text{in } X'.$$

281 (c) For every strongly converging sequence  $(u_k)_k$  in  $X$  it holds

$$u_k \rightarrow u \implies L_k u_k \rightarrow Lu \quad \text{in } Y.$$

282 *Proof.* (a)  $\Rightarrow$  (b): Let  $(v_k)_k$  converge weakly  $*$  to  $v \in Y'$ . By employing the strong convergence of  
 283 the operators  $L_k$ , for every  $u \in X$  we get

$$\langle L_k^* v_k, u \rangle_{X', X} = \langle v_k, L_k u \rangle_{Y', Y} \rightarrow \langle v, Lu \rangle_{Y', Y} = \langle L^* v, u \rangle_{X', X},$$

284 i.e.,  $L_k^* v_k \xrightarrow{*} L^* v$  in  $X'$ .

285

286 (b)  $\Rightarrow$  (c): We first prove the weak convergence of the sequence  $(L_k u_k)_k$ . To this end, note that  
 287 for an arbitrary  $v \in Y'$  we have

$$\langle v, L_k u_k \rangle_{Y', Y} = \langle L_k^* v, u_k \rangle_{X', X} \rightarrow \langle L^* v, u \rangle_{X', X} = \langle v, Lu \rangle_{Y', Y}.$$

288 Here we have employed the strong convergence of  $(u_k)$  and the assumption (b). In particular, we  
 289 have

$$\|Lu\|_Y \leq \liminf_{k \rightarrow \infty} \|L_k u_k\|_Y. \quad (3.12)$$

290 On the other hand, for each  $k$  there exists  $v_k \in Y'$  such that

$$\|L_k u_k\|_Y^2 = \langle v_k, L_k u_k \rangle_{Y', Y}, \quad (3.13)$$

291 and  $\|v_k\|_{Y'} = \|L_k u_k\|_Y$  (see e.g. [6, Corollary 1.3.]).

292 Due to the weak convergence of  $(L_k u_k)_k$ , the sequence of norms  $\|L_k u_k\|_Y$  is bounded, in sequel  
 293 implying boundedness of  $(v_k)_k$  in  $Y'$ . In particular, up to a subsequence,  $(v_k)_k$  converge weakly \*  
 294 to some  $v \in Y'$ , and it holds

$$\|v\|_{Y'} \leq \liminf_{k \rightarrow \infty} \|v_k\|_{Y'} = \liminf_{k \rightarrow \infty} \|L_k u_k\|_Y. \quad (3.14)$$

295 From here it follows

$$\|L_k u_k\|_Y^2 = \langle v_k, L_k u_k \rangle_{Y', Y} = \langle L_k^* v_k, u_k \rangle_{X', X} \longrightarrow \langle L^* v, u \rangle_{X', X} = \langle v, Lu \rangle_{Y', Y}, \quad (3.15)$$

296 where we passed to the limit by using the assumption (b) again.

In particular, by employing (3.15) and (3.14), we obtain

$$\lim_{k \rightarrow \infty} \|L_k u_k\|_Y^2 = Y' \langle v, Lu \rangle_Y \leq \|v\|_{Y'} \|Lu\|_Y \leq \lim_{k \rightarrow \infty} \|L_k u_k\|_Y \|Lu\|_Y,$$

297 implying  $\lim_{k \rightarrow \infty} \|L_k u_k\|_Y \leq \|Lu\|_Y$ . Together with (3.12) this finishes the proof.

298

299 (c)  $\Rightarrow$  (a): Trivial (take a constant sequence  $u_k = u$ ). □

300 Before proving the convergence, we first demonstrate the boundedness properties of the mini-  
 301 mizers.

302 **Lemma 3.9.** *The sequence  $(\mu_k)_k$  of minimizers of functionals  $J_k$  given by (3.10) is bounded*  
 303 *in  $\mathcal{M}(\Theta)^n$ .*

304 *Proof.* We argue by contradiction and assume that

$$\|\mu_k\|_{\mathcal{M}(\Theta)^n} \longrightarrow \infty \quad (3.16)$$

305 (up to a non-relabelled subsequence).

306 Dividing the relation (3.11) by  $\|\mu_k\|_{\mathcal{M}(\Theta)^n}$  and denoting  $\nu_k = \mu_k / \|\mu_k\|_{\mathcal{M}(\Theta)^n}$  we get

$$\|\mu_k\|_{\mathcal{M}(\Theta)^n} \|\Phi_{T_k}^* \nu_k\|_{L^2(0, T)^m}^2 = \langle \nu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon_k. \quad (3.17)$$

307 As all the terms on the right-hand side are bounded (with respect to  $k$ ), the contradictory assump-  
 308 tion (3.16) implies

$$\Phi_{T_k}^* \nu_k \rightarrow 0 \quad \text{strongly in } L^2(0, T)^m. \quad (3.18)$$

309 On the other hand, denoting by  $\nu$  the vague limit of  $(\nu_k)$  (up to a subsequence), the strong  
 310 convergence of operators  $\Phi_{T_k}$  and Lemma 3.8 imply  $\Phi_{T_k}^* \nu_k \rightharpoonup \Phi_T^* \nu$  weakly in  $L^2(0, T)^m$ . Together  
 311 with (3.18), we get  $\Phi_T^* \nu = 0$ , while the ensemble controllability assumption (2.5) implies  $\nu = 0$ .

312 Going back to (3.17) and passing to the limit we get

$$0 > \varepsilon_k - \langle \nu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \longrightarrow \varepsilon.$$

313 Here we use that  $\langle \nu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$  is the dual product of a strongly convergent functions with  
 314 measures converging vaguely to zero, thus vanishing on the limit.

315 The last relation provides the desired contradiction which, in turn, implies the result. □

316 Based on the last two lemmas, we are going to obtain the main result of this paragraph.

317 **Proposition 3.10.** *The sequence of minimizers  $\mu_k$  of the functionals  $J_k$  given by (3.10) converges*  
 318 *vaguely in  $\mathcal{M}(\Theta)^n$  to  $\mu_{opt}$  (the minimizer of the limit functional (3.1)), and  $\|\mu_k\|_{\mathcal{M}(\Theta)^n}$  converges*  
 319 *to  $\|\mu_{opt}\|_{\mathcal{M}(\Theta)^n}$ . In addition, the sequence of optimal ensemble controls  $u_k = \Phi_{T_k}^* \mu_k$  converges*  
 320 *strongly in  $L^2(0, T)^m$  to the solution of the limit problem (P).*

321 *Proof.* By Lemma 3.9, there exists a (non-relabelled) subsequence of the minimizers  $\mu_k$  converging  
 322 vaguely to some measure  $\mu_L \in \mathcal{M}(\Theta)^n$ .

323 In the first part of the proof, we show that the limit  $\mu_L$  is the unique minimizer of the limit  
 324 functional (3.1). Thus, it suffices to show

$$J(\mu_L) \leq J(\mu), \quad \mu \in \mathcal{M}(\Theta)^n. \quad (3.19)$$

325 To this end, we explore the convergence properties of the sequence of minimal functional values

$$\liminf_{k \rightarrow \infty} J_k(\mu_k) = \liminf_{k \rightarrow \infty} \left( \frac{1}{2} \|\Phi_{T_k}^* \mu_k\|_{L^2(0, T)^m}^2 - \langle \mu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon_k \|\mu_k\|_{\mathcal{M}(\Theta)^n} \right). \quad (3.20)$$

326 Due to the weak lower semi-continuity of norms in Banach spaces, the last term is bounded from  
 327 below by  $\varepsilon \|\mu_L\|_{\mathcal{M}(\Theta)^n}$ .

328 The second term on the right-hand side of (3.20) is the product of a strongly and vaguely  
 329 convergent sequence, thus  $\langle \mu_k, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \rightarrow \langle \mu_L, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$ .

330 As for the first term entering functional  $J_k$  by Lemma 3.8 and the strong convergence of the  
 331 operators  $\Phi_{T_k}$ , we have  $\Phi_{T_k}^* \mu_k \rightharpoonup \Phi_T^* \mu_L$  weakly in  $L^2(0, T)^n$ , implying

$$\liminf_{k \rightarrow \infty} \|\Phi_{T_k}^* \mu_k\|_{L^2(0, T)^m}^2 \geq \|\Phi_T^* \mu_L\|_{L^2(0, T)^m}^2. \quad (3.21)$$

Consequently, from (3.20) we get

$$\liminf_{k \rightarrow \infty} J_k(\mu_k) \geq \frac{1}{2} \|\Phi_T^* \mu_L\|_{L^2(0, T)^m}^2 - \langle \mu_L, y \rangle + \varepsilon \|\mu_L\|_{\mathcal{M}(\Theta)^n} = J(\mu_L)$$

332 As  $\mu_k$  minimizes  $J_k$ , for an arbitrary  $\mu \in \mathcal{M}(\Theta)^n$  we have

$$\liminf_{k \rightarrow \infty} J_k(\mu) \geq J(\mu_L). \quad (3.22)$$

333 On the other hand, employing the assumed convergences, (i), (ii) and (iii), for each fixed  $\mu$  we have

$$\begin{aligned} J_k(\mu) &= \frac{1}{2} \|\Phi_{T_k}^* \mu\|_{L^2(0, T)^m}^2 - \langle \mu, y_k \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon_k \|\mu\|_{\mathcal{M}(\Theta)^n} \\ &\rightarrow \frac{1}{2} \|\Phi_T^* \mu\|_{L^2(0, T)^m}^2 - \langle \mu, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} + \varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n} = J(\mu). \end{aligned} \quad (3.23)$$

334 Together with (3.22) this implies (3.19), and in particular  $\mu_k \xrightarrow{*} \mu_{opt}$ .

335 In the second part of the proof, we will show that the sequence  $\|\mu_k\|_{\mathcal{M}(\Theta)^n}$  converges  
 336 to  $\|\mu_{opt}\|_{\mathcal{M}(\Theta)^n}$ .

337 Taking  $\mu = \mu_{opt}$  in (3.23) we have,

$$J(\mu_{opt}) = \lim_{k \rightarrow \infty} J_k(\mu_{opt}) \geq \lim_{k \rightarrow \infty} J_k(\mu_k).$$

338 Combining the obtained inequality with (3.22) (remember that  $\mu_{opt} = \mu_L$ ) we get the convergence  
 339 of the optimal values  $J_k(\mu_k) \rightarrow J(\mu_{opt})$ . Together with (3.21) this convergence implies,

$$\liminf_{k \rightarrow \infty} \|\mu_k\|_{\mathcal{M}(\Theta)^n} \leq \|\mu_{opt}\|_{\mathcal{M}(\Theta)^n}.$$

340 which together with the vague convergence of the sequence  $(\mu_k)$  implies the desired result.

341 The strong convergence of optimal ensemble controls  $u_k = \Phi_{T_k}^* \mu_k$  follows now directly from the  
 342 strong convergence of the operators  $\Phi_{T_k}$  and the vague convergence of the measures  $\mu_k$  (cf. Lem-  
 343 ma 3.8 (c)).  $\square$

## 344 4 Systems with a finite number of parameter values

345 In this section, we restrict the analysis to the case of a finite number of parameters, i.e.,  $\text{card } \Theta < \infty$ .  
 346 Such situation deserves a special attention, not only because in some applications the parameter  
 347 assumes only discrete and bounded values, but also because the most general situations (i.e., those  
 348 in which the parameter ranges over a continuous set of values) are numerically treated and analyzed  
 349 as the limit case of discretized problems as the number of elements in the parameter set goes to  
 350 infinity.

351 The aim is to provide a geometrical description of the problem and its solution, which will  
 352 allow us to discuss the conditions under which the optimal measure (i.e., the minimizer of the  
 353 functional  $J$  given by (3.1)) is sparse.

354 To this end, we assume that the parameter dependent system (2.1) is uniformly ensemble  
 355 controllable, and that we deal with a parameter set  $\Theta$  of finite cardinality. As we consider finite  
 356 dimensional systems only, these assumptions imply that the Gramian operator  $\Lambda_T = \Phi_T \Phi_T^* \in$   
 357  $\mathcal{L}(\mathcal{M}(\Theta)^n; \mathcal{C}(\Theta)^n)$  is not just injective (due to Proposition 2.7), but also of full range. Indeed,  
 358 due to the assumed finite cardinality of the parameter set, both spaces  $\mathcal{M}(\Theta)^n$  and  $\mathcal{C}(\Theta)^n$  are  
 359 finite dimensional and are isomorphic to  $\mathbb{R}^{nK}$  (where  $K = \text{card } \Theta \in \mathbb{N}^*$  denotes the number of  
 360 parameters), thus the statement follows from the rank-nullity theorem.

For  $c \geq 0$ , we introduce the subset  $E_c \subset \mathcal{C}(\Theta)^n$  consisting of all the final states  $x^T$  that can be reached with a control of norm less or equal to  $c$ , i.e.,

$$E_c := \left\{ x^T \in \mathcal{C}(\Theta)^n \mid \exists u \in L^2(0, T)^m \text{ such that } \|u\|_{L^2(0, T)^m} \leq c \text{ and } x^T = \Phi_T u \right\}.$$

As the minimal norm control reaching any state  $x^T$  is of the HUM form  $\Phi_T^* \mu$  for some  $\mu \in \mathcal{M}(\Theta)$  (see e.g. [29]), the introduced set can be characterized as follows

$$E_c = \left\{ x^T \in \mathcal{C}(\Theta)^n \mid \exists \mu \in \mathcal{M}(\Theta)^n \text{ such that } x^T = \Lambda_T \mu \text{ and } \|\Phi_T^* \mu\|_{L^2(0, T)^m} \leq c \right\},$$

where  $\Lambda_T$  stands for the Gramian  $\Phi_T \Phi_T^*$ . As the ensemble controllability assumption ensures the injectivity of the Gramian, the optimal control norm can be rewritten as

$$\|\Phi_T^* \mu\|_{L^2(0, T)^m}^2 = \langle \mu, \Phi_T \Phi_T^* \mu \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} = \langle \Lambda_T^{-1} x^T, x^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n},$$

361 leading to the following characterization of the set  $E_c$ :

$$E_c = \left\{ x^T \in \mathcal{C}(\Theta)^n \mid \langle \Lambda_T^{-1} x^T, x^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \leq c^2 \right\}. \quad (4.1)$$

362 In other words,  $E_c$  is a sublevel set of a smooth function  $F(x^T) = \langle \Lambda_T^{-1} x^T, x^T \rangle$ . In particular,  
 363  $\{E_c, c \geq 0\}$  is a nested family of closed, convex sets with smooth boundary  $\partial E_c = \{x^T \mid F(x^T) =$   
 364  $c^2\}$ . The normal functional at the point  $x^T \in \partial E_c$  is  $n_{E_c}(x^T) = \nabla F(x^T) = 2\Lambda_T^{-1} x^T$ .

365 **Theorem 4.1.** *Let  $x^T, x^1 \in \mathcal{C}(\Theta)^n$  be such that  $|x^1(\theta) - x^T(\theta)| \leq \varepsilon$  for every  $\theta \in \Theta$ .  
 366 Then  $x^T$  is the optimal final state of the problem (P) with the target  $x^1$  and initial datum  $x^0 = 0$   
 367 if and only if the normal functional  $n_{E_c}(x^T)$  is of the form*

$$n_{E_c}(x^T) = (\alpha(x^1 - x^T)) \flat \nu, \quad (4.2)$$

368 where  $\alpha \in \mathcal{C}(\Theta)$  and  $\nu \in \mathcal{M}(\Theta)$  are respectively a non-negative scalar function and a non-negative  
 369 measure, and  $\alpha(\theta) = 0$  whenever  $|x^1(\theta) - x^T(\theta)| < \varepsilon$ .

370 Here  $E_c$  is the sublevel set defined by (4.1) with the constant  $c = \|\Phi_T^* \Lambda_T^{-1} x^T\|_{L^2(0, T)^m}$  being the  
 371 minimal norm required to reach  $x^T$ .

372 **Remark 4.2.** By the choice of the level  $c$ , it follows that  $x^T \in \partial E_c$ . If additionally the relation (4.2)  
373 holds, i.e., if  $x^T$  is the optimal final state of the problem (P), then  $c = \|\Phi_T^* \Lambda_T^{-1} x^T\|_{L^2(0,T)^m}$  is the  
374 smallest level such that the intersection of the sublevel set  $E_c$  with the target box is non-empty  
(see Figure 1).




Figure 1: Set  $E_c$  for  $n = 1$  and  $K = 2$ , together with the target ball, the optimal final state  $x^T$  and the normal functional  $n_{E_c}(x^T)$ . Each  $x \in \mathcal{C}(\Theta)$  is presented by a point  $(x(\theta_1), x(\theta_2)) \in \mathbb{R}^2$ . For this plot, we have set  $A(\theta_1) = 1$ ,  $A(\theta_2) = 5$ ,  $B(\theta_1) = 1$ ,  $B(\theta_2) = 1/50$ ,  $T = 1$ ,  $\varepsilon = 1/4$ , and the plotted set  $E_c$  is for  $c = 1$ .

375

376 *Proof of Theorem 4.1.*  $\implies$  Let us assume that  $x^T$  is the optimal final state of the problem (P).  
377 In that case  $\frac{1}{2}n_{E_c}(x^T) = \Lambda_T^{-1}x^T$  is the optimal measure and the result follows directly from  
378 Proposition 3.3 (ensuring that it is of the form  $\alpha(x^1 - x^T)|\mu_{opt}|$ , with  $\alpha$  and  $|\mu_{opt}|$  being a non-  
379 negative scalar function and measure, respectively) and the description of the support of the  
380 optimal measure (3.3).

381  $\Leftarrow$  Assume that (4.2) holds. In order to prove the statement, it is enough to show that  $x^T$  is the  
382 only element in the intersection of the sublevel set  $E_c$  and the  $\varepsilon$  ball around the target  $x^1 \in \mathcal{C}(\Theta)^n$ .  
383 We assume the contrary, i.e., that there exists some  $\tilde{x}^T \neq x^T$  lying in the above intersection.

384 We split the parameter set into two disjoint subsets, defined as

$$\Theta_b = \{\theta \in \Theta \mid |x^1(\theta) - x^T(\theta)| = \varepsilon\} \quad \text{and} \quad \Theta_i = \{\theta \in \Theta \mid |x^1(\theta) - x^T(\theta)| < \varepsilon\}.$$

385 By the assumption made on  $\alpha$ ,  $\Theta_i$  is a zero measure set for  $n_{E_c}(x^T)$ . On the other hand, for any  
386  $\theta \in \Theta_b$ , the vector  $x^T(\theta)$  lies on the boundary of the  $\varepsilon$  ball (in  $\mathbb{R}^n$ ) around the target  $x^1(\theta)$ , which,  
387 together with the fact that  $\tilde{x}^T(\theta)$  belongs to the same ball, implies that

$$\langle (x^1 - x^T)(\theta), (\tilde{x}^T - x^T)(\theta) \rangle \geq 0 \quad (\theta \in \Theta_b). \quad (4.3)$$

388 Consequently, we get

$$\begin{aligned} \langle n_{E_c}(x^T), \tilde{x}^T - x^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} &= \langle n_{E_c}(x^T), \tilde{x}^T - x^T \rangle_{\mathcal{M}(\Theta^1)^n, \mathcal{C}(\Theta^1)^n} \\ &= \left\langle (\alpha(x^1 - x^T))_b \nu, \tilde{x}^T - x^T \right\rangle_{\mathcal{M}(\Theta^1)^n, \mathcal{C}(\Theta^1)^n} \\ &= \langle \alpha_b \nu, (x^1 - x^T) \cdot (\tilde{x}^T - x^T) \rangle_{\mathcal{M}(\Theta^1)^n, \mathcal{C}(\Theta^1)^n} \geq 0, \end{aligned} \quad (4.4)$$

389 where the final inequality follows by (4.3) and the assumed non-negativity of the measure  $\alpha_b \nu$ .



390 As  $n_{E_c}(x^T)$  is the normal functional, the only way for  $\tilde{x}^T \in E_c \cap \mathcal{B}_\infty(x^1, \varepsilon)$  to satisfy the  
 391 inequality (4.4) is that  $\langle n_{E_c}(x^T), \tilde{x}^T - x^T \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} = 0$ , i.e., that  $\tilde{x}^T$  belongs to the tangential  
 392 space  $T_{E_c}(x^T)$ . As the convex set  $E_c$  intersects the tangential space  $T_{E_c}(x^T)$  only in its boundary  
 393 points, it follows that  $\tilde{x}^T \in \partial E_c$ . In such way we obtain two distinct points  $\tilde{x}^T$  and  $x^T$  lying in  
 394  $\partial E_c \cap \mathcal{B}_\infty(x^1, \varepsilon)$ . However, that would imply existence of two different feasible controls of the same  
 395 norm,  $c$ , which contradicts the uniqueness of the solution to problem (P).  $\square$

396 **Remark 4.3.** Note that the relation (4.2) implies that the optimal measure

$$\mu_{opt} = \Lambda_T^{-1} x^T = \frac{1}{2} n_{E_c}(x^T) \quad (4.5)$$

397 has, in every point  $\theta \in \Theta$ , the same direction as  $(x^1 - x^T)(\theta)$ , which is in accordance with the  
 398 results of Proposition 3.3.

399 The last result allows an alternative, more direct proof of Proposition 3.6, in the case where  
 400 the parameter set is of finite cardinal, by which one can construct a series of problems for which  
 401 the exact solution is known.

402 *Alternative proof of Proposition 3.6.* With the notations and assumptions of Proposition 3.6, let  
 403  $c = \|\Phi_T^* \Lambda_T^{-1} \tilde{x}^T\|_{L^2(0, T)^m}$  be the minimal norm required to reach the state  $\tilde{x}^T$ . Then for the normal  
 404 functional  $n_{E_c}(\tilde{x}^T)$  we have

$$n_{E_c}(\tilde{x}^T) = 2\Lambda_T^{-1} \tilde{x}^T = 2f_b \eta = 2 \left( \frac{|f|}{\varepsilon} (y - \tilde{x}^T) \right)_b \eta,$$

405 where  $\eta \in \mathcal{M}(\Theta)$  and  $f \in \mathcal{C}(\Theta)^n$  are the nonnegative scalar measure and the continuous vector  
 406 function from the statement of Proposition 3.6 such that  $\Lambda_T^{-1} \tilde{x}^T = f_b \eta$ .

407 From here we see that the normal  $n_{E_c}(\tilde{x}^T)$  is of the form (4.2) (note that  $y = x^1$  for the zero  
 408 initial datum). The statement now follows from Theorem 4.1.  $\square$

409 The introduced sets  $E_c$  allow us to describe the set of targets for which the associated optimal  
 410 control vanishes for some parameter  $\theta \in \Theta$ . This allows us to construct examples for which the  
 411 exact solution is known. This is important in numerical implementations, where the detection of  
 412 active parameters plays a crucial role in constructing the optimal measure.

413 For simplicity, the discussion is provided for scalar systems, i.e., we assume  $n = 1$ .

414 Given some optimal final state  $x^T$ , we know that the corresponding optimal measure is of the  
 415 form (4.5). In particular, its  $i$ -th component (with  $i \in \{1, \dots, K\}$  and  $K = \text{card } \Theta$ ) is going to be  
 416 zero measure if and only if  $x^T = \Lambda_T \mu_{opt}$  lies in the hyperplane  $H_i = [\Lambda_1, \dots, \Lambda_{i-1}, \Lambda_{i+1} \dots \Lambda_K]$   
 417 spanned by all the columns of the Gramian matrix except the  $i$ -th one. This happens exactly for  
 418 the states  $x^T$  for which the normal derivative  $n_{E_c}(x^T) = 2\mu_{opt}$  is orthogonal to the  $i$ -th coordinate  
 419 axes (Figure 2).

420 As any target  $x^1$  lies at most  $\sqrt[2]{K} \varepsilon / 2$  from the optimal final state, this implies that the set of  
 421 targets  $N_i$  for which the  $i$ -th component of the optimal measure equals zero is contained within the  
 422 hyperstrip  $H_i^\varepsilon := H_i + B(0, \sqrt[2]{K} \varepsilon / 2)$ . More precise characterization of this set can be obtained in  
 423 the case  $K = 2$ , i.e., when the parameter obtains only two values. In that case, the set of targets  
 424 for which the  $i$ -th component of the optimal measure is zero is given by

$$N_i = \left\{ x^T + \varepsilon \left( \frac{n_{E_c}(x^T)}{|n_{E_c}(x^T)|} + \beta e_i \right) \mid x^T \in H_i, |\beta| \leq 1 \right\} \cup \mathcal{B}_\infty(0, \varepsilon),$$

425 and depicted on Figure 3.




Figure 2: The hyperplanes  $H_1$  and  $H_2$  for  $n = 1$  and  $K = 2$ . Arrows represent  $n_{E_c}(x^T)$  for different values of  $c$  and well-chosen values of  $x^T$ . The parameters used for this plot are the ones of Figure 1.




Figure 3: Sets  $N_1$  and  $N_2$  for  $n = 1$  and  $K = 2$ . The parameters used for this plot are the ones of Figure 1.

426 For numerical implementations, it would be very useful to know whether the given target  
 427 belongs to some set  $N_i$ . This is because revealing the support of the optimal measure is often the  
 428 most challenging part of the numerical process, while, as we shall see in Section 7 (see in particular  
 429 Examples 7.1 and 7.2), tuning the weights of active Dirac distributions turns out to be easier.  
 430 However, the description of the sets  $N_i$  requires computation of the Gramian operator, which is a  
 431 computationally very demanding task (therefore construction of the Gramian is generally avoided  
 432 and instead various iterative procedures are used for solving control optimization problems). For  
 433 this reason, the above description of targets for which the optimal measure vanishes at some points  
 434 cannot directly accelerate numerical procedures. However, it contributes to a better understanding  
 435 of the problem and its solution, and allows us to describe some characteristics of the optimal  
 436 measures, in particular the sparsity.

437 In general, we can say that for the targets that are not contained in the union of hyperstrips  
 438  $\cup_i H_i^\varepsilon$  the optimal measure is going to be fully supported. Of course, as hyperstrips  $H_i^\varepsilon$ ,  $i \in$   
 439  $\{1, \dots, K\}$  cover just a smaller portion of the whole output space  $\mathbb{R}^K$ , this will be the case for  
 440 the most of the targets. This might look surprisingly, having in mind that the term  $\varepsilon \|\mu\|_{\mathcal{M}(\Theta)^n}$   
 441 entering the dual functional (3.1) is supposed to imply the sparsity in the support of the optimal  
 442 measure. However, in the discrete case, the norm  $\|\cdot\|_{\mathcal{M}(\Theta)^n}$  is equivalent to the  $\ell^1$  norm (see  
 443 comments preceding Theorem 3.1), while the size of the measure support (i.e., number of active  
 444 Dirac distributions) is given by the  $\ell^0$  norm. Although the  $\ell^1$  norm might be used to recover sparse  
 445 solutions under suitable assumptions (see e.g. [7]), in general the  $\ell^0$  term is the one that induces  
 446 sparsity.

## 447 5 Discretized problem and convergence

448 In this section, we show that the solution to the problem (P) can be arbitrarily well approximated  
 449 by a solution to a discretized problem taking into account only a finite number of parameter  
 450 values. This is of crucial importance from the numerical point of view, as the minimization of the  
 451 dual functional in the latter case (originally defined on  $\mathcal{M}(\Theta)^n$ ) reduces to the minimization of a  
 452 function on  $\mathbb{R}^{K^n}$  for some large enough  $K \in \mathbb{N}^*$ .

453 To this end, let  $h > 0$  and consider  $\Theta_h$  a compact subset of  $\Theta$  such that  $\text{dist}(\theta, \Theta_h) \leq h$  for  
 454 every  $\theta \in \Theta$ . The discretized version of (P) is

$$\begin{aligned}
 & \min \quad \frac{1}{2} \|u\|_{L^2(0,T)^m}^2, \\
 & \left| \begin{array}{l} u \in L^2(0,T)^m \quad \text{and} \quad \sup_{\theta \in \Theta_h} \left| x^1(\theta) - e^{TA(\theta)} x^0(\theta) - \int_0^T e^{(T-t)A(\theta)} B(\theta) u(t) dt \right| \leq \varepsilon. \end{array} \right. \\
 & \hspace{20em} (P_h)
 \end{aligned}$$

455 Following [18], we have the following results.

456 **Theorem 5.1.** *Assume that (2.1) is uniformly ensemble controllable. Given  $y \in \mathcal{C}(\Theta)^n$  and*  
 457  *$\varepsilon > 0$ , we set  $u_{opt} \in L^2(0,T)^m$  (respectively  $u_h \in L^2(0,T)$ ) the unique minimizer of (P) (respec-*  
 458 *tively  $(P_h)$ ). Let us also set  $\mu_{opt} \in \mathcal{M}(\Theta)^n$  and  $\mu_h \in \mathcal{M}(\Theta_h)^n$  the unique minimizers of  $J$  defined*  
 459 *by (3.1) over  $\mathcal{M}(\Theta)^n$  and  $\mathcal{M}(\Theta_h)^n$  respectively.*

460 *We have  $\|u_h\|_{L^2(0,T)^m} \leq \|u_{opt}\|_{L^2(0,T)^m}$ , for every  $h > 0$ , and in addition,  $u_h \rightarrow u_{opt}$  in  $L^2(0,T)^m$*   
 461 *and  $\mu_h$  vaguely converges to  $\mu_{opt}$  as  $h \rightarrow 0$ .*

462 Before giving the proof of this result, let us first recall that according to Proposition 2.4,  
 463 if (2.1) is uniformly ensemble controllable with parameter set  $\Theta$ , then (2.1) is uniformly ensemble  
 464 controllable with parameter set  $\Theta_h$ . Hence, the existence and uniqueness of  $u_h$  and  $\mu_h$  (and of  $u_{opt}$   
 465 and  $\mu_{opt}$ ) directly follow from Theorem 3.1. Let us also mention that the strong convergence

466 of  $(u_h)_h$  has already been given in [18, §2.2], under the additional assumption that  $A$ ,  $B$  and  $y$  are  
 467 Lipschitz function with respect to  $\theta$ . It seems that this regularity assumption is not required, and  
 468 we give this proof below.

469 *Proof of Theorem 5.1.* We identify  $\mu_h$  to a measure in  $\mathcal{M}(\Theta)$  supported on  $\Theta_h$ . Since  $\Theta_h \subset \Theta$ , it  
 470 is obvious that  $\|u_h\|_{L^2(0,T)^m} \leq \|u_{opt}\|_{L^2(0,T)^m}$ .

**Step 1.** Let us show that for every sequence  $(h_k)_{k \in \mathbb{N}} \in (\mathbb{R}_+^*)^{\mathbb{N}}$  converging to 0,  $\|\mu_{h_k}\|_{\mathcal{M}(\Theta)^n}$  is uniformly bounded with respect to  $k$ .

Assume by contradiction that there exist a sequence  $(h_k)_{k \in \mathbb{N}} \in (\mathbb{R}_+^*)^{\mathbb{N}}$  converging to 0 such that  $\|\mu_{h_k}\|_{\mathcal{M}(\Theta)^n} \rightarrow \infty$  as  $k \rightarrow \infty$ . We then set  $\lambda_k = \|\mu_{h_k}\|_{\mathcal{M}(\Theta)^n}$ , and we assume without loss of generality that  $\lambda_k > 0$  for every  $k \in \mathbb{N}$ , and we set  $\varsigma_k = \mu_{h_k}/\lambda_k$ . By vague compactness (see e.g. [2, Chapter IV]), there exist a subsequence (still denoted by  $(\varsigma_k)_{k \in \mathbb{N}}$ ) such that  $(\varsigma_k)_k$  is vaguely convergent to  $\varsigma_0 \in \mathcal{M}(\Theta)^n$ . According to eq. (3.2), we have

$$\lambda_k \|\Phi_T^* \varsigma_k\|_{L^2(0,T)}^2 = \langle \varsigma_k, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon \|\varsigma_k\|_{\mathcal{M}(\Theta)^n} = \langle \varsigma_k, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon.$$

471 This ensures that  $\|\Phi_T^* \varsigma_k\|_{L^2(0,T)} \rightarrow 0$  as  $k \rightarrow \infty$ , i.e.,  $\Phi_T^* \varsigma_0 = 0$ , and hence,  $\varsigma_0 = 0$  by Propo-  
 472 sition 2.7. Hence, for  $k$  large enough, we have,  $\lambda_k \|\Phi_T^* \varsigma_k\|_{L^2(0,T)}^2 = \langle \varsigma_k, y \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} - \varepsilon < 0$ ,  
 473 which leads to a contradiction with  $\lambda_k \|\Phi_T^* \varsigma_k\|_{L^2(0,T)}^2 \geq 0$ .

474 **Step 2.** Let us consider a sequence  $(h_k)_{k \in \mathbb{N}} \in (\mathbb{R}_+^*)^{\mathbb{N}}$  such that  $h_k \rightarrow 0$  as  $k \rightarrow \infty$ , and assume  
 475 that  $(\mu_{h_k})_{k \in \mathbb{N}}$  converges vaguely in  $\mathcal{M}(\Theta)^n$ . We aim to prove that  $(\mu_{h_k})_{k \in \mathbb{N}}$  converges vaguely  
 476 to  $\mu_{opt}$ , and that the corresponding sequence of controls  $(u_{h_k})_{k \in \mathbb{N}}$  converges strongly to  $u_{opt}$  in  
 477  $L^2(0, T)^m$ .

478 Let us denote by  $\mu_0$  the vague limit of  $(\mu_{h_k})_k$  and we set  $u_0 = \Phi_T^* \mu_0 \in L^2(0, T)^m$ . For every  
 479  $\varphi \in L^2(0, T)^m$ , we have as  $k \rightarrow \infty$ ,

$$\langle u_{h_k}, \varphi \rangle_{L^2(0,T)^m} = \langle \mu_{h_k}, \Phi_T \varphi \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} \longrightarrow \langle \mu_0, \Phi_T \varphi \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n} = \langle u_0, \varphi \rangle_{L^2(0,T)^m}.$$

480 This ensures the weak convergence of  $(u_{h_k})_{k \in \mathbb{N}}$  to  $u_0$ , in particular, we have  $\|u_0\|_{L^2(0,T)^m} \leq$   
 481  $\liminf_{k \rightarrow \infty} \|u_{h_k}\|_{L^2(0,T)^m} \leq \|u_{opt}\|_{L^2(0,T)^m}$ .

Since  $A$  and  $B$  are continuous on the compact set  $\Theta$ , there are uniformly continuous and bounded. In addition, it is easy to see that there exists a constant  $C > 0$ , depending only on  $T$  and on the bounds of  $A$  and  $B$ , such that

$$\left| e^{(T-t)A(\theta_1)} B(\theta_1) - e^{(T-t)A(\theta_0)} B(\theta_0) \right| \leq C (|A(\theta_1) - A(\theta_0)| + |B(\theta_1) - B(\theta_0)|)$$

$$(t \in [0, T], \theta_0, \theta_1 \in \Theta).$$

482 This ensures that for every  $v \in L^2(0, T)^m$  and every  $\theta_0, \theta_1 \in \Theta$  we have

$$|(\Phi_T v)(\theta_1) - (\Phi_T v)(\theta_0)| \leq \sqrt{TC} \|v\|_{L^2(0,T)^m} (|A(\theta_1) - A(\theta_0)| + |B(\theta_1) - B(\theta_0)|).$$

483 The last inequality clearly implies the uniform continuity of  $\Phi_T u_{h_k}$ , more precisely, for every  $\delta > 0$ ,  
 484 there exists  $\eta(\delta) > 0$  (with  $\eta(\delta) \rightarrow 0$  as  $\delta \rightarrow 0$ ) such that for every  $k \in \mathbb{N}$  and every  $\theta_0, \theta_1 \in \Theta$   
 485 satisfying  $|\theta_1 - \theta_0| \leq \delta$ , we have

$$|(\Phi_T u_{h_k})(\theta_1) - (\Phi_T u_{h_k})(\theta_0)| \leq \eta(\delta).$$

486 By taking larger  $\eta(\delta)$  if needed, we can also assume that

$$|y(\theta_1) - y(\theta_0)| \leq \eta(\delta).$$

487 Indeed,  $y$  is also uniformly continuous on  $\Theta$ , since  $y$  is continuous on the compact set  $\Theta$ .

For every  $\theta \in \Theta$  and every  $k \in \mathbb{N}$ , let us set  $\theta_{h_k} \in \Theta_{h_k}$  such that  $|\theta - \theta_{h_k}| \leq h_k$ . We recall that since  $\theta_{h_k} \in \Theta_{h_k}$ , we have  $|(\Phi_T u_{h_k})(\theta_{h_k}) - y(\theta_{h_k})| \leq \varepsilon$ . We then have, for every  $\theta \in \Theta$  and every  $k \in \mathbb{N}$ ,

$$\begin{aligned} |(\Phi_T u_0)(\theta) - y(\theta)| &\leq |(\Phi_T u_0)(\theta) - (\Phi_T u_{h_k})(\theta)| + |(\Phi_T u_{h_k})(\theta) - (\Phi_T u_{h_k})(\theta_{h_k})| \\ &\quad + |(\Phi_T u_{h_k})(\theta_{h_k}) - y(\theta_{h_k})| + |y(\theta_{h_k}) - y(\theta)| \\ &\leq |(\Phi_T u_0)(\theta) - (\Phi_T u_{h_k})(\theta)| + 2\eta(h_k) + \varepsilon. \end{aligned}$$

488 Taking the limit  $k \rightarrow \infty$ , we conclude that  $|(\Phi_T u_0)(\theta) - y(\theta)| \leq \varepsilon$ . This ensures that  $u_0$  solves the  
489 uniform ensemble controllability problem with parameter set  $\Theta$ . Thus, we have  $\|u_0\|_{L^2(0,T)^m} \geq$   
490  $\|u_{opt}\|_{L^2(0,T)^m}$ . But, we already know that  $\|u_0\|_{L^2(0,T)^m} \leq \|u_{opt}\|_{L^2(0,T)^m}$  from the weak conver-  
491 gence of the sequence  $(u_{h_k})_k$ . This ensures that  $\|u_0\|_{L^2(0,T)^m} = \|u_{opt}\|_{L^2(0,T)^m}$ , and by unique-  
492 ness of the minimal  $L^2$ -norm control,  $u_0 = u_{opt}$ . Finally, from  $\|u_{opt}\|_{L^2(0,T)^m} \geq \|u_{h_k}\|_{L^2(0,T)^m}$   
493 and  $\|u_0\|_{L^2(0,T)^m} \leq \liminf_{k \rightarrow \infty} \|u_{h_k}\|_{L^2(0,T)^m}$ , we conclude that  $\lim_{k \rightarrow \infty} \|u_{h_k}\|_{L^2(0,T)^m} =$   
494  $\|u_{opt}\|_{L^2(0,T)^m}$ . This ensures that  $u_{h_k} \rightarrow u_{opt}$  strongly in  $L^2(0,T)^m$  as  $k \rightarrow \infty$ .

495 The fact that  $u_0 = u_{opt}$  also ensures that  $\Phi_T^* \mu_0 = \Phi_T^* \mu_{opt}$  and hence,  $\mu_0 = \mu_{opt}$  by Propo-  
496 sition 2.7.

497 **Conclusion.** Assume by contradiction that  $\mu_h$  is not vaguely convergent to  $\mu_{opt}$  as  
498  $h \rightarrow 0$ . Then, there exists  $f \in \mathcal{C}(\Theta)^n$  such that  $\langle \mu_h, f \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$  does not converge to  
499  $\langle \mu_{opt}, f \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}$  as  $h \rightarrow 0$ . In particular, there exists  $r > 0$  and a sequence  $(h_k)_{k \in \mathbb{N}} \in (\mathbb{R}_+^*)^{\mathbb{N}}$   
500 converging to 0 such that  $|\langle \mu_{h_k} - \mu_{opt}, f \rangle_{\mathcal{M}(\Theta)^n, \mathcal{C}(\Theta)^n}| > r$ . But according to step 1,  $(\mu_{h_k})_{k \in \mathbb{N}}$   
501 is uniformly bounded. Using vague compactness, up to the extraction of a subsequence  $(\mu_{h_k})_k$  is  
502 vaguely convergent. Finally, the step 2 ensures that the only possible vague limit is  $\mu_{opt}$ . This  
503 leads to a contradiction, and concludes the proof.  $\square$

504 **Remark 5.2.** Let us mention that the arguments used in the first step of the proof of Theorem 5.1  
505 are similar to the ones used in Lemma 3.9. This type of similarity is expected since in Theorem 5.1,  
506 we prove continuity of the minimizers of  $J$  with respect to the parameter set  $\Theta$ , while Lemma 3.9  
507 is an ingredient to prove continuity with respect to  $T$ ,  $\varepsilon$  and  $y$  (see Proposition 3.10).

508 Let us now consider a discretized version of the problem. To this end, we consider  $K \in \mathbb{N}^*$ ,  
509  $\theta_1, \dots, \theta_K \in \Theta$  be two by two distinct,  $\Theta_{h_K} = \{\theta_1, \dots, \theta_K\}$ , with  $h_K = \max_{\theta \in \Theta} \text{dist}(\theta, \Theta_{h_K})$ . Let  
510  $\mu_{h_K}$  be the minimizer of  $J$  defined by (3.1) over  $\Theta_{h_K}$ , then  $\mu_{h_K} = \sum_{k=1}^K w_k \delta_{\theta_k}$ , where  $\delta_{\theta_k}$  is the  
511 atomic mass located at  $\theta_k$ , and  $w = (w_k)_{k \in \{1, \dots, K\}} \in (\mathbb{R}^n)^K$  is the unique minimizer of

$$J_{\Theta_{h_K}}(w) = \frac{1}{2} \int_0^T \left| \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \right|^2 dt - \sum_{k=1}^K \langle w_k, y(\theta_k) \rangle + \varepsilon \sum_{k=1}^K |w_k| \quad (5.1)$$

and we have,

$$u_{h_K}(t) = \Phi_T^* \mu_{h_K} = \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \quad (t \in [0, T]).$$

512 In order to prepare § 6.1, let us give the following results.

513 **Proposition 5.3.**  $w = (w_k)_{k \in \{1, \dots, K\}} \in (\mathbb{R}^n)^K$  is a minimizer of  $J_{\Theta_{h_K}}$  if and only if for every  
514  $l \in \{1, \dots, K\}$ ,  $w_l \in \mathbb{R}^n$  is a minimizer of

$$J_l : \hat{w} \in \mathbb{R}^n \mapsto \frac{1}{2} \hat{w}^* \Lambda_{\theta_l} \hat{w} - \left\langle \hat{w}, y(\theta_l) - \int_0^T e^{(T-t)A(\theta_l)} B(\theta_l) u_l(t) dt \right\rangle + \varepsilon |\hat{w}|,$$

with

$$u_l(t) = \sum_{\substack{k=1 \\ k \neq l}}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \quad \text{and} \quad \Lambda_{\theta_l} = \int_0^T e^{(T-t)A(\theta_l)} B(\theta_l) B(\theta_l)^* e^{(T-t)A(\theta_l)^*} dt.$$

*Proof.* According to Theorem A.1,  $w = (w_k)_{k \in \{1, \dots, K\}}$  is a minimizer of  $J_{\Theta_{h_K}}$  if and only if for every  $l \in \{1, \dots, K\}$ ,  $w_l$  is a minimizer of  $\widehat{w} \in \mathbb{R}^n \mapsto J_{\Theta_{h_K}}(\widehat{w})$  with  $\widehat{w}_k = \begin{cases} w_k, & \text{if } k \neq l, \\ \widehat{w}, & \text{otherwise} \end{cases}$ .

But, for every  $l \in \{1, \dots, K\}$ , we have,

$$\begin{aligned} J_{\Theta_{h_K}}(w) &= \frac{1}{2} \int_0^T \left| \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \right|^2 dt - \sum_{k=1}^K \langle w_k, y(\theta_k) \rangle + \varepsilon \sum_{k=1}^K |w_k| \\ &= \frac{1}{2} \int_0^T \left| B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l \right|^2 dt \\ &\quad + \int_0^T \left\langle B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l, \sum_{\substack{k=1 \\ k \neq l}}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \right\rangle dt - \langle w_l, y(\theta_l) \rangle + \varepsilon |w_l| \\ &\quad + \frac{1}{2} \int_0^T \left| \sum_{\substack{k=1 \\ k \neq l}}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k \right|^2 dt - \sum_{\substack{k=1 \\ k \neq l}}^K \langle w_k, y(\theta_k) \rangle + \varepsilon \sum_{\substack{k=1 \\ k \neq l}}^K |w_k|. \end{aligned}$$

515 It is then clear that minimizers of  $\widehat{w} \mapsto J_{\Theta_{h_K}}(\widehat{w})$  are minimizers of  $J_l$ . □

516 Let us now give condition on the minimizers of  $J_l$  defined in Proposition 5.3.

517 **Proposition 5.4.** *Let  $\Lambda \in \mathbb{R}^{n \times n}$ , be a positive definite matrix,  $z \in \mathbb{R}^n$  and  $\varepsilon > 0$ , the minimizer*  
 518  *$p_{opt} \in \mathbb{R}^n$  of  $\mathcal{J} : p \in \mathbb{R}^n \mapsto \frac{1}{2} p^* \Lambda p - \langle p, z \rangle + \varepsilon |p|$  over  $\mathbb{R}^n$  is unique and satisfies  $p_{opt} = 0$  if  $|z| < \varepsilon$ ,*  
 519 *and otherwise  $p_{opt} = \rho q$  where,*

- 520
  - $|q| = 1$ ;
  - $\text{shrink}(|z|, \varepsilon)/C_\Lambda \leq \rho \leq \text{shrink}(|z|, \varepsilon)/c_\Lambda$ ;
  - $(\varepsilon I + \rho \Lambda)q = z$ ,

523 where  $0 < c_\Lambda \leq C_\Lambda$  are such that

$$c_\Lambda |p|^2 \leq p^* \Lambda p \leq C_\Lambda |p|^2 \quad (p \in \mathbb{R}^n)$$

524 and

$$\text{shrink}(s, \mu) = \begin{cases} s + \mu & \text{if } s < -\mu, \\ 0 & \text{if } -\mu \leq s \leq \mu, \\ s - \mu & \text{if } \mu < s \end{cases} \quad (s \in \mathbb{R}, \mu \in \mathbb{R}_+).$$

525 **Remark 5.5.** When  $n = 1$ , we have  $p_{opt} = \text{shrink}(z, \varepsilon)/\Lambda$ .

526 *Proof.* First,  $\mathcal{J}$  is strictly convex and coercive, hence admits a unique minimizer  $p_{opt}$  on  $\mathbb{R}^n$ .  
 527 Furthermore,  $p_{opt}$  is characterized by  $0 \in \partial \mathcal{J}(p_{opt})$ . But, we have

$$\partial \mathcal{J}(p) = \begin{cases} \Lambda p - z + \varepsilon \frac{p}{|p|} & \text{if } p \neq 0, \\ \overline{B}(0, \varepsilon) - z & \text{if } p = 0, \end{cases}$$

528 recall that  $\bar{B}(0, \varepsilon) \subset \mathbb{R}^n$  is the closed ball of  $\mathbb{R}^n$  centered on 0 and of radius  $\varepsilon$ . In particular, we  
 529 have  $0 \in \partial \mathcal{J}(0)$  if  $|z| \leq \varepsilon$ . Let us then assume that  $|z| > \varepsilon$ , we then have  $p_{opt} \neq 0$ , and we set  
 530  $\rho = |p_{opt}|$  and  $q = p_{opt}/\rho$ . We thus have  $(\varepsilon I + \rho \Lambda)q = z$ , and hence  $\rho > 0$  shall be such that  
 531  $1 = |q|^2 = |(\varepsilon I + \rho \Lambda)^{-1}z|^2$ . This leads to the bounds on  $\rho$ .  $\square$

Before entering the algorithmic computation of a minimizer of  $J_{\Theta_{h_K}}$ , let us give the sub-  
 differential of  $J_{\Theta_{h_K}}$ . For every  $k \in \{1, \dots, K\}$ , we have,

$$\partial_{w_k} J_{\Theta_{h_K}}(w) = \begin{cases} y(\theta_k) - \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) u(t) dt + \varepsilon \frac{w_k}{|w_k|}, & \text{if } w_k \neq 0, \\ y(\theta_k) - \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) u(t) dt + \mathcal{B}(0, \varepsilon), & \text{if } w_k = 0, \end{cases}$$

532 with  $u(t) = \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k$ . In particular, if  $0 \in \partial_{w_k} J_{\Theta_{h_K}}(w)$ , we have

$$\text{shrink} \left( \left| y(\theta_k) - \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) u(t) dt \right|, \varepsilon \right) = 0.$$

533 In addition, when  $w_k = 0$ , we have

$$\text{shrink} \left( \left| y(\theta_k) - \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) u(t) dt \right|, \varepsilon \right) = \min \{ |z| \mid z \in \partial_{w_k} J(w) \}.$$

534 Alternatively to the algorithm proposed in § 6.1, one can use a direct optimization based on  
 535 the optimality conditions given in Proposition 3.3. We have the following direct corollary of Pro-  
 536 position 3.3.

537 **Corollary 5.6.**  $w = (w_k)_{k \in \{1, \dots, K\}} \in (\mathbb{R}^n)^K$  is a minimizer of  $J_{\Theta_{h_K}}$  if there exists  $(\alpha_k)_{k \in \{1, \dots, K\}} \in$   
 538  $\mathbb{R}^K$  such that for every  $k \in \{1, \dots, K\}$ ,

- 539 •  $\alpha_k \geq 0$ , and  $\alpha_k = 0$  if  $|y(\theta_k) - (\Phi_T \Phi_T^* \mu_{h_K})(\theta_k)| < \varepsilon$ ;
- 540 •  $w_k = \alpha_k (y(\theta_k) - (\Phi_T \Phi_T^* \mu_{h_K})(\theta_k))$ ,

541 where we have set  $\mu_{h_K} = \sum_{k=1}^K w_k \delta_{\theta_k}$ .

542 These condition will lead to the minimization problem given in § 6.2.

## 543 6 Numerical strategies

544 In this section, we propose three ways to numerically compute a minimizer of  $J_{\Theta_h}$ .

545 The first one, S.0 is basically using existing solvers without taking in consideration differentia-  
 546 bility issues that emerge from non-smoothness of the dual functional. The second one, S.1 is based  
 547 on the optimality conditions given in Corollary 5.6, see § 6.2. The last ones, S.2.0 and S.2.1, use  
 548 the *greedy coordinate descent method* proposed in [22], see § 6.1.

549 Finally, the strategies S.2.0 and S.2.1 can also be combined with the strategies S.0 and S.1. All  
 550 the possibility are given in § 6.3, and compared on some numerical examples in Section 7.

### 551 6.1 Greedy coordinate descent

552 Inspired from [22], we propose the following *greedy coordinate descent* algorithm (Algorithm 1).  
 553 This algorithm is a coordinate descent algorithm, but instead of considering each coordinate se-  
 554 quentially, we select at each iteration the coordinate for which the decrease of the cost function is  
 555 the most important.

556 The convergence of Algorithm 1 is ensured by the following result.

557 **Proposition 6.1.** Let  $f_1, \dots, f_K \in \mathcal{C}(\mathbb{R}^n)$  and  $F \in \mathcal{C}^1((\mathbb{R}^n)^K)$  be all convex. We consider the cost  
 558 function

$$\mathfrak{J}(w_1, \dots, w_K) = F(w_1, \dots, w_K) + \sum_{k=1}^K f_k(w_k) \quad (w_1, \dots, w_K \in \mathbb{R}^n),$$

559 and we assume that  $\mathfrak{J}$  is coercive and strictly convex.

560 Given  $w^0 = (w_1^0, \dots, w_K^0) \in (\mathbb{R}^n)^K$ , we define the sequences  $(k_i)_{i \in \mathbb{N}} \in \{1, \dots, K\}^{\mathbb{N}}$  and  $(w^i)_{i \in \mathbb{N}} \in$   
 561  $((\mathbb{R}^n)^K)^{\mathbb{N}}$  by:

$$k_i \in \operatorname{argmin}_{k \in \{1, \dots, K\}} \left( \min_{\widehat{w} \in \mathbb{R}^n} J(w_1^i, \dots, w_{k-1}^i, \widehat{w}, w_{k+1}^i, \dots, w_K^i) \right), \quad (6.1)$$

and

$$\begin{aligned} w_{k_i}^{i+1} &\in \operatorname{argmin}_{\widehat{w} \in \mathbb{R}^n} J(w_1^i, \dots, w_{k_i-1}^i, \widehat{w}, w_{k_i+1}^i, \dots, w_K^i), \\ w_k^{i+1} &= w_k^i \quad (\forall k \in \{1, \dots, K\} \setminus \{k_i\}), \end{aligned}$$

562 Then  $\lim_{i \rightarrow \infty} w^i = w^{opt} \in (\mathbb{R}^n)^K$ , where  $w^{opt}$  is the minimizer of  $\mathfrak{J}$ .

563 The proof of this result follows the arguments given in [22, Section 2.4] and is given in Appen-  
 564 dix A for the sake of completeness. Note that in [22] the proof is given for  $n = 1$ .

565 **Remark 6.2.** We expected that the convergence result would also hold by selecting

$$k_i \in \operatorname{argmax}_{k \in \{1, \dots, K\}} \operatorname{dist}(0, \partial_{w_k} \mathfrak{J}(w^i)). \quad (6.2)$$

566 But, we did not succeed to prove this result without additional regularity assumptions on the  
 567 functions  $f_k$ . However, in practice, it seems that using the selection rule (6.2), give similar results  
 568 compared to the selection rule (6.1), but is much faster (see examples in Section 7).

---

**Algorithm 1** Greedy coordinate descent for uniform ensemble control.

---

**Require:**  $T > 0, \varepsilon > 0, (A, B, x^0, x^1) \in \mathcal{C}(\Theta)^{n \times n} \times \mathcal{C}(\Theta)^{n \times m} \times \mathcal{C}(\Theta)^n \times \mathcal{C}(\Theta)^n$ , with  $n, m \in \mathbb{N}^*$

**Require:**  $\Theta_h \subset \Theta$  with  $K := \operatorname{card}(\Theta_h) \in \mathbb{N}^*$  and  $\Theta_h = \{\theta_1, \dots, \theta_K\}$

**Require:**  $\eta > 0$

$\triangleright$  A tolerance parameter.

**Ensure:**  $u \in L^2(0, T)^m$  is such that  $\sup_{\theta \in \Theta_h} \left| x^1(\theta) - e^{TA(\theta)} x^0(\theta) - \int_0^T e^{(T-t)A(\theta)} B(\theta) u(t) dt \right| \leq \varepsilon + \delta$

---



---

**Algorithm 1** (continued)

---

```
1: for all  $k \in \{1, \dots, K\}$  do
2:    $w_k \leftarrow 0$ 
3:    $y_k \leftarrow x^1(\theta_k) - e^{TA(\theta_k)}x^0(\theta_k)$ 
4:    $G_k \leftarrow \text{shrink}(|y_k|, \varepsilon)$   $\triangleright G_k = \text{dist}\{0, \partial_{w_k} J_{\Theta_h}(w)\}$ .
5: end for
6:  $u(t) \leftarrow 0$   $\triangleright u(t) = \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k$ .
7:  $c \leftarrow 0$   $\triangleright c = J_{\Theta_h}(w)$ .
8: while  $\sup_{k \in \{1, \dots, K\}} |y_k| > \varepsilon + \eta$  do
9:   for all  $l \in \{1, \dots, K\}$  do  $\triangleright$  Look for the best descent coordinate.
10:    if  $G_l \neq 0$  then
11:     for all  $k \in \{1, \dots, K\}$  do  $\hat{w}_k \leftarrow w_k$ 
12:     end for
13:      $\hat{u}(t) \leftarrow B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l$ 
14:      $\hat{y}_l \leftarrow y_l + \int_0^T e^{(T-t)A(\theta_l)} B(\theta_l) \hat{u}(t) dt$ 
 $\triangleright$  We have  $\hat{y}_l = x^1(\theta_l) - e^{TA(\theta_l)}x^0(\theta_l) - \int_0^T e^{(T-t)A(\theta_l)} B(\theta_l) \sum_{\substack{k=1 \\ k \neq l}}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k dt$ .
15:      $\hat{w}_l \leftarrow \text{APPROXIMATE CONTROL}(\varepsilon, T, A(\theta_l), B(\theta_l), \hat{y}_l)$   $\triangleright$  See Algorithm 2.
16:      $\hat{c}_l \leftarrow J_{\Theta_h}(\hat{w})$ 
17:     else  $\hat{c}_l \leftarrow c$ 
18:     end if
19:   end for
20:    $l \leftarrow \text{an element of } \underset{k \in \{1, \dots, K\}}{\text{argmax}} \hat{c}_k$   $\triangleright l$  is the best coordinate to update.
21:    $\hat{u}(t) \leftarrow B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l$ 
22:    $\hat{y}_l \leftarrow y_l + \int_0^T e^{(T-t)A(\theta_l)} B(\theta_l) \hat{u}(t) dt$ 
23:    $w_l \leftarrow \text{APPROXIMATE CONTROL}(\varepsilon, T, A(\theta_l), B(\theta_l), \hat{y}_l)$ 
24:    $u(t) \leftarrow u(t) - \hat{u}(t) + B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l$   $\triangleright$  We have  $u(t) = \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k$ .
25:    $c \leftarrow J_{\Theta_h}(w)$ 
26:   for all  $k \in \{1, \dots, K\}$  do
27:      $y_k \leftarrow x^1(\theta_k) - e^{TA(\theta_k)}x^0(\theta_k) - \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) u(t) dt$ 
28:     if  $w_k = 0$  then  $G_k \leftarrow \text{shrink}(|y_k|, \varepsilon)$ 
29:     else  $G_k \leftarrow \left| y_k - \varepsilon \frac{w_k}{|w_k|} \right|$ 
30:     end if
31:   end for
32: end while
33: return  $u(t)$ 
```

---

569 **Remark 6.3.** Here are some comments on Algorithm 1.

- 570 • The loop, lines 9 to 19, can be done in parallel.

- 571 • What is computed in the loop, lines 9 to 19, can be stored and reused lines 21 to 23 and 25.
- 572 • Algorithm 2 will be called many times, and hence has to be fast.
- 573 • This algorithm is adapted from [22], where the aim is to find sparse solutions. Thus, as we
- 574 will see on some examples in Section 7, will lead to sparse minimizers. Consequently, it might
- 575 not be optimal when we are in the situation described in Proposition 3.6 and Corollary 3.7,
- 576 i.e., when the optimal measure is fully supported. However, we shall see that the controls
- 577 obtained in this way still provide a fairly good approximation of the optimal one.

---

**Algorithm 2** Computation of an approximate control for the system  $\dot{x} = Ax + Bu$ .

---

**Require:**  $T > 0$ ,  $\varepsilon > 0$  and  $(A, B, y) \in \mathbb{R}^{n \times n} \times \mathbb{R}^{n \times m} \times \mathbb{R}^n$  with  $n, m \in \mathbb{N}^*$

**Ensure:**  $w \in \mathbb{R}^n$  is such that  $\left| y - \int_0^T e^{(T-t)A} B u(t) dt \right| \leq \varepsilon$  with  $u(t) = B^* e^{(T-t)A^*} w$

```

1: function APPROXIMATE CONTROL( $\varepsilon, T, A, B, y$ )
2:   if  $|y| \leq \varepsilon$  then  $w \leftarrow 0$  ▷ See Remark 5.5.
3:   else ▷ In that case, we have  $w \neq 0$ .
4:     if  $n = 1$  then
5:        $\Lambda \leftarrow \int_0^T e^{(T-t)A} B B^* e^{(T-t)A^*} dt$ 
6:        $w \leftarrow \text{shrink}(y, \varepsilon) / \Lambda$  ▷ See Remark 5.5.
7:     else if the control Gramian is “computable” then ▷ e.g. for  $n$  “small”.
8:        $\Lambda \leftarrow \int_0^T e^{(T-t)A} B B^* e^{(T-t)A^*} dt$ 
9:        $(r, q) \leftarrow$  solution of  $r > 0$ ,  $q \in \mathbb{R}^n$ ,  $|q| = 1$  and  $(\varepsilon I + r\Lambda)q(r) = y$ 
10:       $w \leftarrow r q(r)$  ▷ See Proposition 5.4.
11:    else
12:       $w \leftarrow \underset{\nu \in \mathbb{R}^n}{\text{argmin}} \left( \frac{1}{2} \int_0^T |B^* e^{(T-t)A^*} \nu|^2 dt - \langle \nu, y \rangle + \varepsilon |\nu| \right)$ 
13:    end if
14:  end if
15:  return  $w$ 
16: end function

```

---

- 578 **Remark 6.4.** • In Algorithm 2 (line 9), finding a solution of  $r > 0$ ,  $q \in \mathbb{R}^n$ ,  $|q| = 1$  and
- 579  $(\varepsilon I + r\Lambda)q(r) = y$  can be done using a dichotomy search.
- 580 • To find a minimizer of  $J$  (line 12 of Algorithm 2), one can use a gradient descent method.
- 581 In fact, note that  $J$  is only not differentiable at 0, and note that 0 cannot be the minimizer
- 582 in the situation of line 12 this is because  $|y| > \varepsilon$ . A way to avoid the evaluation of  $\mathcal{J}(\nu) =$
- 583  $\frac{1}{2} \int_0^T |B^* e^{(T-t)A^*} \nu|^2 dt - \langle \nu, y \rangle + \varepsilon |\nu|$  at 0 is to initialize the gradient descent method with
- 584  $\nu \in \mathbb{R}^n$  such that  $\mathcal{J}(\nu) < \mathcal{J}(0) = 0$ . This is for instance possible by setting  $\nu = \alpha y$ , with
- 585  $\alpha = \text{shrink}(|y|^2, \varepsilon |y|) / C$ , with  $C = \int_0^T |B^* e^{(T-t)A^*} y|^2 dt$ . We thus have  $\nu \neq 0$  and  $\mathcal{J}(\nu) < 0$ ,
- 586 see Remark 5.5.
- 587 • Computing the Gramian matrix might be resource consuming. Hence, instead of computing
- 588 it in Algorithm 2, it can be computed and stored in Algorithm 1.

589 **6.2 Reduced minimization problem**

590 For this strategy, we will use the optimality condition given in Corollary 5.6. More precisely, we  
 591 consider the following minimization problem (recall that we have set  $y(\theta) = x^1(\theta) - e^{TA(\theta)}x^0(\theta)$ ),

$$\min \frac{1}{2} \left\| \sum_{k=1}^K B(\theta_k)^* e^{(T-\cdot)A(\theta_k)^*} w_k \right\|_{L^2(0,T)^m}^2 \quad (6.3a)$$

592 with respect to  $w_1, \dots, w_K \in \mathbb{R}^n$ , subject to the following constraints

$$\forall k \in \{1, \dots, K\}, \exists \alpha_k \geq 0 \quad \text{such that:} \quad \begin{cases} |y(\theta_k) - y^T(\theta_k)|^2 \leq \varepsilon^2, \\ \alpha_k \left( |y(\theta_k) - y^T(\theta_k)|^2 - \varepsilon^2 \right) \geq 0, \\ w_k = \alpha_k (y(\theta_k) - y^T(\theta_k)), \end{cases} \quad (6.3b)$$

593 where we have set

$$y^T(\theta_k) = \int_0^T e^{(T-t)A(\theta_k)} B(\theta_k) \left( \sum_{l=1}^K B(\theta_l)^* e^{(T-t)A(\theta_l)^*} w_l \right) dt. \quad (6.3c)$$

594 Observe that the last three constraints coincide with the optimality condition given in Corollary 5.6,  
 595 and the control is given by  $u(t) = \sum_{k=1}^K B(\theta_k)^* e^{(T-t)A(\theta_k)^*} w_k$ . Using Corollary 5.6, it is obvious  
 596 that minimizers of  $J_{\Theta_{h_K}}$  coincide with minimizers of (6.3).

597 **Remark 6.5.** In practice the minimization problem (6.3) can be expressed as a minimization only  
 598 on  $(\alpha_k)_{k \in \{1, \dots, K\}}$ , however, from a numerical point of view, it is easier to consider the minimiza-  
 599 tion problem (6.3) as a minimization with respect to  $(w_k, \alpha_k, y(\theta_k))_{k \in \{1, \dots, K\}}$  since gradient and  
 600 Hessians can be easily computed.

601 **6.3 Proposed numerical strategies**

602 Based on the previous consideration, we can propose the following numerical strategies.

603 S.0. Use existing nonlinear solvers, and try to directly minimize  $J_{\Theta_{h_K}}$ , without considering po-  
 604 tential issues related to the non-differentiability of the cost functional. In the examples of  
 605 Section 7, we use the `fminunc` function of `Matlab` with the quasi-Newton algorithm.

606 S.1. Use existing nonlinear solvers to find a minimizer of (6.3). In the examples of Section 7, we  
 607 use the `fmincon` function of `Matlab` with the interior-point algorithm.

608 In addition, one can use algorithms based on Algorithm 1. This leads to two strategies:

609 S.2.0. Use directly Algorithm 1.

610 S.2.1. Use Algorithm 1 but instead of using the selection rule given by (6.1), we use the one given  
 611 by (6.2) (see Remark 6.2).

612 As we will see in Section 7, with strategies S.2.0 and S.2.1, it appears that many steps are between  
 613 non-zeros values. Hence, we propose a refinement of these two algorithms by running S.0 or S.1 after  
 614 some component of  $(w_k)_k$  switches from null to a non-null value, or vice versa. More precisely,  
 615 using notations of Algorithm 1, if  $w_l$  was 0 and is updated to a non-null value or if  $w_l$  was  
 616 non-null and is updated to 0, we run S.0 or S.1 for the same minimization problem, but with  
 617  $\tilde{\Theta} = \{\theta_k \in \Theta_h, \forall k \in \{1, \dots, K\} \text{ s.t. } w_k \neq 0\}$  in place of  $\Theta_h$ . In other words, we try to update all  
 618 the non-zero values of  $(w_k)_k$  in one shoot. This leads to the four additional strategies, S.2.i-S.2.j  
 619 (with  $i, j \in \{0, 1\}$ ).

620 Let us recall that we have analytical guaranties of convergence only for the strategy S.2.0.

## 7 Numerical examples

Unless explicitly said otherwise, for all the following examples

- the final time is  $T = 1$ ;
- $\Theta = [1, 2]$ , and  $\Theta_h$  is a uniform discretization of  $\Theta$  with step  $h = 10^{-2}$ , leading to  $K = \text{card } \Theta_h = 101$ ;
- $\varepsilon = 5 \times 10^{-2}$ ;
- the time integrals are computed using the trapezoidal rule with a uniform time step of  $10^{-4}$ . In particular, to compute  $\Phi_T$ , we use the trapezoidal rule, and the matrix exponentials are computed numerically. Another alternative could have been to solve the ordinary differential equation using a numerical scheme;
- the stopping criterion on the gradient (i.e.,  $\delta$  in Algorithm 1) is set to  $10^{-12}$ , we also set a step tolerance of  $10^{-20}$  (i.e., step sizes shall be greater than this number) and a maximal number of iterations of  $10^5$ .

Examples 7.1 and 7.2 correspond to situation where the optimal control and the adjoint measure are known (see Proposition 3.6 and Corollary 3.7). Example 7.3 corresponds to a situation where the target cannot be reached (see Appendix B). Finally, Example 7.4 tackled the uniform ensemble controllability for a discretized heat equation.

In the tables used for the comparison of the different numerical strategies,  $G = (G_1, \dots, G_K) \in (\mathbb{R}_+)^K$  is given by lines 27 to 29 of Algorithm 1 that is  $G_k = \text{dist}(0, \partial_{w_k} J_{\Theta_h}(w))$ .

The codes are run using MatLab version R2020b on a 12 core processor of 64 bits and 3.20 GHz.

**Example 7.1.** We consider a first basic example, where  $n = m = 1$ ,  $A(\theta) = -\theta$  and  $B(\theta) = 1$ . It is well-known that this system is ensemble controllable (see [21, 27]). We consider  $x^0(\theta) = 0$ ,  $d\mu_{opt}(\theta) = \chi_\omega(\theta) d\theta$ , where  $\omega$  is an open subset of  $\Theta$ . According to Corollary 3.7, we know that  $u_{opt} = \Phi_T^* \mu_{opt}$  is the minimizer of (P) when the target is set to  $x^1(\theta) = (\Phi_T \Phi_T^* \mu_{opt})(\theta) + \varepsilon$ .

In practice, since  $\Theta_h$  (introduced in Section 5) is chosen as a uniform discretization of  $\Theta$ , we set  $\mu_{opt,h} = h \sum_{\theta \in \Theta_h \cap \omega} \delta_\theta$ , and it is clear that  $\mu_{opt,h}$  is vaguely convergent to  $\mu_{opt}$ . Also, the target  $x_h^1$  is set accordingly, i.e.,  $x_h^1(\theta) = (\Phi_T \Phi_T^* \mu_{opt,h})(\theta) + \varepsilon$ , and the discretized optimal control is  $u_{opt,h} = \Phi_T^* \mu_{opt,h}$ .

The numerical simulations are run with  $\omega = (4/3, 7/4)$  (we recall that the other parameters are given in the preamble of Section 7). A comparison of the different strategies listed in § 6.3 is given in Table 1.

Strategy	CPU time (s)	running time (s)	nb. iter.	max $G$	$\frac{\ u_h - u_{opt,h}\ _{L^2}}{\ u_{opt,h}\ _{L^2}}$	$\frac{\ \mu_h - \mu_{opt,h}\ _{\mathcal{M}}}{\ \mu_{opt,h}\ _{\mathcal{M}}}$
S.0	6.07	0.63	7	$2.553 \times 10^{-5}$	$1.561 \times 10^{-3}$	<b>0.959</b>
S.1	4.59	2.82	83	$1.293 \times 10^{-13}$	$4.384 \times 10^{-8}$	1.271
S.2.0	1409.58	201.93	100000	$2.387 \times 10^{-8}$	$2.467 \times 10^{-5}$	1.952
S.2.0-S.0	39.29	8.49	11	$8.161 \times 10^{-10}$	$4.432 \times 10^{-6}$	1.952
S.2.0-S.1	11.19	2.47	14	<b><math>7.452 \times 10^{-15}</math></b>	<b><math>6.833 \times 10^{-9}</math></b>	1.902
S.2.1	198.18	20.01	100000	$3.389 \times 10^{-7}$	$1.140 \times 10^{-3}$	1.800
S.2.1-S.0	<b>1.06</b>	<b>0.19</b>	14	$3.043 \times 10^{-9}$	$1.617 \times 10^{-5}$	1.905
S.2.1-S.1	4.28	1.19	12	$3.757 \times 10^{-14}$	$2.951 \times 10^{-8}$	1.857

(a) Errors, time of computation and number of iterations.

Table 1: Comparison of the strategies listed in § 6.3 for Example 7.1.

Strategy	Stopping reason
S.0	Cannot decrease the objective function along the current search direction.
S.1	Local minimum possible. Constraints satisfied.
S.2.0	Maximal number of iteration reached.
S.2.0-S.0	Cost is not decreasing.
S.2.0-S.1	Cost is not decreasing.
S.2.1	Maximal number of iteration reached.
S.2.1-S.0	Cost is not decreasing.
S.2.1-S.1	Cost is not decreasing.

(b) Stopping reasons.

Table 1: (continued)

652 In Table 1b, for strategies S.2.i-S.j ( $i, j \in \{0, 1\}$ ) the last cost modifications is of order  $10^{-17}$ ,  
653 which is close to the numerical 0 (indeed, by denoting  $c_{opt}$  the minimal cost, we have numerically  
654  $c_{opt} + 10^{-18} = c_{opt}$ ). Let us also mention that for all simulations, we have  $|x^1(\theta) - x(T, \theta)| = \varepsilon$   
655 up to an error of order  $10^{-8}$ . We also see that the strategy S.2.0-S.1 gives the best results and  
656 strategy S.2.1-S.0 is the fastest one. As claimed in § 6.3, for strategy S.2.0 and S.2.1, most of the  
657 time is spent in updating non-zero values in fact for S.2.0, for all  $10^5$  steps the selected index is  
658 in  $\{1, 55, 101\}$ , and for S.2.1, after some iterations the selected index loops between 1, 42 and 101  
659 and then between 42, 69 and 101. In Table 1a, we see that even if the control is closed to the  
660 optimal ones, the adjoint is rather far from the optimal one, this is also illustrated on Figure 4. As  
661 we can see from Figure 4 and Table 1a, wide variation on the adjoint does not necessarily produce  
662 large variations on the control. One can see from Figure 4, that the strategies S.0 and S.1 lead to  
663 adjoints that are fully supported on  $\Theta_h$ , while the other strategies lead to adjoints for which the  
664 support consist in only few points of  $\Theta_h$ . These facts will be discussed later on.




Figure 4: Adjoints obtained with strategies listed in § 6.3 for Example 7.1.




Figure 4: (continued)

665 **Example 7.2.** We consider the system and data given in Example 7.1, except that we set  $\mu_{opt} =$   
 666  $\delta_{\tilde{\theta}_1} + 2\delta_{\tilde{\theta}_2}$ . In that case, for  $i \in \{1, 2\}$ , we set  $\tilde{\theta}_{i,h} \in \operatorname{argmin} \left\{ \left| \tilde{\theta}_i - \theta \right|, \theta \in \Theta_h \right\}$  and  $\mu_{opt,h} =$   
 667  $\delta_{\tilde{\theta}_{1,h}} + 2\delta_{\tilde{\theta}_{2,h}}$ .

668 The numerical simulation is run with  $\tilde{\theta}_1 = 7/6$  and  $\tilde{\theta}_2 = 17/10$  (we recall that the other  
 669 parameters are given in the preamble of Section 7). A comparison of the different strategies listed  
 in § 6.3 is given in Table 2.

Strategy	CPU time (s)	running time (s)	nb. iter.	max $G$	$\frac{\ u_h - u_{opt,h}\ _{L^2}}{\ u_{opt,h}\ _{L^2}}$	$\frac{\ \mu_h - \mu_{opt,h}\ _{\mathcal{M}}}{\ \mu_{opt,h}\ _{\mathcal{M}}}$
S.0	4.11	0.44	19	$2.431 \times 10^{-8}$	$2.421 \times 10^{-5}$	<b>1.960</b>
S.1	2.92	1.78	76	<b><math>2.058 \times 10^{-13}</math></b>	<b><math>2.675 \times 10^{-8}</math></b>	1.981
S.2.0	1478.55	213.11	100000	$7.575 \times 10^{-7}$	$1.065 \times 10^{-4}$	2.000
S.2.0-S.0	0.55	0.10	10	$4.783 \times 10^{-9}$	$1.999 \times 10^{-5}$	2.000
S.2.0-S.1	5.60	1.38	11	$2.553 \times 10^{-12}$	$3.665 \times 10^{-7}$	2.000
S.2.1	194.10	19.49	100000	$2.495 \times 10^{-6}$	$3.447 \times 10^{-4}$	2.000
S.2.1-S.0	<b>0.28</b>	<b>0.04</b>	5	$2.188 \times 10^{-8}$	$4.409 \times 10^{-5}$	2.000
S.2.1-S.1	3.01	0.78	8	$9.378 \times 10^{-12}$	$5.331 \times 10^{-7}$	2.000

(a) Errors, time of computation and number of iterations

Strategy	Stopping reason
S.0	Cannot decrease the objective function along the current search direction.
S.1	Local minimum possible. Constraints satisfied.
S.2.0	Maximal number of iteration reached.
S.2.0-S.0	Step size is less than the step tolerance.
S.2.0-S.1	Step size is less than the step tolerance.
S.2.1	Maximal number of iteration reached.
S.2.1-S.0	Cost is not decreasing.
S.2.1-S.1	Cost is not decreasing.

(b) Stopping reasons.

Table 2: Comparison of the strategies listed in § 6.3 for Example 7.2.

670 We see from Table 2 that almost all the conclusion made for Example 7.1 also holds for Exam-  
 671 ple 7.1, except that here the strategy S.1 give slightly better results than the strategy S.2.0-S.1. As  
 672 for Example 7.1, we display on Figure 5 the adjoints numerically computed. The comments made  
 673 in Example 7.1 also applies here, wide variation on the adjoint does not necessarily produce large  
 674

675 variations on the control, and the strategies **S.0** and **S.1** lead to adjoints that are fully supported  
 676 on  $\Theta_h$ , while the other strategies lead to adjoints for which the support consist in only few points  
 of  $\Theta_h$ .




Figure 5: Adjoints obtained with strategies listed in § 6.3 for Example 7.1.

677

678 Based on the results given in Examples 7.1 and 7.2, we can make the following comments.

679

- 680 • All strategy lead to some acceptable control.
- 681 • Strategies **S.0** and **S.1** lead to adjoint measure, which are everywhere non-zero. In fact, they  
 682 are not adapted for sparsity, contrary to strategies **S.2.0** and **S.2.1** and their derivatives. In  
 683 addition, for strategy **S.1** since an interior-point algorithm is used, the fact that the obtained  
 684 solution is non-zero everywhere is expected.
- 685 • Even if the strategy **S.2.1-S.1** did not give the best results, it is relatively fast and provide  
 good results for the control. We will then use this strategy for the following examples.

686

**Example 7.3.** We consider the system and data given in Example 7.1, but we set  $x^1(\theta) = 1$ . In  
 687 that case, it is known from Proposition B.1 that the target state in not reachable. Hence, it is  
 688 expected that the norm of the minimal  $L^2$ -norm control goes to  $\infty$  as  $\varepsilon$  goes to 0.

689

This fact is illustrated on Figure 6. On Figure 7, we also display the obtained results for  
 690  $\varepsilon = 2 \times 10^{-4}$  (we recall that the other parameters are given in the preamble of Section 7). In  
 691 particular, Figures 7b and 7d confirm the claim of Corollary 3.5, i.e., the sign of the optimal measure

692 coincide with the sign of  $x^1 - x^T$  and the measure is only supported on the set of parameters  $\theta$  such  
 693 that  $|x^1(\theta) - x^T(\theta)| = \varepsilon$ . The results for this example have been obtained using strategy S.2.1–S.1.




Figure 6: Norm of the control, number of Dirac masses in the adjoint state and absolute value of the adjoint state, with respect to  $\varepsilon$ , for Example 7.3.




Figure 7: Results obtained with strategy S.2.1–S.1 for the system given in Example 7.3, with  $\varepsilon = 2 \times 10^{-4}$ .



**Example 7.4.** For this example, we consider a discretized version of the family indexed by  $\theta \in \Theta$  of systems of 1D heat equation:

$$\dot{y}(t, \theta, x) = \partial_x^2 y(t, \theta, x) - (\theta - 1)y(t, \tau, x) + u(t, x) \quad (t > 0, x \in (0, 1)), \quad (7.1a)$$

$$y(t, \theta, 0) = y(t, \theta, 1) = 0 \quad (t > 0), \quad (7.1b)$$

$$y(0, \theta, x) = y^0(\theta, x) \quad (x \in (0, 1)). \quad (7.1c)$$

694 One can refer to [9] for some ensemble controllability results related to parabolic systems.

695 Given  $n \in \mathbb{N}^*$ , using centered finite differences, the discretized version of (7.1) is

$$\dot{Y}(t, \theta) = A(\theta)Y(t, \theta) + B(\theta)U(t), \quad Y(0, \theta) = Y^0(\theta), \quad (7.2)$$

696 with

$$A(\theta) = \frac{1}{(n+1)^2} \begin{pmatrix} -2 & 1 & 0 & \dots & 0 \\ 1 & \ddots & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & 1 \\ 0 & \dots & 0 & 1 & -2 \end{pmatrix} - (\theta - 1)I_n \in \mathbb{R}^{n \times n} \quad \text{and} \quad B(\theta) = I_n \in \mathbb{R}^{n \times n}$$

697 and where  $Y_i(t, \theta) \simeq y(t, \theta, \frac{i}{n+2})$ ,  $U_i(t) \simeq u(t, \frac{i}{n+2})$  and  $Y_i^0(\theta) = y^0(\theta, \frac{i}{n+2})$ .

698 Let us mention that using the trapezoidal rule,  $\|y(t, \theta, \cdot)\|_{L^2(0,1)}$  is approximated by  $\sqrt{\frac{1}{n+1}} |Y(t, \theta)|$

699 hence, the value of  $\varepsilon$  will be adapted accordingly. For the control problem, we set  $y^0(\theta, x) =$

700  $\sin(x + \theta)$  and null target, i.e.,  $y^1(\theta) = 0$ . The goal is to find a control  $u \in L^2(0, T; L^2(0, 1))$

701 such that  $\sup_{\theta \in \Theta} \|y(T, \theta, \cdot)\|_{L^2(0,1)} \leq \varepsilon$ ,  $\varepsilon, T, \Theta$  and discretization parameters are defined in the

702 preamble of Section 7. For the numerical illustration, we use the strategy S.2.1–S.1, with  $n = 19$ .

703 The obtained results are displayed on Figure 8.

## 704 A Proof of Proposition 6.1

705 For this proof, we follow [22, Section 2.4]. We consider a cost function of the form

$$\mathfrak{J}(w_1, \dots, w_K) = F(w_1, \dots, w_K) + \sum_{k=1}^K f_k(w_k) \quad (w_1, \dots, w_K \in \mathbb{R}^n),$$

706 where  $f_1, \dots, f_K$  are continuous convex function and  $F$  is a convex function of class  $\mathcal{C}^1$ . For every

707  $k \in \{1, \dots, K\}$ , we define  $\tilde{w}_k = (w_1, \dots, w_{k-1}, w_{k+1}, \dots, w_K) \in (\mathbb{R}^n)^{K-1}$ , and  $\mathfrak{J}_{k, \tilde{w}_k}(w_k) = \mathfrak{J}(w)$

708 and  $F_{k, \tilde{w}_k}(w_k) = F(w)$ .

709 **Theorem A.1.** Using the notations and assumptions introduced above,  $w = (w_1, \dots, w_K)$  is a

710 minimizer of  $\mathfrak{J}$  if and only if for every  $k \in \{1, \dots, K\}$ ,  $w_k$  is a minimizer of  $\mathfrak{J}_{k, \tilde{w}_k}$ .


711 *Proof.* If  $w$  is a minimizer of  $\mathfrak{J}$ , then for every  $\delta \in \mathbb{R}^n$  and every  $k$ , we have

$$\mathfrak{J}(w_1, \dots, w_{k-1}, w_k + \delta, w_{k+1}, \dots, w_K) \geq \mathfrak{J}(w_1, \dots, w_{k-1}, w_k, w_{k+1}, \dots, w_K),$$


712 That is to say that  $\mathfrak{J}_{k, \tilde{w}_k}(w_k + \delta) \geq \mathfrak{J}_{k, \tilde{w}_k}(w_k)$ , i.e.,  $w_k$  minimizes  $\mathfrak{J}_{k, \tilde{w}_k}$ .

713 Reciprocally, if  $w_k$  minimizes  $\mathfrak{J}_{k, \tilde{w}_k}$ , for every  $\delta \in \mathbb{R}^n$  and every  $t \in [0, 1]$ , we have,


$$0 \leq \mathfrak{J}_{k, \tilde{w}_k}(w_k + t\delta) - \mathfrak{J}_{k, \tilde{w}_k}(w_k) = F_{k, \tilde{w}_k}(w_k + t\delta) - F_{k, \tilde{w}_k}(w_k) + f_k(w_k + t\delta) - f_k(w_k).$$




(a) Obtained control ( $\|u\|_{L^2((0,T)\times(0,1))} \simeq 3.4857$ ).



(b) State at final time.



(c)  $L^2$ -norm of the time dependent solutions for some values of  $\theta$ . On this graph, we plot  $\|y(t, \theta, \cdot)\|_{L^2(0,1)}$ .



(d)  $L^2$ -norm of the obtained adjoint measure.

Figure 8: Results obtained with strategy S.2.1–S.1 for the system given in Example 7.4.

714 But, since  $f_k$  is convex, we have  $f_k(w_k + t\delta) = f_k((1-t)w_k + t(w_k + \delta)) \leq (1-t)f_k(w_k) + tf_k(w_k +$   
 715  $\delta)$ , and we get

$$f_k(w_k + t\delta) - f_k(w_k) \leq t(f_k(w_k + \delta) - f_k(w_k)) \quad (t \in [0, 1]).$$

716 This ensures

$$0 \leq \frac{1}{t} (F_{k, \check{w}_k}(w_k + t\delta) - F_{k, \check{w}_k}(w_k)) + f_k(w_k + \delta) - f_k(w_k) \quad (t \in (0, 1)).$$

717 Since  $F$  is of class  $\mathcal{C}^1$ ,  $F_{k, \check{w}_k}$  is also of class  $\mathcal{C}^1$  and taking the limit  $t \rightarrow 0$  in the above equation,  
 718 we get

$$0 \leq \langle \nabla F_{k, \check{w}_k}(w_k), \delta \rangle + f_k(w_k + \delta) - f_k(w_k).$$

Now, assume that  $w = (w_1, \dots, w_K)$  is such that  $w_k$  minimizes  $F_{k, \check{w}_k}$  for every  $k \in \{1, \dots, K\}$ .

For every  $\delta = (\delta_1, \dots, \delta_K) \in (\mathbb{R}^n)^K$ , we have, using the convexity (and  $\mathcal{C}^1$  regularity) of  $F$ ,

$$\begin{aligned}
\mathfrak{J}(w + \delta) - \mathfrak{J}(w) &= F(w + \delta) - F(w) + \sum_{k=1}^K (f_k(w_k + \delta_k) - f_k(w_k)) \\
&\geq \langle \nabla F(w), \delta \rangle + \sum_{k=1}^K (f_k(w_k + \delta_k) - f_k(w_k)) \\
&= \sum_{k=1}^K (\langle \nabla_{w_k} F(w), \delta_k \rangle + f_k(w_k + \delta_k) - f_k(w_k)) \\
&= \sum_{k=1}^K (\langle \nabla F_{k, \tilde{w}_k}(w_k), \delta_k \rangle + f_k(w_k + \delta_k) - f_k(w_k)) \\
&\geq 0.
\end{aligned}$$

719 That is to say that  $w$  is a minimizer of  $\mathfrak{J}$ . □

720 **Lemma A.2.** *Assume  $\mathfrak{J}$  is strictly convex and coercive. Then, for every  $k \in \{1, \dots, K\}$ , the*  
721 *function  $g_k : (\mathbb{R}^n)^{K-1} \rightarrow \mathbb{R}^n$  defined by  $g_k(\tilde{w}) = \operatorname{argmin} \mathfrak{J}_{k, \tilde{w}}$  is continuous.*

722 *Proof.* Firstly, since we assumed that  $\mathfrak{J}$  is strictly convex and coercive, we have that  $\mathfrak{J}_{k, \tilde{w}}$  is also  
723 strictly convex and coercive, this holds for every  $\tilde{w} \in (\mathbb{R}^n)^{K-1}$  and every  $k \in \{1, \dots, K\}$ . This  
724 ensures the existence and uniqueness of minimizers of  $\mathfrak{J}_{k, \tilde{w}}$ , i.e., the function  $g_k$  is well-defined.

725 Let  $\tilde{w} \in (\mathbb{R}^n)^{K-1}$  and  $(\tilde{w}^i)_{i \in \mathbb{N}} \in \left( (\mathbb{R}^n)^{K-1} \right)^{\mathbb{N}}$  be a sequence converging to  $\tilde{w}$ , for every  
726  $i \in \mathbb{N}$ , we define  $w_k^i = g_k(\tilde{w}^i)$  and  $w_k = g_k(\tilde{w})$ . We then have,  $\mathfrak{J}_{k, \tilde{w}^i}(w_k^i) \leq \mathfrak{J}_{k, \tilde{w}^i}(w_k)$ , and  
727  $\lim_{i \rightarrow \infty} \mathfrak{J}_{k, \tilde{w}^i}(w_k) = \mathfrak{J}_{k, \tilde{w}}(w_k)$ . This ensures that the sequence  $(\mathfrak{J}_{k, \tilde{w}^i}(w_k^i))_i$  is bounded. But  
728 since  $\mathfrak{J}$  is coercive (and recalling that  $\mathfrak{J}_{k, \tilde{w}^i}(w_k^i) = \mathfrak{J}(w_k^i)$ ), this implies that the sequence  $(w_k^i)_i$  is  
729 bounded, and hence admits a convergent subsequence (still denote by  $(w_k^i)_i$ ), and we denote by  
730  $w_k^\infty$  its limit. We then have  $\mathfrak{J}_{k, \tilde{w}}(w_k^\infty) = \lim_{i \rightarrow \infty} \mathfrak{J}_{k, \tilde{w}^i}(w_k^i) \leq \lim_{i \rightarrow \infty} \mathfrak{J}_{k, \tilde{w}^i}(w_k) = \mathfrak{J}_{k, \tilde{w}}(w_k)$ ,  
731 this implies  $w_k = w_k^\infty$ , by uniqueness of the minimizer of  $\mathfrak{J}_{k, \tilde{w}}$ . We have thus proved that  
732  $\lim_{i \rightarrow \infty} g_k(\tilde{w}^i) = g_k(\tilde{w})$ . □

733 We are now in position to prove the convergence of the algorithm.

734 *Proof of Proposition 6.1.* It is trivial that

$$\min_{(\mathbb{R}^n)^K} \mathfrak{J} \leq \mathfrak{J}(w^{i+1}) \leq \mathfrak{J}(w^i) \quad (i \in \mathbb{N}). \tag{A.1}$$

735 This ensures that  $(\mathfrak{J}(w^i))_i$  is bounded and convergent. Due to the coercivity of  $\mathfrak{J}$ , we also have, by  
736 compactness, the existence of  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  increasing and  $w^\infty \in (\mathbb{R}^n)^K$  such that  $\lim_{i \rightarrow \infty} w^{\phi(i)} =$   
737  $w^\infty$ . The continuity of  $\mathfrak{J}$  ensures that  $\mathfrak{J}(w^\infty) = \lim_{i \rightarrow \infty} \mathfrak{J}(w^{\phi(i)}) = \lim_{i \rightarrow \infty} \mathfrak{J}(w^i)$  (the last equality  
738 follows from the uniqueness of limits).

739 For every  $k \in \{1, \dots, K\}$  and every  $w \in (\mathbb{R}^n)^K$ , we define  $\Delta \mathfrak{J}_k(w) = \mathfrak{J}(w_1, \dots, w_K) -$   
740  $\mathfrak{J}(w_1, \dots, w_{k-1}, g_k(\tilde{w}_k), w_{k+1}, \dots, w_K) \geq 0$ , where  $g_k$  is defined in Lemma A.2. From Lemma A.2,  
741  $g_k$  is continuous, hence  $\Delta \mathfrak{J}_k$  is also continuous. We then have  $\Delta \mathfrak{J}_k(w^\infty) = \lim_{i \rightarrow \infty} \Delta \mathfrak{J}_k(w^{\phi(i)})$ .  
742 We also have, by definition of the sequence,

$$0 \leq \Delta \mathfrak{J}_k(w^{\phi(i)}) \leq \Delta \mathfrak{J}_{k, \phi(i)}(w^{\phi(i)}) = \mathfrak{J}(w^{\phi(i)}) - \mathfrak{J}(w^{\phi(i)+1}) \quad (i \in \mathbb{N}, k \in \{1, \dots, K\}).$$

743 which goes to 0 as  $i$  goes to  $\infty$ . Hence, taking the limit  $i \rightarrow \infty$ , we conclude that

$$0 = \Delta \mathfrak{J}_k(w^\infty) \quad (k \in \{1, \dots, K\}).$$

744 This ensures, using the definition of  $\Delta \mathfrak{J}_k$  and Theorem A.1, that  $w^\infty$  is the minimizer of  $\mathfrak{J}$ . Note  
 745 that the existence and uniqueness of the minimizer of  $\mathfrak{J}$  is ensured by the coercivity and strict  
 746 convexity of  $\mathfrak{J}$ .

747 Finally, the uniqueness of minimizers, the coercivity of  $\mathfrak{J}$ , and inequality (A.1) lead to  
 748  $\lim_{i \rightarrow \infty} w^i = w^\infty$ .  $\square$

## 749 B Some unreachable states

750 In this paragraph, we will show, in the case  $m = 1$ , that for every  $k \in \mathbb{N}$ ,  $\theta \in \Theta \mapsto A(\theta)^k B(\theta) \in \mathbb{R}^n$   
 751 is unreachable for the system (2.1). To get this result, we will assume that the system (2.1) is  
 752 uniformly ensemble controllable, and that  $\Theta$  has a nonempty interior. Let us first define the set of  
 753 reachable points in time  $T > 0$ ,

$$\mathcal{R}_{A,B}(T) = \left\{ \theta \in \Theta \mapsto \int_0^T e^{(T-t)A(\theta)} B(\theta) u(t) dt \mid u \in L^2(0, T)^m \right\}.$$

754 Before entering the proof of this result, let us mention that (2.1) is uniformly ensemble control-  
 755 lable if and only if  $\text{Span} \{ \theta \in \Theta \mapsto A(\theta)^k B(\theta) v \mid k \in \mathbb{N}, v \in \mathbb{R}^m \}$  is dense in  $\mathcal{C}(\Theta)$ . This fact can  
 756 be found in [28] (see also [11]).

757 Let us first prove our result in the case  $n = 1$ .

758 **Proposition B.1.** *Let  $\Theta \subset \mathbb{R}$  be a compact set and assume that there exists  $\theta_0 \in \Theta$  such that*  
 759  *$\theta_0$  is an accumulation point of  $\Theta$ . Let  $a, b \in \mathcal{C}(\Theta)$  and assume that  $(a, b)$  is uniformly ensemble*  
 760 *controllable. Then for every  $T > 0$ , and every  $k \in \mathbb{N}$ ,  $a^k b \notin \mathcal{R}_{a,b}(T)$ .*

761 *Proof.* Let us set  $\tilde{\Theta} = a(\Theta)$ , we have  $\tilde{\Theta} \subset \mathbb{R}$  is a compact set,  $\tilde{\theta}_0 = a(\theta_0) \in \tilde{\Theta}$  is an accumulation  
 762 point of  $\tilde{\Theta}$ , and we set  $\iota_{\tilde{\Theta}}$  the identity map on  $\tilde{\Theta}$ . Since the pair  $(a, b)$  is ensemble controllable, we  
 763 necessarily have that  $a$  is injective (see [11, Proposition 4]) and  $b(\theta) \neq 0$  for every  $\theta \in \Theta$  (see Re-  
 764 mark 2.5). This ensures that if  $y \in \mathcal{R}_{a,b}(T)$  then  $y \circ a^{-1} \in \mathcal{R}_{\iota_{\tilde{\Theta}}, b \circ a^{-1}}(T) = (b \circ a^{-1}) \mathcal{R}_{\iota_{\tilde{\Theta}}, 1}(T)$ .  
 765 Consequently, it is enough to prove this result for  $a = \iota_{\tilde{\Theta}}$  and  $b = 1$ .

766 Let us assume by contradiction that there exists  $k \in \mathbb{N}$  and  $u \in L^2(0, T)$  such that  
 767  $\int_0^T e^{(T-t)\theta} u(t) dt = \theta^k$  for every  $\theta \in \Theta$ . Since  $\Theta$  admits an accumulation point, by analytic  
 768 continuation, we have

$$\int_0^T e^{(T-t)\theta} u(t) dt = \theta^k \quad (\theta \in \mathbb{R}).$$

769 For every  $\ell \in \mathbb{N}$ , we also have by derivation with respect to  $\theta$ ,

$$\int_0^T (T-t)^{k+1+\ell} e^{(T-t)\theta} u(t) dt = 0 \quad (\theta \in \mathbb{R}).$$

770 Finally, using Stone-Weierstrass Theorem, we conclude that

$$(T-t)^{k+1} e^{(T-t)\theta} u(t) = 0 \quad (\theta \in \mathbb{R}, t \in [0, T] \text{ a.e.})$$

771 and using the fact that  $(T-t)^{k+1} e^{(T-t)\theta}$  does not vanish on  $(0, T)$ , we conclude that  $u = 0$ . This  
 772 leads to a contradiction with  $\int_0^T e^{(T-t)\theta} u(t) dt = \theta^k$ .  $\square$

773 Even if the above result is given in the real context, it can be easily extended to the complex  
774 case (i.e., with  $a(\theta), b(\theta) \in \mathbb{C}$ , complex controls and  $\Theta \subset \mathbb{C}$ ). Its complex version will be used in  
775 the next result.

776 We are now in position to give the general result.

777 **Proposition B.2.** *Let  $\Theta \subset \mathbb{R}$  be a compact set and assume that  $\text{int } \Theta \neq \emptyset$ . Let  $A \in \mathcal{C}(\Theta)^{n \times n}$  and  
778  $B \in \mathcal{C}(\Theta)^n$ , and assume that  $(A, B)$  is uniformly ensemble controllable. Then for every  $T > 0$  and  
779 every  $k \in \mathbb{N}$ , we have*

$$A^k B \notin \mathcal{R}_{A,B}(T).$$

780 *Proof.* Following [10, Lemma A.2], there exists  $\theta_0 \in \text{int } \Theta$ ,  $\delta > 0$ ,  $\lambda \in \mathcal{C}([\theta_0 - \delta, \theta_0 + \delta], \mathbb{C})$  and  
781  $v \in \mathcal{C}([\theta_0 - \delta, \theta_0 + \delta], \mathbb{C})^n$  such that  $v(\theta) \neq 0$  and  $A(\theta)^* v(\theta) = \lambda(\theta)v(\theta)$  for every  $\theta \in [\theta_0 - \delta, \theta_0 + \delta]$ .  
782 We now notice that if the system (2.1) is ensemble controllable on  $\Theta$ , then the system given by the  
783 pair  $(\lambda, v^* B)$  is also uniformly ensemble controllable on  $[\theta_0 - \delta, \theta_0 + \delta]$ . According to Proposition B.1,  
784 we have that  $\lambda^k v^* B \notin \mathcal{R}_{\lambda, v^* B}(T)$  for every  $T > 0$ . This clearly ensures that  $A^k B \notin \mathcal{R}_{A,B}(T)$  for  
785 every  $T > 0$ .  $\square$

786 **Remark B.3.** Let us mention that in the case  $m > 1$ , the situation is not so trivial.  
787 For instance, let  $T > 0$ ,  $v \in \mathcal{C}^\infty([0, T])$  such that  $v^{(k)}(0) = v^{(k)}(T) = 0$  for every  $k \in \mathbb{N}$ , and such  
788 that  $\int_0^T e^{(T-t)\theta} v(t) dt \neq 0$ , and define  $\beta(\theta) = \left( \int_0^T e^{(T-t)\theta} v(t) dt \right)^{-1}$ . For every  $k \in \mathbb{N}$ , we have  
789  $\theta^k \in \mathcal{R}_{\theta, (1/\beta)}(T)$ . In fact, we clearly have  $1 = \int_0^T e^{(T-t)\theta} \beta(\theta) v(t) dt$ , from this relation and by  
790 integration by parts, one can deduce that  $\theta^k = \int_0^T e^{(T-t)\theta} \beta(\theta) v^{(k)}(t) dt$  for every  $k \in \mathbb{N}$ .

## 791 References

- 792 [1] A. Agrachev, Y. Baryshnikov, and A. Sarychev. Ensemble controllability by Lie algebraic  
793 methods. *ESAIM, Control Optim. Calc. Var.*, 22(4):921–938, 2016.
- 794 [2] H. Bauer. *Measure and integration theory. Transl. from the German by Robert B. Burckel*,  
795 volume 26 of *De Gruyter Stud. Math.* Berlin: de Gruyter, 2001.
- 796 [3] K. Beauchard, J.-M. Coron, and P. Rouchon. Controllability issues for continuous-spectrum  
797 systems and ensemble controllability of Bloch equations. *Commun. Math. Phys.*, 296(2):525–  
798 557, 2010.
- 799 [4] A. Becker and T. Bretl. Approximate steering of a unicycle under bounded model perturbation  
800 using ensemble control. *IEEE Transactions on Robotics*, 28(3):580–591, 2012.
- 801 [5] J. M. Borwein and Q. J. Zhu. *Techniques of variational analysis*, volume 20 of *CMS Books*  
802 *Math./Ouvrages Math. SMC*. New York, NY: Springer, 2005.
- 803 [6] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext.  
804 Springer, New York, 2011.
- 805 [7] E. J. Candès, J. Romberg, and T. Tao. Robust uncertainty principles: exact signal reconstruc-  
806 tion from highly incomplete frequency information. *IEEE Trans. Inform. Theory*, 52(2):489–  
807 509, 2006.
- 808 [8] X. Chen. Controllability issues of linear ensemble systems over multi-dimensional parameter-  
809 ization spaces, 2020. preprint.

- 810 [9] B. Danhane and J. Lohéac. Ensemble controllability of parabolic type equations. *Syst. Control*  
811 *Lett.*, 183:10, 2024. Id/No 105683.
- 812 [10] B. Danhane, J. Lohéac, and M. Jungers. Conditions for uniform ensemble output control-  
813 lability, and obstruction to uniform ensemble controllability. *Math. Control Relat. Fields*,  
814 14(3):1128–1175, 2024.
- 815 [11] G. Dirr and M. Schönlein. Uniform and  $L^q$ -ensemble reachability of parameter-dependent  
816 linear systems. *J. Differ. Equations*, 283:216–262, 2021.
- 817 [12] I. Ekeland and R. Témam. *Convex analysis and variational problems.*, volume 28 of *Classics*  
818 *Appl. Math.* Philadelphia, PA: Society for Industrial and Applied Mathematics, unabridged,  
819 corrected republication of the 1976 English original edition, 1999.
- 820 [13] C. Esteve, B. Geshkovski, D. Pighin, and E. Zuazua. Large-time asymptotics in deep learning,  
821 2021.
- 822 [14] G. B. Folland. *Real analysis.* Pure and Applied Mathematics (New York). John Wiley &  
823 Sons, Inc., New York, second edition, 1999. Modern techniques and their applications, A  
824 Wiley-Interscience Publication.
- 825 [15] P. A. Guth, K. Kunisch, and S. S. Rodrigues. Tracking optimal feedback control under  
826 uncertain parameters. *Phys. D*, 467:Paper No. 134245, 13, 2024.
- 827 [16] U. Helmke and M. Schönlein. Uniform ensemble controllability for one-parameter families of  
828 time-invariant linear systems. *Syst. Control Lett.*, 71:69–77, 2014.
- 829 [17] E. Kreindler and P. Sarachik. Constrained linear-quadratic optimization problems with  
830 parameter-dependent entries. *Journal of Optimization Theory and Applications*, 198(2):781–  
831 804, 2023.
- 832 [18] M. Lazar and J. Lohéac. Chapter 8 - control of parameter dependent systems. In E. Trélat  
833 and E. Zuazua, editors, *Numerical Control: Part A*, volume 23 of *Handbook of Numerical*  
834 *Analysis*, pages 265–306. Elsevier, 2022.
- 835 [19] M. Lazar and C. Molinari. Optimal distributed control of linear parabolic equations by spectral  
836 decomposition. *Optimal Control Appl. Methods*, 42(4):891–926, 2021.
- 837 [20] J.-S. Li and N. Khaneja. Control of inhomogeneous quantum ensembles. *Phys. Rev. A*,  
838 73:030302, Mar 2006.
- 839 [21] J.-S. Li and J. Qi. Ensemble control of time-invariant linear systems with linear parameter  
840 variation. *IEEE Trans. Autom. Control*, 61(10):2808–2820, 2016.
- 841 [22] Y. Li and S. Osher. Coordinate descent optimization for  $\ell^1$  minimization with application to  
842 compressed sensing; a greedy algorithm. *Inverse Probl. Imaging*, 3(3):487–503, 2009.
- 843 [23] W. Miao, G. Cheng, and J.-S. Li. On uniform ensemble controllability of diagonalizable linear  
844 ensemble systems. preprint, 2021.
- 845 [24] J. Peypouquet. *Convex optimization in normed spaces. Theory, methods and examples.*  
846 SpringerBriefs Optim. Cham: Springer, 2015.
- 847 [25] M. Schönlein. Computation of open-loop inputs for uniformly ensemble controllable systems.  
848 *Math. Control Relat. Fields*, 12(3):813–829, 2022.

- 849 [26] M. Schönlein. Polynomial methods to construct inputs for uniformly ensemble reachable linear  
850 systems. *Math. Control Signals Syst.*, 36(2):251–296, 2024.
- 851 [27] M. Schönlein and U. Helmke. Controllability of ensembles of linear dynamical systems. *Math.*  
852 *Comput. Simul.*, 125:3–14, 2016.
- 853 [28] R. Triggiani. Controllability and observability in Banach space with bounded operators. SIAM  
854 *J. Control* 13, 462–491 (1975)., 1974.
- 855 [29] E. Zuazua. Controllability and observability of partial differential equations: some results  
856 and open problems. In *Handbook of differential equations: evolutionary equations. Vol. III*,  
857 *Handb. Differ. Equ.*, pages 527–621. Elsevier/North-Holland, Amsterdam, 2007.