



Preparation of nanoparticle based aerogel scintillators for real-time radioactive gas detection

Yannis Cheref, Raphaël Marie-Luce, Pavlo Mai, Frédéric Lerouge, Sylvie Pierre, Benoit Sabot, Frédéric Chaput, Christophe Dujardin

► To cite this version:

Yannis Cheref, Raphaël Marie-Luce, Pavlo Mai, Frédéric Lerouge, Sylvie Pierre, et al.. Preparation of nanoparticle based aerogel scintillators for real-time radioactive gas detection. 22nd international Sol-Gel conference, Sep 2024, Berlin (Germany), Germany. p229. hal-04863394

HAL Id: hal-04863394

<https://hal.science/hal-04863394v1>

Submitted on 23 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nanoparticle-based scintillating aerogels for real-time radioactive gas detection

Tuesday, 3rd September - 17:45: Nanoparticles synthesis, self-assembly - Oral

Dr. Yannis Cheref¹, Mr. Raphaël Marie-Luce², Mr. Pavlo Mai³, Dr. Frédéric Lerouge², Mrs. Sylvie Pierre⁴, Dr. Benoit Sabot⁴, Dr. Frédéric Chaput², Prof. Christophe Dujardin³

1. ENS Lyon and ILM, Université Claude Bernard Lyon 1, CNRS, **2.** ENS Lyon, Université Claude Bernard Lyon 1, CNRS, **3.** ILM, Université Claude Bernard Lyon 1, CNRS, **4.** LIST, Laboratoire National Henri Becquerel, Université Paris-Saclay, CEA

The measurement of pure β emitting radioactive gases such as ${}^3\text{H}$ and ${}^{85}\text{Kr}$ is already of major importance for the nuclear safety authorities, and will meet an increasing demand with the expansion of nuclear-based energy production. Due to the short penetration length of β electrons in air, gaseous β emitters must be mixed with radiosensitive elements to enable detection. These are either gas-gas mixtures in an ionization chamber, or gas-liquid mixtures in liquid scintillation. However, none of these existing methods combine real-time analysis, sensitivity to multiple gases, and ease of deployment for on-site measurements.

We demonstrate a new gas-solid mixture approach using inorganic aerogels as nanoporous scintillators. This method combines the above criteria, allowing efficient real-time measurement of ${}^3\text{H}$ and ${}^{85}\text{Kr}$ activity. This presentation will focus on the preparation of scintillating aerogels, from the 10g-scale synthesis of scintillating nanoparticles to supercritical drying of nanoparticle gels into aerogels. This preparation method has been successfully applied to different scintillating nanomaterials.