

**Supplemental Table 1. Characteristics of studies included in this analysis from the INHANCE consortium pooled data version 1.5, by region and information on caffeinated coffee, decaffeinated coffee, or tea consumption**

<u>Study Location</u>	<u>Recruitment Period</u>	<u>Source</u>	<u>Cases</u> (n=9,548)	<u>Controls</u> (n=15,783)	<u>Matched Factors</u>	<u>Information Collected</u>		
						Caffeinated Coffee	Decaffeinated Coffee	Tea
<b>Europe</b>								
Milan, Italy	1984-1989	Hospital-based	416	1,531	---	Yes	Yes	Yes
France	1987-1992	Hospital-based	323	234	age, sex, hospital	Yes	No	Yes
Italy Multicenter	1990-1999	Hospital-based	1,261	2,716	---	Yes	Yes	Yes
Switzerland	1991-1997	Hospital-based	516	883	---	Yes	Yes	Yes
Saarland, Germany*	2001-2003	Population-based	94	94	age, sex	Yes	No	Yes
France Multicenter (Paris)*	2001-2007	Population-based	2,237	3,555	age, sex, region	Yes	No	Yes
Milan, Italy*	2006-2009	Hospital-based	370	755	---	Yes	Yes	Yes
<b>North America</b>								
Los Angeles, CA	1999-2004	Population-based	417	1,005	age, sex, neighborhood	Yes	Yes	Yes
Boston, MA	1999-2003	Population-based	1,034	1,251	age, sex, neighborhood	Yes	Yes	Yes
US Multicenter	1983-1984	Population-based	1,114	1,268	age, sex, race/ethnicity	Yes	No	Yes
Memorial Sloan-Kettering Cancer Center	1992-1994	Hospital-based	134	169	age, sex	Yes	Yes	Yes
Seattle, WA*	1983-1987	Population-based	650	547	age, sex	Yes	Yes	Yes
Buffalo, NY*	1982-1998	Hospital-based	632	1,254	age, sex	Yes	Yes	Yes
<b>Latin America</b>								
Puerto Rico	1992-1995	Population-based	350	521	age, sex	Yes	No	Yes
*Additional studies added for this updated pooled analysis								