



HAL
open science

Characteristics and Popularity of Videos of Abusive Head Trauma Prevention: Systematic Appraisal

Luc Goethals, Victoria Prokofieva Nelson, Fabien Fenouillet, Karine Chevreul, Manon Bergerat, Christine Lebreton, Yacine Refes, Flora Blangis, Martin Chalumeau, Enora Le Roux

► To cite this version:

Luc Goethals, Victoria Prokofieva Nelson, Fabien Fenouillet, Karine Chevreul, Manon Bergerat, et al.. Characteristics and Popularity of Videos of Abusive Head Trauma Prevention: Systematic Appraisal. Journal of Medical Internet Research, 2024, 26, pp.e60530. 10.2196/60530 . hal-04872565

HAL Id: hal-04872565

<https://hal.science/hal-04872565v1>

Submitted on 8 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Original Paper

Characteristics and Popularity of Videos of Abusive Head Trauma Prevention: Systematic Appraisal

Luc Goethals^{1,2}, PhD; Victoria Prokofieva Nelson³, PhD; Fabien Fenouillet³, PhD; Karine Chevreul^{2,4}, MD, PhD; Manon Bergerat^{1,5}, MD; Christine Lebreton⁶, MSc; Yacine Refes¹, PhD; Flora Blangis^{1,5}, PhD; Martin Chalumeau^{1,5*}, MD, PhD; Enora Le Roux^{2,7*}, PhD

¹Obstetrical, Perinatal and Pediatric Epidemiology Research team (EPOPé), Centre of Research in Epidemiology and Statistics (CRESS), Université Paris Cité, Inserm, Paris, France

²Epidémiologie clinique-évaluation économique appliqué aux populations vulnérables (ECEVE), Université Paris Cité, Inserm, Paris, France

³Learning, Physical Activity and Health, Interdisciplinary Laboratory in Neuroscience, Physiology and Psychology, Université Paris Nanterre, Nanterre, France

⁴Kastafiore, Paris, France

⁵Child protection unit, Department of General Pediatrics, Necker-Enfants malades hospital, Assistance Publique des Hôpitaux de Paris (AP-HP), Université Paris Cité, Paris, France

⁶Synakène, Ivry-sur-Seine, France

⁷Unit of Clinical Epidemiology (CIC-EC 1426), Hôpital Universitaire Robert-Debré, Assistance Publique des Hôpitaux de Paris (AP-HP), Inserm, Paris, France

* these authors contributed equally

Corresponding Author:

Luc Goethals, PhD

Obstetrical, Perinatal and Pediatric Epidemiology Research team (EPOPé), Centre of Research in Epidemiology and Statistics (CRESS)

Université Paris Cité

Inserm

123 Bd de Port-Royal

Paris, 75014

France

Phone: 33 0699222119

Email: luc.goethals@inserm.fr

Abstract

Background: Numerous strategies for preventing abusive head trauma (AHT) have been proposed, but controlled studies failed to demonstrate their effectiveness. Digital tools may improve the effectiveness of AHT prevention strategies by reaching a large proportion of the adult population.

Objective: This study aimed to describe the characteristics of videos of AHT prevention published on the internet, including their quality content, and to study their association with popularity.

Methods: From a systematic appraisal performed in June 2023, we identified videos addressing the primary prevention of AHT in children younger than 2 years that were published in English or French on the internet by public organizations or mainstream associations. We analyzed the characteristics of the videos; their quality with the Global Quality Scale (GQS); and their association with an index of popularity, the Video Power Index, using multivariable quasi-Poisson modeling.

Results: We included 53 (6.6%) of the 804 videos identified. Videos were mainly published by public organizations (43/53, 81%). The median time spent on the web was 6 (IQR 3-9) years, the median length was 202 (IQR 94-333) seconds, and the median GQS score was 4 (IQR 3-4). Infants were often depicted (42/53, 79%), including while crying (35/53, 66%) and being shaken (21/53, 40%). The characterization of shaking as an abuse and its legal consequences were cited in 47% (25/53) and 4% (2/53) of videos, respectively. The main prevention strategies in the videos were to raise awareness of the noxious outcome of shaking (49/53, 93%) and convince viewers of the effectiveness of coping strategies for infants' cries (45/53, 85%). The Video Power Index was positively correlated with the GQS ($r=0.38$; $P=.007$) and was independently associated with depicting an infant being shaken ($P=.03$; $\beta=1.74$, 95% CI 1.06-2.85) and the use of text or headers ($P=.04$; $\beta=2.15$, 95% CI 1.08-4.26).

Conclusions: AHT prevention videos had high quality but did not frequently deal with parental risk factors. The characteristics identified as being associated with the popularity of AHT prevention videos could help improve the impact of future prevention programs by enhancing their popularity.

(*J Med Internet Res* 2024;26:e60530) doi: [10.2196/60530](https://doi.org/10.2196/60530)

KEYWORDS

abusive head trauma; child physical abuse; shaken baby syndrome; SBS; primary prevention; web-based videos; digital tools; head trauma; prevention; video; internet; infant; mortality; morbidity; parent; caregivers; communication

Introduction

Abusive head trauma (AHT), the most severe form of child physical abuse, is defined as “an injury to the skull or intracranial contents of an infant or young child due to inflicted blunt impact and/or violent shaking” [1]. AHT most often occurs in infants younger than 2 years, at a mean age of 2 to 4 months [2-5], and its incidence has been estimated at 14 to 34 per 100,000 infants per year in countries with developed economies (CDE) [6-9]. Perpetrators of AHT are most often adult men acting alone [10]. Persistent crying is reported as the primary trigger for most AHT perpetration [11]. AHT primary prevention is pivotal because its mortality rate ranges from 7% to 26% [12-14], and nonlethal forms are associated with severe long-term morbidity, notably neurodevelopmental impairment [15].

Numerous AHT prevention strategies have been proposed, targeting the public, parents, or caregivers of children through mass communication campaigns (brochures, television, and radio) or specific programs that mainly rely on educating about the management of infants’ cries and the dangers of shaking a child [16,17]. These prevention programs also emphasize emotional regulation, particularly through the promotion of coping strategies when dealing with stress or frustration in front of infants’ persistent crying [16]. The effectiveness of these strategies has been evaluated in several controlled trials, and although some individual studies have reported reductions in AHT incidence [18-20], meta-analyses and the US Preventive Services Task Force have concluded no significant summed preventive effects [16,21-24]. This discrepancy may be due to differences in the study design used to measure the impact of the intervention and the level of proof used to define effectiveness [24]. More generally, despite all these efforts, the incidence of AHT is not declining in CDE, particularly in the wake of the COVID-19 pandemic and its prevention and mitigation measures [25,26]. Consequently, there is a pressing need for innovative approaches that can reach a wider population and lead to effective behavioral change among the parents and caregivers of tens of millions of newborns each year [27].

Public organizations increasingly diffuse prevention videos on wide-audience digital platforms to support and educate new parents [28]. This approach can reach a large part of the population and become an effective vector for health promotion and prevention [29]. Thus, internet videos could participate in the effective prevention of AHT [30,31]. Young adults are major consumers of internet videos [32] and frequently use the internet to search for health information [33,34]. Consequently, knowing the characteristics of AHT prevention videos that are associated

with a wide reach (ie, popularity) will help build more impactful public health interventions. Although measuring the direct impact of such videos on the incidence of AHT is complex and requires large-scale studies [35,36], understanding the factors that contribute to their popularity can serve as an intermediate step toward developing innovative prevention programs. The effectiveness of these programs could ultimately be evaluated by the reduction in AHT incidence. We aimed to describe the characteristics of videos of AHT prevention published on the internet, including their quality content, and to study their association with popularity.

Methods

Ethical Considerations

This study was determined to be exempt from review by the Research Project Ethics Evaluation Committee of the Robert Debré Hospital (IRB 00006477).

General Design

We performed a systematic appraisal of videos available on the internet whose main topic was the primary prevention of AHT in infants <2 years old. Because of the lack of methodological guidelines to conduct and report this type of systematic appraisal, we followed, after adaptation, the Center for Reviews and Dissemination guidance [37] and the PRISMA (Preferred Reporting Items for Systematic Review and Meta-Analysis) statement ([Multimedia Appendix 1](#)) [38].

Search Strategy and Selection Criteria

The search was conducted on June 29, 2023. We included all videos addressing the primary prevention [39] of AHT in children <2 years old that were published in English or French (arbitrary) on the internet by public organizations (medical or scientific societies, governments or agencies, hospitals) or mainstream associations in the field of child physical abuse. We excluded commercial videos, videos not specific to children <2 years old, duplicates, and videos proposed by professionals or individuals acting outside organizations or mainstream associations.

Two investigators (a public health researcher [LG] and an educational sciences researcher [VN]) systematically searched potentially eligible videos on official websites of the national pediatric and public health societies and ministries of health of CDE [40] and on the YouTube platform, the most-used website in the world to upload and share videos [41]. On YouTube, one investigator (LG) sequentially used the terms “child physical abuse,” “shaken baby syndrome,” “abusive head trauma,” and “crying baby” in English and French to identify videos. The

YouTube search engine was used in “incognito status,” with no filters, customized search methods, or personal log-in. The results were sorted by relevance according to the default YouTube search, and the first 100 videos of each search were extracted as suggested in the literature [42-45].

We first screened video titles and descriptions to identify and exclude content that was not specific to children under 2 years of age. We then evaluated each selected video in its entirety to determine its relevance to our study’s focus on AHT in infants, excluding commercial content.

Data Extraction and Collection Process

Because of the lack of guidelines to identify the variables to be extracted in the analysis of prevention videos on the internet, the authors established a list based on previous research syntheses of health videos [45-48], public health behavior theoretical models [49-52], and AHT literature [2,3,6,10,12,15,25,53]. This list was revised and enriched after the visualization of a sample of 5 eligible videos and then discussed among researchers until a consensus was reached.

The final data extraction form included 6 sections, 28 independent items, and 3 scales (Multimedia Appendices 2-4). The first section related to the general characteristics of the videos as follows: author (public organizations or associations); author’s country (United States, France, or other); video duration; persistence (years since web publication); and the number of views, likes, and dislikes. The second section related to video creation features, including the type of narrative style used (interview, monologue, or conversational), the use of storytelling, the use of metaphor, the type of language used (colloquial and formal), and the use of audiovisual elements (sound, music, image, photograph, text or headers, subtitles, or motion design). The third section related to the characters depicted in the videos: the presence of an infant, the infant’s age (younger than 6 months), whether the infant was crying or represented being shaken, the presence of the infant’s parents and family, the professionals (legal, health, or early childhood ones) and association members involved, and the perceived origin of these characters classified as suggested by the French regulation authority for audiovisual and numeric contents [54]. The fourth section related to AHT information delivered by the videos, including the risk factors for AHT related to the infant (male sex and disability) [55] and the parents (low socioeconomic level [56], parental vulnerability [2], and poor parent-infant bonding [57]), symptoms and medical consequences of AHT (temporary or long term), characterization of shaking as an abuse, and its penal consequences. The fifth

section was related to the perceived purpose of the video, which included the prevention strategies used (fear-appeal, persuasion of the severity of the depicted events, increasing the awareness of the noxious outcome of shaking, and persuasion of the effectiveness of coping strategies for infants’ cries) [49] and the perceived general outcomes searched (informational, educational, giving solution, call for action, or call to prevent from action). The sixth section included the Patient Education Materials Assessment Tool (PEMAT) Audio Visual A/V score [58,59] and the Global Quality Scale (GQS; Multimedia Appendix 4) [60].

Two investigators (LG and VN) extracted the data for each video independently. In case of disagreement between the 2 investigators, consensus was sought with the help of a third investigator (a public health researcher [EL]).

Analyses


First, we calculated the popularity of the videos by using the Video Power Index (VPI; Multimedia Appendix 4), as suggested [61]. Second, the Cohen κ value was calculated to measure the interinvestigator agreement on each extracted variable [62]. Third, we described the characteristics of the analyzed videos. Fourth, we studied the associations between the VPI and the videos’ characteristics and GQS and PEMAT scores using first, univariate tests and correlation analyses, and then a multivariable quasi-Poisson regression (due to the overdispersion of residuals in our regression model) including all variables statistically significant ($P < .05$) in univariate analyses. The GQS and PEMAT scores were not included in the multivariable model given the redundancy of their information (Multimedia Appendix 4) with some included variables. Hypothesis tests were 2-sided, with a priori level of significance of .05. R (version 4.3.0; R Foundation for Statistical Computing) was used for all statistical analysis [63].

Results

Results of the Search

Among the 804 videos identified by the search strategy, 53 (6.6%; Multimedia Appendix 5) met the inclusion criteria and were analyzed (Figure 1). All 53 videos were available on the YouTube platform; however, 2 (4%) had not been identified by our defined search on YouTube, but by searching the official websites of public organizations. The terms used to tag videos on the YouTube platform were “shaken baby syndrome” (42/53, 82%), “abusive head trauma” (11/53, 22%), and “crying baby” (2/53, 4%); none used the term “child physical abuse.”

Figure 1. PRISMA (Preferred Reporting Items for Systematic Review and Meta-Analysis) flowchart.



General Characteristics, Quality, and Popularity of Videos

Videos were published by public organizations (43/53, 81%) and associations (10/53, 19%) from the United States (34/53, 64%), France (11/53, 21%), or other countries (8/53, 15%; [Table 1](#)). They had a median length of 202 (IQR 94-333) seconds and had remained accessible via the web for a median of 6 years (IQR 3.3-9.3; [Table 2](#)). Videos accumulated a median of 7917 (IQR 1389-45,272) views, 21 (IQR 4-147) likes, and 2 (IQR 0-13) dislikes. The VPI was calculated for 48 (91%) of the 53 videos included due to missing data (number of likes and dislikes not available). The median VPI was 3 (IQR 1-14; mean 29, SD 113).

For the 28 video items rated, the median Cohen κ value was 0.65 (IQR 0.47-0.80; [Multimedia Appendix 3](#)). The videos included audiovisual elements such as text or headers (43/53, 81%), music (39/53, 74%), images (26/53, 49%), motion design (18/53, 34%), photographs (11/53, 21%), and subtitles (5/53,

9%; [Table 1](#)). Characters depicted were infants (42/53, 79%; various forms of presentation were used: real infants, photographs, motion design, and dolls), including while crying (35/53, 66%) and being shaken (21/53, 40%). The process of shaking was illustrated by motion design, the use of dolls, and metaphorical representations of an infant’s brain. Other characters included parents and family (36/53, 68%), and professionals (40/53, 75%). In 57% (30/53) of videos, these characters were from diverse origins. The cited AHT risk factors referred mainly to the infant (44/53, 83%) and then the parents (8/53, 15%). The characterization of shaking as an abuse and its penal consequences were mentioned in 47% (25/53) and 4% (2/53) of the videos, respectively. The main prevention strategies used in the videos aimed at increasing awareness of the noxious outcomes of shaking (49/53, 93%) and the persuasion of the effectiveness of coping strategies for infants’ cries (45/53, 85%).

The median understandability and actionability PEMAT scores were 83% (IQR 70%-89%) and 100% (IQR 67%-100%), respectively. The median GQS score was 4 (IQR 3-4; [Table 2](#)).

Table 1. Univariate associations between Video Power Index (VPI) and video characteristics.

Video characteristics	Videos (n=53), n (%)	VPI (n=48), median (IQR)	<i>P</i> value ^a
Video creation feature			
Author of the video			.41
Public organizations	43 (81)	3.0 (1.2-14.8)	
Association	10 (19)	3.9 (0.7-6.9)	
Author's country			.28
United States	34 (64)	6.6 (1.2-17.6)	
France	11 (21)	1.9 (1.6-3.2)	
Other	8 (15)	1.8 (0.9-5.7)	
Narrative style			
Interview			.96
Yes	20 (38)	4.8 (1.4-14.3)	
No	33 (62)	2.9 (1.1-13.6)	
Monologue			.02
Yes	45 (85)	3.9 (1.3-14.8)	
No	8 (15)	0.5 (0.2-1.6)	
Conversational			.41
Yes	14 (26)	7.3 (2.7-14.4)	
No	39 (74)	2.3 (1.1-12.4)	
Use of storytelling			.31
Yes	12 (23)	8.0 (2.2-14.5)	
No	41 (77)	2.7 (1.0-11.6)	
Use of metaphor			.12
Yes	4 (8)	35.1 (3.1-78.6)	
No	49 (92)	2.9 (1.1-13.1)	
Type of language used			
Colloquial			.22
Yes	41 (77)	3.9 (1.2-14.8)	
No	12 (19)	1.5 (0.7-7.0)	
Formal			.17
Yes	46 (87)	4.6 (1.2-14.8)	
No	7 (13)	1.9 (0.9-2.9)	
Audiovisual element			
Sound			.22
Yes	34 (64)	2.7 (0.9-10.5)	
No	19 (36)	4.6 (1.8-16.5)	
Music			.12
Yes	39 (74)	5.6 (1.3-14.4)	
No	14 (26)	1.5 (0.6-6.1)	
Image			.01
Yes	26 (49)	9.8 (2.8-23.7)	
No	27 (51)	1.7 (0.7-6.6)	

Video characteristics	Videos (n=53), n (%)	VPI (n=48), median (IQR)	P value ^a
Photograph			.02
Yes	11 (21)	14.5 (8.1-23.7)	
No	42 (79)	2.3 (0.8-8.0)	
Text or headers			.02
Yes	43 (81)	6.5 (1.5-15.6)	
No	10 (19)	1.1 (0.7-1.8)	
Subtitles			.94
Yes	5 (9)	3.6 (2.4-5.5)	
No	48 (91)	3.0 (1.1-14.8)	
Motion design			.003
Yes	18 (34)	14.8 (3.9-25.0)	
No	35 (66)	1.8 (0.8-6.6)	
Characters depicted in the videos			
Infant			.07
Yes	42 (79)	5.6 (1.3-14.8)	
No	11 (21)	1.5 (0.7-2.9)	
Infant's age younger than 6 months			.12
Yes	37 (70)	6.5 (1.3-14.8)	
No	16 (30)	1.7 (1.0-3.9)	
Representation of an infant being shaken			<.001
Yes	21 (40)	13.5 (5.4-25.2)	
No	32 (60)	1.8 (0.7-5.7)	
Representation of a crying infant			.28
Yes	35 (66)	6.5 (1.1-16.4)	
No	18 (34)	1.8 (1.3-5.1)	
Infant's parents and family			.47
Yes	36 (68)	6.5 (1.2-14.8)	
No	17 (32)	2.7 (0.9-5.6)	
Professional			.57
Yes	40 (75)	3.0 (1.2-14.8)	
No	13 (25)	3.1 (0.6-8.2)	
Perceived origin of the characters			.08
Only white	23 (43)	1.9 (0.8-6.6)	
Mixed origin	30 (57)	6.5 (1.3-22.9)	
AHT^b information delivered in the videos			
Infant risk factors			.19
Yes	44 (83)	1.9 (1.0-13.5)	
No	9 (17)	6.6 (2.7-16.5)	
Parent risk factors			.30
Yes	8 (15)	8.0 (3.9-17.7)	
No	45 (85)	2.7 (1.0-12.7)	

Video characteristics	Videos (n=53), n (%)	VPI (n=48), median (IQR)	<i>P</i> value ^a
Symptoms of AHT			.049
Yes	18 (34)	10.3 (2.6-22.3)	
No	35 (66)	1.9 (0.9-8.1)	
Temporary medical consequences of AHT			.11
Yes	25 (47)	7.3 (1.5-16.0)	
No	28 (53)	2.3 (0.8-7.7)	
Long-term medical consequences of AHT			.17
Yes	42 (80)	3.9 (1.2-16.0)	
No	11 (20)	1.8 (0.8-5.8)	
Characterization of shaking as abuse			.09
Yes	25 (47)	8.0 (2.7-14.8)	
No	28 (53)	1.8 (0.9-7.5)	
Penal consequences spelled out			.86
Yes	2 (4)	4.6 (4.6-4.6)	
No	51 (96)	3.0 (1.1-14.5)	
Video perceived purposes			
Prevention strategy used			
Fear-appeal, persuasion of severity of the depicted events			.14
Yes	26 (49)	6.6 (1.2-18.8)	
No	27 (51)	1.9 (1.1-8.0)	
Increasing the awareness of the noxious outcome of shaking			.06
Yes	49 (93)	3.9 (1.2-14.8)	
No	4 (7)	0.9 (0.0-2.1)	
Persuasion of the effectiveness of coping strategies for infants' cries			.68
Yes	45 (85)	3.0 (0.9-14.8)	
No	8 (15)	3.7 (2.3-7.9)	
Perceived general outcome searched			
Educational			.91
Yes	49 (93)	3.1 (1.1-14.8)	
No	4 (7)	2.7 (2.4-4.1)	
Informational			.37
Yes	50 (94)	3.1 (1.2-14.2)	
No	3 (6)	0.2 (0.1-56.6)	
Giving solution			.56
Yes	43 (81)	3.1 (1.1-14.8)	
No	10 (19)	2.7 (1.2-6.1)	
Call for action			.12
Yes	43 (81)	4.9 (1.2-16.0)	
No	10 (19)	2.0 (0.9-4.1)	
Call to prevent from action			.38
Yes	48 (91)	3.0 (1.2-14.8)	

Video characteristics	Videos (n=53), n (%)	VPI (n=48), median (IQR)	P value ^a
No	5 (9)	3.1 (0.7-6.6)	

^aMann-Whitney and Kruskal-Wallis tests.

^bAHT: abusive head trauma.

Table 2. General characteristics of the included videos and association with the Video Power Index (VPI).

Video features	Value (n=53), n (%)	Value (n=48), median (IQR)	r (n=48)	P value ^a (n=48)
Duration (seconds)	53 (100)	202 (94-333)	0.36	.01
Number of years accessible via the web	53 (100)	6 (3-9)	N/A ^b	N/A
Number of views	53 (100)	7917 (1389-45,272)	N/A	N/A
Number of likes	48 (91)	21 (4-147)	N/A	N/A
Number of dislikes	48 (91)	2 (0-13)	N/A	N/A
VPI	48 (91)	3 (1-14)	N/A	N/A
GQS ^c score	53 (100)	4 (3-4)	0.38	.007
PEMAT ^d understandability score (%)	53 (100)	83 (70-89)	0.27	.06
PEMAT actionability score (%)	53 (100)	100 (67-100)	0.27	.07

^aSpearman correlation test.

^bNot applicable.

^cGQS: Global Quality Scale.

^dPEMAT: Patient Education Materials Assessment Tool.

Determinants of Popularity

In the univariate analyses (Table 1), only monologue used as narrative style was significantly associated with increased VPI (median 3.9 vs 0.5 for no monologue; $P=.02$). The VPI was significantly increased when videos contained images (median 9.8 vs 1.7; $P=.01$), photographs (median 14.5 vs 2.3; $P=.02$), text or headers (median 6.5 vs 1.1; $P=.02$), and motion design (median 14.8 vs 1.8; $P=.003$) as audiovisual elements. The presence of an infant in the videos was not associated with the VPI. The VPI was significantly increased with the depiction of an infant being shaken (median 13.5 vs 1.8; $P<.001$). Character types represented in the video (infant's parents and family, professionals) and their perceived origin were not significantly associated with the VPI (all $P\geq.05$). The VPI was significantly

increased when symptoms of AHT were mentioned (median 10.3 vs 1.9; $P=.049$) but not when shaking was characterized as an abuse ($P=.09$) or when its penal consequences were stated ($P=.86$). None of the prevention strategies used in the videos and the perceived video's general outcomes were associated with increased VPI. The VPI was significantly positively correlated with the duration of the videos ($r=0.36$; $P=.01$) and the GQS score ($r=0.38$; $P=.007$) but not PEMAT scores (all $P\geq.05$).

In the multivariable analysis (Table 3), increased VPI was independently significantly associated only with the use of text or headers in the videos ($P=.04$; $\beta=2.15$, 95% CI 1.08-4.26) and the representation of an infant being shaken ($P=.03$; $\beta=1.74$, 95% CI 1.06-2.85).

Table 3. Multivariable analysis of the association of variables with Video Power Index (n=48).

Video features	β	95% CI	P value
Duration (seconds)	1.00	1.00-1.00	.60
Narrative style: monologue	2.44	0.98-6.10	.06
Audiovisual element			
Image	0.85	0.50-1.45	.55
Photograph	1.26	0.79-2.02	.34
Text or headers	2.15	1.08-4.26	.04
Motion design	1.00	0.58-1.75	.99
Representation of an infant being shaken	1.74	1.06-2.85	.03
Symptoms of abusive head trauma	1.35	0.85-2.13	.21

Discussion

Principal Findings

This systematic appraisal highlights the relationship between the characteristics and content of health web-based prevention videos and their popularity. It leads to practical implications for developing more effective prevention campaigns. In our study, VPIs were significantly increased for videos that included images, photographs, text or headers, and motion design. Although numerous studies examined the popularity of preventive health videos or health topics, none analyzed the relation between visual elements within the videos and their popularity [46,47,64-66]. In another educational field, instructional videos, one study had the same conclusion as ours on the determinants of popularity [67]. In our study, the representation of an infant being shaken was associated with increased VPI, a result providing additional evidence to support considering the use of shocking content on this topic. Shocking content is often used in prevention campaigns because it is supposed to better capture the audience's attention [68-70], but there is no consensus on its effectiveness [68-72]. Finally, in our study, as in other studies in various fields [73-75], the quality of the videos was positively correlated with their popularity.

In our study, the overall content of 53 videos of AHT prevention had good quality as indicated by a median GQS score of 4. This result can be explained in part by the videos being mainly published by public organizations, whose videos generally have higher GQS score than those published by nonprofessionals [48,64,76]. The videos we analyzed mostly contained information about the symptoms and medical consequences of AHT as well as the dangerous consequences of shaking an infant. This observation is consistent with the main content of AHT prevention interventions that do not use the internet [16,21]. The videos were mostly educational and used a strategy of persuasion about the effectiveness of coping, which is consistent with the strategies used in most AHT prevention interventions that attempt to convince parents of the efficacy of coping strategies in managing the response to infant crying (eg, walk away, leave the infant alone, or call a support person) [16,21]. However, some information was not addressed, particularly regarding parental risk factors, a finding similar to the content of other AHT prevention interventions [16,21]. By focusing solely on child risk factors and neglecting parental and

more generally perpetrator risk factors, the videos analyzed could induce a deleterious shift of the responsibility for shaking from the adults to the infant, a mechanism similar to victim blaming [77-79]. Including perpetrators' risk factors in prevention programs would help raise awareness of the personal responsibility of perpetrators and limit their ability to claim to be victims themselves, passive and will-less in their wrongdoing [78]. The characterization of shaking as an abuse and its penal consequences were often missing in the analyzed videos and were not found to be associated with popularity. The use of these arguments in prevention campaigns is debated. Emphasizing legal consequences might decrease audience participation in prevention campaigns, particularly because of stigma [80,81]. However, some studies suggest that education about the legality, consequences, and responsibilities associated with abusive behavior might enhance the target audience's understanding and commitment to preventing such behavior [82-84]. The optimal balance between legal deterrence and educational strategies should be adapted according to the objectives of the prevention campaigns.

The analyzed videos had a mean VPI of 29. There is no validated threshold to interpret a VPI value. The mean VPI in our study seems to be rather low compared with VPIs reported in the medical literature, which range from <1 (eg, 0.17 in health prevention videos) [46] to >1000 (eg, 1078 in chronic medical conditions videos) [47,64-66], and are lower when the publisher is a public organization [46,64,66]. The video platform that we used, YouTube, may itself use a recommendation algorithm that will not prioritize or suggest preventive AHT content because of potential violent content linked to the representation of a shaken infant. Furthermore, only 4% (2/53) of the analyzed videos resulted from the search term "crying baby," whereas crying is reported as the primary trigger in most perpetrations of AHT [11] and is probably the most frequent term used by parents rather than "shaken baby syndrome," the most frequent tag used in the analyzed videos. In the future, publishers should systematically tag their videos with the term "crying baby".

Limitations

The first limitation is that we included only videos published by public organizations in CDE and mainstream associations, on YouTube and on public organization websites, in English and French. As well, the search was restricted to the first 100

videos for each search term. The characteristics and popularity of videos posted on other popular platforms, such as Facebook, Instagram, and TikTok, and by other publishers may differ. Our findings, which are limited to videos using English and French languages, may not be generalized to all contexts. The direction and magnitude of the probable selection biases introduced by these criteria are unknown. Second, because of these strict selection criteria, the sample size resulted in limited statistical power to identify the determinants of VPI. Third, the median Cohen κ indicated only substantial agreement between investigators, thus highlighting the subjectivity of quoted criteria. Finally, although the GQS and VPI are increasingly used tools in the literature, they are not validated and there is no threshold for their interpretation.

Implications

As a first step to developing an effective AHT prevention program, this systematic appraisal of available videos suggests that developing high-quality content for AHT prevention videos can have a significant positive impact on their popularity. Future videos may also include specific features such as text or headers and the depiction of an infant being shaken to improve their popularity. However, popularity alone does not ensure the effectiveness of these videos in preventing AHT. Therefore, understanding popularity must be complemented by evaluating the effectiveness of the content and the needs and expectations of the target audience. Increasing popularity could lead to more impactful prevention programs.

Another implication is the need for the development and validation of methodological tools to scientifically conduct and

report health video synthesis research. Videos will probably become the main channel for public health programs in the coming years, and the standardization of methods is critical. Currently, there is an urgent need to validate or refine the PEMAT, GQS, and VPI as tools to evaluate the quality and popularity of videos. Furthermore, consensus is needed on core items that must be evaluated when analyzing videos. Methods to maximize the objectivity of the assessments should also be produced and made available to the research community.

By establishing an association between video quality and popularity, this research provides insights for the development of more effective digital prevention strategies that could be of use beyond AHT. The methodology developed to study the videos, in the absence of a recommended approach, could serve as a basis for further methodological developments that would improve research practices in this field.

Conclusion

Videos addressing the primary prevention of AHT in children <2 years old that were published on the internet in English or French by public organizations or mainstream associations were of high quality but did not frequently deal with parental risk factors. The inclusion of characteristics identified as associated with the popularity of AHT prevention videos could help improve the impact of future prevention programs by enhancing their popularity. Finally, there is an urgent need for the development and validation of methodological tools to standardize the conduct and report of studies on prevention videos.

Acknowledgments

The authors thank Pauline Scherdel, PhD, and Marianne Jacques, PhD student, of the Inserm for their technical advice on the statistical analysis of this study. They did not receive compensation for their contribution. This study was supported by a grant from the health sponsorship program of Mutuelles AXA. The funder had no role in the study design; the collection, analysis, and interpretation of data; the writing of the report; and the decision to submit the paper for publication.

Data Availability

The datasets generated and analyzed during this study are available from the corresponding author on reasonable request ([Multimedia Appendix 6](#)).

Authors' Contributions

LG contributed to conceptualization, methodology, validation, formal analysis, investigation, data curation, writing the original draft, reviewing and editing the manuscript, visualization, and project administration. VPN was involved in conceptualization, methodology, formal analysis, investigation, writing the original draft, and reviewing and editing the manuscript. FF contributed to conceptualization, methodology, formal analysis, writing the original draft, and reviewing and editing the manuscript. KC and MB reviewed and edited the manuscript. CL and YR contributed to conceptualization, methodology, writing the original draft, reviewing and editing the manuscript, and project administration. FB was responsible for conceptualization, methodology, software, formal analysis, writing the original draft, and reviewing and editing the manuscript. MC was involved in conceptualization, methodology, validation, formal analysis, resources, writing the original draft, reviewing and editing the manuscript, visualization, supervision, project administration, and securing funding. ELR contributed to conceptualization, methodology, software, validation, formal analysis, investigation, data curation, writing the original draft, reviewing and editing the manuscript, visualization, supervision, and project administration.

Conflicts of Interest

None declared.

Multimedia Appendix 1

PRISMA (Preferred Reporting Items for Systematic Review and Meta-Analysis) checklist.

[\[DOCX File , 21 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Metrics of the videos analyzed.

[\[DOCX File , 33 KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Video characteristics and their κ coefficient for investigator agreement.

[\[DOCX File , 20 KB-Multimedia Appendix 3\]](#)

Multimedia Appendix 4

Description of the Global Quality Scale (GQS), Patient Education Materials Assessment Tool Audio Visual (PEMAT) A/V scale, and Video Power Index (VPI) .

[\[DOCX File , 18 KB-Multimedia Appendix 4\]](#)

Multimedia Appendix 5

Videos included in the appraisal.

[\[DOCX File , 23 KB-Multimedia Appendix 5\]](#)

Multimedia Appendix 6

Data sharing statement.

[\[DOCX File , 13 KB-Multimedia Appendix 6\]](#)

References

1. Parks SE, Annett JL, Hill HA, Karch DL. Pediatric abusive head trauma : recommended definitions for public health surveillance and research. Centers for Disease Control and Prevention. URL: <https://stacks.cdc.gov/view/cdc/26243> [accessed 2023-05-09]
2. Rebbe R, Mienko JA, Martinson ML. Incidence and risk factors for abusive head trauma: a population-based study. *Child Abuse Rev.* 2020;29(3):195-207. [FREE Full text] [doi: [10.1002/car.2630](https://doi.org/10.1002/car.2630)] [Medline: [33071539](https://pubmed.ncbi.nlm.nih.gov/33071539/)]
3. Lee C, Barr RG, Catherine N, Wicks A. Age-related incidence of publicly reported shaken baby syndrome cases: is crying a trigger for shaking? *J Dev Behav Pediatr.* 2007;28(4):288-293. [doi: [10.1097/DBP.0b013e3180327b55](https://doi.org/10.1097/DBP.0b013e3180327b55)] [Medline: [17700080](https://pubmed.ncbi.nlm.nih.gov/17700080/)]
4. Paget LM, Gilard - Pioc S, Cottenet J, Beltzer N, Quantin C. Identification of infant victims of abusive head trauma hospitalised in France from 2015 to 2017. *Child Abuse Review.* 2020;29(3):182-194. [doi: [10.1002/car.2639](https://doi.org/10.1002/car.2639)]
5. Fujiwara T, Barr RG, Brant RF, Rajabali F, Pike I. Using international classification of diseases, 10th edition, codes to estimate abusive head trauma in children. *Am J Prev Med.* 2012;43(2):215-220. [doi: [10.1016/j.amepre.2012.04.028](https://doi.org/10.1016/j.amepre.2012.04.028)] [Medline: [22813688](https://pubmed.ncbi.nlm.nih.gov/22813688/)]
6. Keenan HT, Runyan DK, Marshall SW, Nocera MA, Merten DF, Sinal SH. A population-based study of inflicted traumatic brain injury in young children. *JAMA.* 2003;290(5):621-626. [doi: [10.1001/jama.290.5.621](https://doi.org/10.1001/jama.290.5.621)] [Medline: [12902365](https://pubmed.ncbi.nlm.nih.gov/12902365/)]
7. Emrick BB, Smith E, Thompson L, Mullett C, Pino E, Snyder K, et al. Epidemiology of abusive head trauma in West Virginia children <24 months: 2000-2010. *Child Abuse Negl.* 2019;93:215-221. [FREE Full text] [doi: [10.1016/j.chiabu.2019.05.011](https://doi.org/10.1016/j.chiabu.2019.05.011)] [Medline: [31125851](https://pubmed.ncbi.nlm.nih.gov/31125851/)]
8. Parrish J, Baldwin-Johnson C, Volz M, Goldsmith Y. Abusive head trauma among children in Alaska: a population-based assessment. *Int J Circumpolar Health.* 2013;72:eCollection 2013. [FREE Full text] [doi: [10.3402/ijch.v72i0.21216](https://doi.org/10.3402/ijch.v72i0.21216)] [Medline: [23986886](https://pubmed.ncbi.nlm.nih.gov/23986886/)]
9. Fanconi M, Lips U. Shaken baby syndrome in Switzerland: results of a prospective follow-up study, 2002-2007. *Eur J Pediatr.* Aug 2010;169(8):1023-1028. [doi: [10.1007/s00431-010-1175-x](https://doi.org/10.1007/s00431-010-1175-x)] [Medline: [20213304](https://pubmed.ncbi.nlm.nih.gov/20213304/)]
10. Edwards GA, Maguire SA, Gaither JR, Leventhal JM. What do confessions reveal about abusive head trauma? a systematic review. *Child Abuse Rev.* 2020;29(3):253-268. [FREE Full text] [doi: [10.1002/car.2627](https://doi.org/10.1002/car.2627)] [Medline: [37982093](https://pubmed.ncbi.nlm.nih.gov/37982093/)]
11. Barr RG, Trent RB, Cross J. Age-related incidence curve of hospitalized shaken baby syndrome cases: convergent evidence for crying as a trigger to shaking. *Child Abuse Negl.* 2006;30(1):7-16. [doi: [10.1016/j.chiabu.2005.06.009](https://doi.org/10.1016/j.chiabu.2005.06.009)] [Medline: [16406023](https://pubmed.ncbi.nlm.nih.gov/16406023/)]

12. Niederkrotenthaler T, Xu L, Parks SE, Sugerman DE. Descriptive factors of abusive head trauma in young children--United States, 2000-2009. *Child Abuse Negl.* 2013;37(7):446-455. [doi: [10.1016/j.chiabu.2013.02.002](https://doi.org/10.1016/j.chiabu.2013.02.002)] [Medline: [23535075](https://pubmed.ncbi.nlm.nih.gov/23535075/)]
13. Hobbs C, Childs AM, Wynne J, Livingston J, Seal A. Subdural haematoma and effusion in infancy: an epidemiological study. *Arch Dis Child.* 2005;90(9):952-955. [FREE Full text] [doi: [10.1136/adc.2003.037739](https://doi.org/10.1136/adc.2003.037739)] [Medline: [16113132](https://pubmed.ncbi.nlm.nih.gov/16113132/)]
14. Ferguson NM, Sarnaik A, Miles D, Shafi N, Peters MJ, Truemper E, et al. Abusive head trauma and mortality-an analysis from an international comparative effectiveness study of children with severe traumatic brain injury. *Crit Care Med.* 2017;45(8):1398-1407. [FREE Full text] [doi: [10.1097/CCM.0000000000002378](https://doi.org/10.1097/CCM.0000000000002378)] [Medline: [28430697](https://pubmed.ncbi.nlm.nih.gov/28430697/)]
15. Chevignard MP, Lind K. Long-term outcome of abusive head trauma. *Pediatr Radiol.* 2014;44 Suppl 4:S548-S558. [doi: [10.1007/s00247-014-3169-8](https://doi.org/10.1007/s00247-014-3169-8)] [Medline: [25501726](https://pubmed.ncbi.nlm.nih.gov/25501726/)]
16. Scott LJ, Wilson R, Davies P, Lyttle MD, Mytton J, Dawson S, et al. Educational interventions to prevent paediatric abusive head trauma in babies younger than one year old: a systematic review and meta-analyses. *Child Abuse Negl.* 2022;134:105935. [FREE Full text] [doi: [10.1016/j.chiabu.2022.105935](https://doi.org/10.1016/j.chiabu.2022.105935)] [Medline: [36308894](https://pubmed.ncbi.nlm.nih.gov/36308894/)]
17. Poole MK, Seal DW, Taylor CA. A systematic review of universal campaigns targeting child physical abuse prevention. *Health Educ Res.* 2014;29(3):388-432. [FREE Full text] [doi: [10.1093/her/cyu012](https://doi.org/10.1093/her/cyu012)] [Medline: [24711483](https://pubmed.ncbi.nlm.nih.gov/24711483/)]
18. Altman RL, Canter J, Patrick PA, Daley N, Butt NK, Brand DA. Parent education by maternity nurses and prevention of abusive head trauma. *Pediatrics.* 2011;128(5):e1164-e1172. [doi: [10.1542/peds.2010-3260](https://doi.org/10.1542/peds.2010-3260)] [Medline: [22025587](https://pubmed.ncbi.nlm.nih.gov/22025587/)]
19. Barr RG, Barr M, Rajabali F, Humphreys C, Pike I, Brant R, et al. Eight-year outcome of implementation of abusive head trauma prevention. *Child Abuse Negl.* 2018;84:106-114. [doi: [10.1016/j.chiabu.2018.07.004](https://doi.org/10.1016/j.chiabu.2018.07.004)] [Medline: [30077049](https://pubmed.ncbi.nlm.nih.gov/30077049/)]
20. Dias MS, Smith K, DeGuehery K, Mazur P, Li V, Shaffer ML. Preventing abusive head trauma among infants and young children: a hospital-based, parent education program. *Pediatrics.* 2005;115(4):e470-e477. [doi: [10.1542/peds.2004-1896](https://doi.org/10.1542/peds.2004-1896)] [Medline: [15805350](https://pubmed.ncbi.nlm.nih.gov/15805350/)]
21. Chang HY, Chang YC, Chang YT, Chen YW, Wu PY, Feng JY. The effectiveness of parenting programs in preventing abusive head trauma: a systematic review and meta-analysis. *Trauma Violence Abuse.* 2024;25(1):354-368. [doi: [10.1177/15248380231151690](https://doi.org/10.1177/15248380231151690)] [Medline: [36762510](https://pubmed.ncbi.nlm.nih.gov/36762510/)]
22. Viswanathan M, Rains C, Hart LC, Doran E, Sathe N, Hudson K, et al. Primary care interventions to prevent child maltreatment: evidence report and systematic review for the US preventive services task force. *JAMA.* 2024;331(11):959-971. [doi: [10.1001/jama.2024.0276](https://doi.org/10.1001/jama.2024.0276)] [Medline: [38502070](https://pubmed.ncbi.nlm.nih.gov/38502070/)]
23. Barry MJ, Nicholson WK, Silverstein M, Chelmsow D, Coker TR, Davis EM, et al. Primary care interventions to prevent child maltreatment: US preventive services task force recommendation statement. *JAMA.* 2024;331(11):951-958. [doi: [10.1001/jama.2024.1869](https://doi.org/10.1001/jama.2024.1869)] [Medline: [38502069](https://pubmed.ncbi.nlm.nih.gov/38502069/)]
24. Schilling S, Wood JN, Christian CW. Struggling to stem the tide of child maltreatment. *JAMA.* 2024;331(11):918-919. [FREE Full text] [doi: [10.1001/jama.2024.0899](https://doi.org/10.1001/jama.2024.0899)] [Medline: [38502084](https://pubmed.ncbi.nlm.nih.gov/38502084/)]
25. Lăzărescu AM, Benichi S, Blauwblomme T, Beccaria K, Bourgeois M, Roux CJ, et al. Abusive head trauma in infants during the COVID-19 pandemic in the Paris Metropolitan area. *JAMA Netw Open.* 2022;5(8):e2226182. [FREE Full text] [doi: [10.1001/jamanetworkopen.2022.26182](https://doi.org/10.1001/jamanetworkopen.2022.26182)] [Medline: [36040743](https://pubmed.ncbi.nlm.nih.gov/36040743/)]
26. Cercone DJ, Berger RP, Manole MD, Soung JK, Coombs CM, Noorbakhsh KA. Increased severity of abusive head trauma during the first year of the COVID-19 pandemic. *Child Abuse Negl.* 2023;135:105971. [FREE Full text] [doi: [10.1016/j.chiabu.2022.105971](https://doi.org/10.1016/j.chiabu.2022.105971)] [Medline: [36427395](https://pubmed.ncbi.nlm.nih.gov/36427395/)]
27. Total births by sex. United Nations, Department of Economic and Social Affairs, Population Division. URL: <https://tinyurl.com/4kwaw3yb> [accessed 2024-03-18]
28. Wiley M, Schultheis A, Francis B, Tiyyagura G, Leventhal JM, Rutherford HJV, et al. Parents' perceptions of infant crying: a possible path to preventing abusive head trauma. *Acad Pediatr.* 2020;20(4):448-454. [doi: [10.1016/j.acap.2019.10.009](https://doi.org/10.1016/j.acap.2019.10.009)] [Medline: [31629119](https://pubmed.ncbi.nlm.nih.gov/31629119/)]
29. Chatterjee A, Prinz A, Gerdes M, Martinez S. Digital interventions on healthy lifestyle management: systematic review. *J Med Internet Res.* 2021;23(11):e26931. [FREE Full text] [doi: [10.2196/26931](https://doi.org/10.2196/26931)] [Medline: [34787575](https://pubmed.ncbi.nlm.nih.gov/34787575/)]
30. Fujiwara T, Isumi A, Sampei M, Yamada F, Miyazaki Y. Effectiveness of using an educational video simulating the anatomical mechanism of shaking and smothering in a home-visit program to prevent self-reported infant abuse: a population-based quasi-experimental study in Japan. *Child Abuse Negl.* 2020;101:104359. [doi: [10.1016/j.chiabu.2020.104359](https://doi.org/10.1016/j.chiabu.2020.104359)] [Medline: [31945514](https://pubmed.ncbi.nlm.nih.gov/31945514/)]
31. Coster D. Changing parents' behaviour using a psycho - educational film as a preventative measure to reduce the risk of non - accidental head injury. *Child Abuse Rev.* 2017;26(6):465-476. [doi: [10.1002/car.2469](https://doi.org/10.1002/car.2469)]
32. Global time spent with online video 2023. Statista. URL: <https://www.statista.com/statistics/611707/online-video-time-spent/> [accessed 2023-09-28]
33. EU citizens: over half seek health information online. Eurostat. URL: <https://ec.europa.eu/eurostat/web/products-eurostat-news/-/edn-20220406-1> [accessed 2023-09-28]
34. Finney Rutten LJ, Blake KD, Greenberg-Worisek AJ, Allen SV, Moser RP, Hesse BW. Online health information seeking among US adults: measuring progress toward a healthy people 2020 objective. *Public Health Rep.* 2019;134(6):617-625. [doi: [10.1177/0033354919874074](https://doi.org/10.1177/0033354919874074)] [Medline: [31513756](https://pubmed.ncbi.nlm.nih.gov/31513756/)]

35. Dias MS, Rottmund CM, Cappos KM, Reed ME, Wang M, Stetter C, et al. Association of a postnatal parent education program for abusive head trauma with subsequent pediatric abusive head trauma hospitalization rates. *JAMA Pediatr.* 2017;171(3):223-229. [doi: [10.1001/jamapediatrics.2016.4218](https://doi.org/10.1001/jamapediatrics.2016.4218)] [Medline: [28135348](https://pubmed.ncbi.nlm.nih.gov/28135348/)]
36. Barr RG, Barr M, Fujiwara T, Conway J, Catherine N, Brant R. Do educational materials change knowledge and behaviour about crying and shaken baby syndrome? a randomized controlled trial. *CMAJ.* 2009;180(7):727-733. [FREE Full text] [doi: [10.1503/cmaj.081419](https://doi.org/10.1503/cmaj.081419)] [Medline: [19255065](https://pubmed.ncbi.nlm.nih.gov/19255065/)]
37. Centre for Reviews and Dissemination. *Systematic Reviews: CRD's Guidance for Undertaking Reviews in Health Care.* Layerthorpe, England. University of York; 2009.
38. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. [FREE Full text] [doi: [10.1136/bmj.n71](https://doi.org/10.1136/bmj.n71)] [Medline: [33782057](https://pubmed.ncbi.nlm.nih.gov/33782057/)]
39. Health promotion and disease prevention through population-based interventions, including action to address social determinants and health inequity. World Health Organization; Eastern Mediterranean Region. URL: <http://www.emro.who.int/about-who/public-health-functions/health-promotion-disease-prevention.html> [accessed 2023-07-27]
40. Country classifications. United Nations. 2022. URL: https://www.un.org/development/desa/dpad/wp-content/uploads/sites/45/WESP2022_ANNEX.pdf [accessed 2023-05-09]
41. Top websites ranking-most visited websites. Similar Web. URL: <https://www.similarweb.com/top-websites/> [accessed 2024-01-02]
42. Morra S, Collà Ruvolo C, Napolitano L, La Rocca R, Celentano G, Califano G, et al. YouTube as a source of information on bladder pain syndrome: a contemporary analysis. *Neurourol Urodyn.* 2022;41(1):237-245. [doi: [10.1002/nau.24802](https://doi.org/10.1002/nau.24802)] [Medline: [34559920](https://pubmed.ncbi.nlm.nih.gov/34559920/)]
43. Cilio S, Collà Ruvolo C, Turco C, Creta M, Capece M, La Rocca R, et al. Analysis of quality information provided by "Dr. YouTube™" on Phimosis. *Int J Impot Res.* 2022;35(4):398-403. [doi: [10.1038/s41443-022-00557-5](https://doi.org/10.1038/s41443-022-00557-5)] [Medline: [35332276](https://pubmed.ncbi.nlm.nih.gov/35332276/)]
44. Capece M, Di Giovanni A, Cirigliano L, Napolitano L, La Rocca R, Creta M, et al. YouTube as a source of information on penile prosthesis. *Andrologia.* 2022;54(1):e14246. [doi: [10.1111/and.14246](https://doi.org/10.1111/and.14246)] [Medline: [34519075](https://pubmed.ncbi.nlm.nih.gov/34519075/)]
45. Chen Z, Pan S, Zuo S. TikTok and YouTube as sources of information on anal fissure: a comparative analysis. *Front Public Health.* 2022;10:1000338. [doi: [10.3389/fpubh.2022.1000338](https://doi.org/10.3389/fpubh.2022.1000338)] [Medline: [36407987](https://pubmed.ncbi.nlm.nih.gov/36407987/)]
46. Kiss A, Soós S, Temesi Á, Unger-Plasek B, Lakner Z, Tompa O. Evaluation of the reliability and educational quality of YouTube™ videos on sport nutrition topics. *J Int Soc Sports Nutr.* 2023;20(1):2278632. [doi: [10.1080/15502783.2023.2278632](https://doi.org/10.1080/15502783.2023.2278632)] [Medline: [37953602](https://pubmed.ncbi.nlm.nih.gov/37953602/)]
47. Kwak D, Park JW, Won Y, Kwon Y, Lee JI. Quality and reliability evaluation of online videos on carpal tunnel syndrome: a YouTube video-based study. *BMJ Open.* 2022;12(4):e059239. [doi: [10.1136/bmjopen-2021-059239](https://doi.org/10.1136/bmjopen-2021-059239)] [Medline: [35428646](https://pubmed.ncbi.nlm.nih.gov/35428646/)]
48. Li HOY, Pastukhova E, Brandts-Longtin O, Tan MG, Kirchhof MG. YouTube as a source of misinformation on COVID-19 vaccination: a systematic analysis. *BMJ Glob Health.* Mar 2022;7(3):e008334. [doi: [10.1136/bmjgh-2021-008334](https://doi.org/10.1136/bmjgh-2021-008334)] [Medline: [35264318](https://pubmed.ncbi.nlm.nih.gov/35264318/)]
49. Rogers RW, Cacioppo JR, Petty RE. Cognitive and physiological processes in fear appeals and attitude change: a revised theory of protection motivation. In: *Social Psychology: A Source Book.* New York City, NY. Guilford Press; 1983:153-176.
50. Maddux JE, Rogers RW. A protection motivation theory of fear appeals and attitude change. *J Psychol.* 1975;91(1):93-114. [doi: [10.1080/00223980.1975.9915803](https://doi.org/10.1080/00223980.1975.9915803)] [Medline: [28136248](https://pubmed.ncbi.nlm.nih.gov/28136248/)]
51. Ajzen I, Fishbein M. *Understanding Attitudes and Predicting Social Behavior.* Englewood Cliffs, NJ. Prentice-Hall; 1980.
52. Mayer RE, Moreno R. Nine ways to reduce cognitive load in multimedia learning. *Educ Psychol.* 2010;38(1):43-52. [doi: [10.1207/S15326985EP3801_6](https://doi.org/10.1207/S15326985EP3801_6)]
53. Notrica DM, Kirsch L, Misra S, Kelly C, Greenberg J, Ortiz JB, et al. Evaluating abusive head trauma in children <5 years old: risk factors and the importance of the social history. *J Pediatr Surg.* 2021;56(2):390-396. [doi: [10.1016/j.jpedsurg.2020.10.019](https://doi.org/10.1016/j.jpedsurg.2020.10.019)] [Medline: [33220974](https://pubmed.ncbi.nlm.nih.gov/33220974/)]
54. Représentation de la diversité. Autorité de Régulation de la Communication Audiovisuelle et Numérique. URL: <https://www.arcom.fr/nous-connaître-nos-missions/garantir-le-pluralisme-et-la-cohesion-sociale/representation-de-la-diversite> [accessed 2024-01-22]
55. Sullivan PM, Knutson JF. Maltreatment and disabilities: a population-based epidemiological study. *Child Abuse Negl.* 2000;24(10):1257-1273. [doi: [10.1016/s0145-2134\(00\)00190-3](https://doi.org/10.1016/s0145-2134(00)00190-3)] [Medline: [11075694](https://pubmed.ncbi.nlm.nih.gov/11075694/)]
56. Judd N, Hughes K, Bellis MA, Hardcastle K, Amos R. Is parental unemployment associated with increased risk of adverse childhood experiences? a systematic review and meta-analysis. *J Public Health (Oxf).* 2023;45(4):829-839. [doi: [10.1093/pubmed/fdad069](https://doi.org/10.1093/pubmed/fdad069)] [Medline: [37253685](https://pubmed.ncbi.nlm.nih.gov/37253685/)]
57. Loiseau M, Cottenet J, François-Pursell I, Bechraoui-Quantin S, Jud A, Gilard-Pioc S, et al. Hospitalization for physical child abuse: associated medical factors and medical history since birth. *Child Abuse Negl.* 2023;146:106482. [doi: [10.1016/j.chiabu.2023.106482](https://doi.org/10.1016/j.chiabu.2023.106482)] [Medline: [37776729](https://pubmed.ncbi.nlm.nih.gov/37776729/)]
58. Shoemaker SJ, Wolf MS, Brach C. Development of the patient education materials assessment tool (PEMAT): a new measure of understandability and actionability for print and audiovisual patient information. *Patient Educ Couns.* 2014;96(3):395-403. [doi: [10.1016/j.pec.2014.05.027](https://doi.org/10.1016/j.pec.2014.05.027)] [Medline: [24973195](https://pubmed.ncbi.nlm.nih.gov/24973195/)]

59. Shoemaker S, Wolf M, Brach C. The Patient Education Materials Assessment Tool (PEMAT) and user's guide. Agency for Healthcare Research and Quality. URL: <https://www.ahrq.gov/health-literacy/patient-education/pemat.html> [accessed 2023-04-05]
60. Bernard A, Langille M, Hughes S, Rose C, Leddin D, Veldhuyzen van Zanten S. A systematic review of patient inflammatory bowel disease information resources on the world wide web. *Am J Gastroenterol*. 2007;102(9):2070-2077. [doi: [10.1111/j.1572-0241.2007.01325.x](https://doi.org/10.1111/j.1572-0241.2007.01325.x)] [Medline: [17511753](https://pubmed.ncbi.nlm.nih.gov/17511753/)]
61. Erdem MN, Karaca S. Evaluating the accuracy and quality of the information in kyphosis videos shared on YouTube. *Spine (Phila Pa 1976)*. 2018;43(22):E1334-E1339. [doi: [10.1097/BRS.0000000000002691](https://doi.org/10.1097/BRS.0000000000002691)] [Medline: [29664816](https://pubmed.ncbi.nlm.nih.gov/29664816/)]
62. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159. [doi: [10.2307/2529310](https://doi.org/10.2307/2529310)]
63. R Core Team (2023). *R Found Stat Comput Vienna Austria*. R: The R Project for Statistical Computing. URL: <https://www.r-project.org/> [accessed 2024-04-24]
64. Kunze KN, Krivicich LM, Verma NN, Chahla J. Quality of online video resources concerning patient education for the meniscus: a YouTube-based quality-control study. *Arthroscopy*. 2020;36(1):233-238. [doi: [10.1016/j.arthro.2019.07.033](https://doi.org/10.1016/j.arthro.2019.07.033)] [Medline: [31864582](https://pubmed.ncbi.nlm.nih.gov/31864582/)]
65. Etzel CM, Bokshan SL, Forster TA, Owens BD. A quality assessment of YouTube content on shoulder instability. *Phys Sportsmed*. 2022;50(4):289-294. [doi: [10.1080/00913847.2021.1942286](https://doi.org/10.1080/00913847.2021.1942286)] [Medline: [34121601](https://pubmed.ncbi.nlm.nih.gov/34121601/)]
66. Zhang X, Yang Y, Shen YW, Zhang KR, Ma LT, Ding C, et al. Quality of online video resources concerning patient education for neck pain: a YouTube-based quality-control study. *Front Public Health*. 2022;10:972348. [doi: [10.3389/fpubh.2022.972348](https://doi.org/10.3389/fpubh.2022.972348)] [Medline: [36211682](https://pubmed.ncbi.nlm.nih.gov/36211682/)]
67. Hove PT, Meij H. Like it or not. what characterizes YouTube's more popular instructional videos? *Semantic Scholar*. 2015. URL: <https://www.semanticscholar.org/paper/Like-It-or-Not.-What-Characterizes-YouTube's-More-Hove-Meij/db1038640b6f95675c66e2be533785e8f006d721> [accessed 2024-02-02]
68. Witte K, Allen M. A meta-analysis of fear appeals: implications for effective public health campaigns. *Health Educ Behav*. 2000;27(5):591-615. [doi: [10.1177/109019810002700506](https://doi.org/10.1177/109019810002700506)] [Medline: [11009129](https://pubmed.ncbi.nlm.nih.gov/11009129/)]
69. Ruiter RAC, Kessels LTE, Peters JGY, Kok G. Sixty years of fear appeal research: current state of the evidence. *Int J Psychol*. 2014;49(2):63-70. [doi: [10.1002/ijop.12042](https://doi.org/10.1002/ijop.12042)] [Medline: [24811876](https://pubmed.ncbi.nlm.nih.gov/24811876/)]
70. Tannenbaum MB, Hepler J, Zimmerman RS, Saul L, Jacobs S, Wilson K, et al. Appealing to fear: a meta-analysis of fear appeal effectiveness and theories. *Psychol Bull*. 2015;141(6):1178-1204. [doi: [10.1037/a0039729](https://doi.org/10.1037/a0039729)] [Medline: [26501228](https://pubmed.ncbi.nlm.nih.gov/26501228/)]
71. Dahl DW, Frankenberger KD, Manchanda RV. Does it pay to shock? reactions to shocking and nonshocking advertising content among university students. *J Adv Res*. 2003;43(03):268-280. [doi: [10.1017/s0021849903030332](https://doi.org/10.1017/s0021849903030332)]
72. Lull RB, Bushman BJ. Do sex and violence sell? a meta-analytic review of the effects of sexual and violent media and ad content on memory, attitudes, and buying intentions. *Psychol Bull*. 2015;141(5):1022-1048. [doi: [10.1037/bu10000018](https://doi.org/10.1037/bu10000018)] [Medline: [26191956](https://pubmed.ncbi.nlm.nih.gov/26191956/)]
73. Güvener O, Da' F, Sarikaya P. Popularity, quality and reliability of YouTube videos as an information resource on kinesio taping. *Dicle T'p Derg Dicle Üniversitesi*. 2023;50(1):22-29. [doi: [10.5798/dicletip.1266703](https://doi.org/10.5798/dicletip.1266703)]
74. Aydin CK, Gunec HG. Quality of information on YouTube about artificial intelligence in dental radiology. *J Dent Educ*. 2020;84(10):1166-1172. [doi: [10.1002/jdd.12362](https://doi.org/10.1002/jdd.12362)] [Medline: [32813894](https://pubmed.ncbi.nlm.nih.gov/32813894/)]
75. Nishizaki N, Hirano D, Oishi K, Shimizu T. YouTube videos in Japanese as a source of information on nocturnal enuresis: a content-quality and reliability analysis. *Pediatr Int*. 2022;64(1):e15049. [doi: [10.1111/ped.15049](https://doi.org/10.1111/ped.15049)] [Medline: [34747553](https://pubmed.ncbi.nlm.nih.gov/34747553/)]
76. Li HOY, Bailey A, Huynh D, Chan J. YouTube as a source of information on COVID-19: a pandemic of misinformation? *BMJ Glob Health*. 2020;5(5):e002604. [doi: [10.1136/bmjgh-2020-002604](https://doi.org/10.1136/bmjgh-2020-002604)] [Medline: [32409327](https://pubmed.ncbi.nlm.nih.gov/32409327/)]
77. Suarez E, Gadalla TM. Stop blaming the victim: a meta-analysis on rape myths. *J Interpers Violence*. 2010;25(11):2010-2035. [doi: [10.1177/0886260509354503](https://doi.org/10.1177/0886260509354503)] [Medline: [20065313](https://pubmed.ncbi.nlm.nih.gov/20065313/)]
78. Lamb S. *The Trouble with Blame: Victims, Perpetrators, and Responsibility*. Cambridge, MA. Harvard University Press; 1996.
79. Strömwall LA, Landström S, Alfredsson H. Perpetrator characteristics and blame attributions in a stranger rape situation. *Eur J Psychol Appl Leg Context*. 2014;6(2):63-67. [doi: [10.1016/j.ejpal.2014.06.002](https://doi.org/10.1016/j.ejpal.2014.06.002)]
80. Kemshall H, Moulden HM. Communicating about child sexual abuse with the public: learning the lessons from public awareness campaigns. *J Sex Aggress*. 2016;23(2):124-138. [doi: [10.1080/13552600.2016.1222004](https://doi.org/10.1080/13552600.2016.1222004)]
81. Beier KM, Neutze J, Mundt IA, Ahlers CJ, Goecker D, Konrad A, et al. Encouraging self-identified pedophiles and hebephiles to seek professional help: first results of the Prevention Project Dunkelfeld (PPD). *Child Abuse Negl*. 2009;33(8):545-549. [doi: [10.1016/j.chiabu.2009.04.002](https://doi.org/10.1016/j.chiabu.2009.04.002)] [Medline: [19765825](https://pubmed.ncbi.nlm.nih.gov/19765825/)]
82. Schober DJ, Fawcett SB, Bernier J. The enough abuse campaign: building the movement to prevent child sexual abuse in Massachusetts. *J Child Sex Abus*. 2012;21(4):456-469. [doi: [10.1080/10538712.2012.675423](https://doi.org/10.1080/10538712.2012.675423)] [Medline: [22809049](https://pubmed.ncbi.nlm.nih.gov/22809049/)]
83. Bailey A, Allen L, Stevens E, Dervley R, Findlater D, Wefers S. Pathways and prevention for indecent images of children offending: a qualitative study. *Sex Offending*. 2022;17:1-25. [doi: [10.5964/sotrap.6657](https://doi.org/10.5964/sotrap.6657)]

84. Newman EF, Efthymiadou E, Quayle E, Squire T, Denis D, Wortley R, et al. The impact of a public health campaign to deter viewing of child sexual abuse images online: a case study of the UK stop it now! campaign. *Sex Abuse*. 2024;36(6):635-661. [doi: [10.1177/10790632231205784](https://doi.org/10.1177/10790632231205784)] [Medline: [37837444](https://pubmed.ncbi.nlm.nih.gov/37837444/)]

Abbreviations

AHT: abusive head trauma

CDE: countries with developed economies

GQS: Global Quality Scale

PEMAT: Patient Education Materials Assessment Tool

PRISMA: Preferred Reporting Items for Systematic Review and Meta-Analysis

VPI: Video Power Index

Edited by F Xie; submitted 14.05.24; peer-reviewed by VJ Palusci, O Berthold, X Jiang; comments to author 17.07.24; revised version received 02.09.24; accepted 25.09.24; published 10.12.24

Please cite as:

Goethals L, Prokofieva Nelson V, Fenouillet F, Chevreur K, Bergerat M, Lebreton C, Refes Y, Blangis F, Chalumeau M, Le Roux E
Characteristics and Popularity of Videos of Abusive Head Trauma Prevention: Systematic Appraisal

J Med Internet Res 2024;26:e60530

URL: <https://www.jmir.org/2024/1/e60530>

doi: [10.2196/60530](https://doi.org/10.2196/60530)

PMID:

©Luc Goethals, Victoria Prokofieva Nelson, Fabien Fenouillet, Karine Chevreur, Manon Bergerat, Christine Lebreton, Yacine Refes, Flora Blangis, Martin Chalumeau, Enora Le Roux. Originally published in the Journal of Medical Internet Research (<https://www.jmir.org>), 10.12.2024. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research (ISSN 1438-8871), is properly cited. The complete bibliographic information, a link to the original publication on <https://www.jmir.org/>, as well as this copyright and license information must be included.