



Borrelia, Leishmania, and Babesia: An Emerging Triad of Vector-Borne Co-Infections?

Eliannte Piloto-Sardiñas, Ana Laura Cano-Argüelles, Alejandro Cabezas-Cruz

► To cite this version:

Eliannte Piloto-Sardiñas, Ana Laura Cano-Argüelles, Alejandro Cabezas-Cruz. Borrelia, Leishmania, and Babesia: An Emerging Triad of Vector-Borne Co-Infections?. *Pathogens*, 2025, 14 (1), pp.36. 10.3390/pathogens14010036 . hal-04873329

HAL Id: hal-04873329

<https://hal.science/hal-04873329v1>

Submitted on 8 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Editorial

Borrelia, Leishmania, and Babesia: An Emerging Triad of Vector-Borne Co-Infections?

Elianee Piloto-Sardiñas ^{1,2,*}, Ana Laura Cano-Argüelles ³ and Alejandro Cabezas-Cruz ^{2,*}

¹ Direction of Animal Health, National Center for Animal and Plant Health, Carretera de Tapaste y Autopista Nacional, Apartado Postal 10, San José de las Lajas 32700, Cuba

² ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale, F-94700 Maisons-Alfort, France

³ Parasitology Laboratory, Institute of Natural Resources and Agrobiology of Salamanca (IRNASA, CSIC), Cordel de Merinas, 40-52, 37008 Salamanca, Spain; ana.cano@irnasa.csic.es

* Correspondence: elianne9409@gmail.com (E.P.-S.); alejandro.cabezas@vet-alfort.fr (A.C.-C.)

Canine leishmaniosis (CanL), caused by the protozoan *Leishmania infantum* and transmitted primarily by phlebotomine sand flies, poses significant challenges for zoonotic disease management [1], with dogs serving as reservoirs, facilitating transmission to humans [2]. Host exposure to sand fly vectors, as well as ticks carrying other pathogens, increases the risk of co-infection with *Leishmania*, *Borrelia*, and *Babesia* [3]. These co-infections may exacerbate CanL progression due to synergistic interactions between pathogens that manipulate host immune responses [3].

The dynamics of zoonotic diseases are increasingly influenced by the overlapping habitats of vectors such as sand flies and ticks [4–6]. Climate change and habitat alterations are driving these vectors into new territories [7,8], creating conditions in which ticks (carrying *Borrelia* and *Babesia*) and sand flies (carrying *Leishmania*) can coexist [9–15]. This overlap enhances the likelihood of hosts, particularly dogs, becoming co-infected with multiple pathogens [9–16]. Pathogen interactions during co-infections—synergistic, antagonistic, or neutral—affect virulence, pathogenicity, and colonization success [17,18], affecting disease progression, and ultimately the risk of human infection [11,19].

A recent study by Pessôa-Pereira et al. [4] highlights how *Borrelia burgdorferi*, the agent of Lyme disease, exacerbates *Leishmania* infection in co-infected dogs. This study underscores the broader implications of co-infections involving vector-borne pathogens with overlapping hosts and/or vectors. *Babesia*, a protozoan parasite transmitted by ticks, causes babesiosis [20] and has been reported in co-infections with *Leishmania* in dogs [3,9,21]. Similarly, in humans, co-infection with *Babesia* and *Borrelia* worsens the clinical course of both Lyme disease and babesiosis [22]. Given these findings, it raises the possibility that co-infection with *Borrelia* might similarly affect dogs with *Leishmania*, potentially enhancing CanL progression. This hypothesis is supported by the study of Pessôa-Pereira et al. [4], which showed that *Borrelia* co-infection increases *Leishmania* survival in macrophages by altering immune responses. Combined immune modulation by *Borrelia*, *Leishmania*, and *Babesia*—each employing different mechanisms to evade or manipulate host defenses [3]—could amplify disease severity and complicate treatment.

Despite their structural differences and distinct pathogenesis mechanisms, bacteria and protozoa can establish ecological interactions within biological systems [23,24]. In terrestrial (e.g., soil) or aquatic ecosystems, bacterial communities can interfere with protozoan colonization or predation, often fostering antagonistic relationships [23–25]. These antagonisms arise through intracellular mechanisms (e.g., survival and replication within



Received: 25 December 2024

Accepted: 3 January 2025

Published: 6 January 2025

Citation: Piloto-Sardiñas, E.; Cano-Argüelles, A.L.; Cabezas-Cruz, A. *Borrelia*, *Leishmania*, and *Babesia*: An Emerging Triad of Vector-Borne Co-Infections? *Pathogens* **2025**, *14*, 36. <https://doi.org/10.3390/pathogens14010036>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

protozoan cells) or extracellular adaptations (e.g., altered cell morphology, increased motility, or biofilm formation) [26], as well as the production of bacteriocins with antiprotozoal activity [27]. However, these interactions differ significantly when subjected to the biology, physiology, and immune pressures of hosts and vectors.

During co-infections in hosts or vectors, pathogens may interact at various stages of their life cycles, targeting the same cells or eliciting overlapping immune responses. For instance, shared immune cell tropism between vector-borne bacteria and protozoa or the modulation of immune functions, such as macrophage activity, can allow bacterial infections to enhance protozoal infections [11]. This overlap enables *Borrelia*, *Leishmania*, and *Babesia* to exploit host immune systems, exacerbating disease severity. For instance, *Borrelia* promotes a skewed Th17 immune response that is inflammatory but insufficient for clearance [28], *Leishmania* inhibits macrophage reactive oxygen species (ROS) production [29], and *Babesia* suppresses adaptive immunity against *Borrelia*, worsening Lyme disease severity [30]. Together, these immune evasion strategies create a feedback loop of suppression, chronic infection, and increased zoonotic risk.

The proactive monitoring of co-infections in regions with vector overlap is vital. The surveillance of multi-pathogen exposure in dogs and wildlife can provide an early warning of zoonotic threats, aiding veterinarians and public health officials in mitigating risks. Dogs are sentinel hosts in zoonotic disease ecology [31], moving between environments shared with humans and wildlife. Co-infections in dogs highlight potential spillover risks to humans [19,32], especially in regions where expanding tick and sand fly populations overlap.

Managing co-infections involving *Leishmania*, *Borrelia*, and *Babesia* necessitates an integrated, multi-vector approach targeting both sand fly and tick populations. Effective measures, such as vector control, habitat management, and innovative immune-modulating therapies that address the compounded effects of co-infections, are crucial to mitigating disease severity in dogs and reducing zoonotic transmission risks. The study by Pessôa-Pereira et al. [4] highlights the critical implications of co-infections among vector-borne pathogens that share vectors and hosts, emphasizing how these interactions accelerate disease progression and complicate treatment. In an era of shifting ecological boundaries, understanding these pathogen interactions within hosts like dogs is vital for anticipating and mitigating zoonotic risks. By integrating vector control strategies, expanding pathogen surveillance in dogs and wildlife, and advancing targeted treatment approaches for co-infected hosts, we can better manage the complex dynamics of vector-borne diseases. This holistic approach is essential to protecting the health of both animal populations and the human communities with whom they share their environments.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Morales-Yuste, M.; Martín-Sánchez, J.; Corpas-Lopez, V. Canine leishmaniasis: Update on epidemiology, diagnosis, treatment, and prevention. *Vet. Sci.* **2022**, *9*, 387. [[CrossRef](#)] [[PubMed](#)]
2. Ribeiro, R.R.; Michalick, M.S.M.; da Silva, M.E.; Dos Santos, C.C.P.; Frézard, F.J.G.; da Silva, S.M. Canine leishmaniasis: An overview of the current status and strategies for control. *BioMed Res. Int.* **2018**, *1*, 3296893. [[CrossRef](#)] [[PubMed](#)]
3. Beasley, E.A.; Pessôa-Pereira, D.; Scorza, B.M.; Petersen, C.A. Epidemiologic, clinical and immunological consequences of co-infections during canine leishmaniosis. *Animals* **2021**, *11*, 3206. [[CrossRef](#)]
4. Pessôa-Pereira, D.; Scorza, B.M.; Cyndari, K.I.; Beasley, E.A.; Petersen, C.A. Modulation of Macrophage Redox and Apoptotic Processes to *Leishmania infantum* during Coinfection with the Tick-Borne Bacteria *Borrelia burgdorferi*. *Pathogens* **2023**, *12*, 1128. [[CrossRef](#)] [[PubMed](#)]
5. LaDeau, S.L.; Allan, B.F.; Leishman, P.T.; Levy, M.Z. The ecological foundations of transmission potential and vector-borne disease in urban landscapes. *Funct. Ecol.* **2015**, *29*, 889–901. [[CrossRef](#)]
6. Martina, B.E.; Barzon, L.; Pijlman, G.P.; de la Fuente, J.; Rizzoli, A.; Wammes, L.J.; Takken, W.; van Rij, R.P.; Papa, A. Human to human transmission of arthropod-borne pathogens. *Curr. Opin. Virol.* **2017**, *22*, 13–21. [[CrossRef](#)] [[PubMed](#)]

7. Sutherst, R.W. Global change and human vulnerability to vector-borne diseases. *Clin. Microbiol. Rev.* **2004**, *17*, 136–173. [[CrossRef](#)] [[PubMed](#)]
8. Ortiz, D.I.; Piche-Ovares, M.; Romero-Vega, L.M.; Wagman, J.; Troyo, A. The impact of deforestation, urbanization, and changing land use patterns on the ecology of mosquito and tick-borne diseases in Central America. *Insects* **2021**, *13*, 20. [[CrossRef](#)] [[PubMed](#)]
9. Cardoso, L.; Yisaschar-Mekuzas, Y.; Rodrigues, F.T.; Costa, Á.; Machado, J.; Diz-Lopes, D.; Baneth, G. Canine babesiosis in northern Portugal and molecular characterization of vector-borne co-infections. *Parasites Vectors* **2010**, *3*, 27. [[CrossRef](#)] [[PubMed](#)]
10. Ortúñoz, M.; Nachum-Biala, Y.; García-Bocanegra, I.; Resa, M.; Berriatua, E.; Baneth, G. An epidemiological study in wild carnivores from Spanish Mediterranean ecosystems reveals association between *Leishmania infantum*, *Babesia* spp. and *Hepatozoon* spp. infection and new hosts for *Hepatozoon martis*, *Hepatozoon canis* and *Sarcocystis* spp. *Transbound. Emerg. Dis.* **2022**, *69*, 2110–2125. [[CrossRef](#)] [[PubMed](#)]
11. Toepp, A.J.; Monteiro, G.R.; Coutinho, J.F.; Lima, A.L.; Larson, M.; Wilson, G.; Grinnage-Pulley, T.; Bennett, C.; Mahachi, K.; Anderson, B.; et al. Comorbid infections induce progression of visceral leishmaniasis. *Parasites Vectors* **2019**, *12*, 54. [[CrossRef](#)] [[PubMed](#)]
12. Little, S.; Braff, J.; Place, J.; Buch, J.; Dewage, B.G.; Knupp, A.; Beall, M. Canine Infection with *Dirofilaria immitis*, *Borrelia burgdorferi*, *Anaplasma* spp., and *Ehrlichia* spp. in the United States, 2013–2019. *Parasites Vectors* **2021**, *14*, 10. [[CrossRef](#)] [[PubMed](#)]
13. Attipa, C.; Solano-Gallego, L.; Leutenegger, C.M.; Papasouliotis, K.; Soutter, F.; Balzer, J.; Carver, S.; Buch, J.S.; Tasker, S. Associations between Clinical Canine Leishmaniosis and Multiple Vector-Borne Co-Infections: A Case-Control Serological Study. *BMC Vet. Res.* **2019**, *15*, 331. [[CrossRef](#)]
14. Miró, G.; Montoya, A.; Roura, X.; Gálvez, R.; Sainz, A. Seropositivity Rates for Agents of Canine Vector-Borne Diseases in Spain: A Multicentre Study. *Parasites Vectors* **2013**, *6*, 117. [[CrossRef](#)] [[PubMed](#)]
15. Evaristo, A.M.C.; Santos, P.T.T.; Sé, F.S.; Collere, F.C.M.; Silva, B.B.; Cardoso, E.R.; Kakimori, M.T.A.; Vieira, T.S.W.J.; Krawczak, F.S.; Moraes-Filho, J.; et al. Co-infection by tick-borne pathogens and *Leishmania* spp. in dogs with clinical signs suggestive of leishmaniasis from an endemic area in northeastern Brazil. *Pesqui. Veterinária Bras.* **2024**, *44*, e07437. [[CrossRef](#)]
16. De Sousa, K.C.M.; André, M.R.; Herrera, H.M.; de Andrade, G.B.; Jusi, M.M.G.; dos Santos, L.L.; Barreto, W.T.G.; Machado, R.Z.; de Oliveira, G.P. Molecular and Serological Detection of Tick-Borne Pathogens in Dogs from an Area Endemic for *Leishmania infantum* in Mato Grosso Do Sul, Brazil. *Rev. Bras. Parasitol. Vet.* **2013**, *22*, 525–531. [[CrossRef](#)] [[PubMed](#)]
17. Blondel, J. Guilds or functional groups: Does it matter? *Oikos* **2003**, *100*, 223–231. [[CrossRef](#)]
18. Lello, J.; Hussell, T. Functional group/guild modelling of inter-specific pathogen interactions: A potential tool for predicting the consequences of co-infection. *Parasitology* **2008**, *135*, 825–839. [[CrossRef](#)]
19. Baxarias, M.; Álvarez-Fernández, A.; Martínez-Orellana, P.; Montserrat-Sangrà, S.; Ordeix, L.; Rojas, A.; Nachum-Biala, Y.; Baneth, G.; Solano-Gallego, L. Does co-infection with vector-borne pathogens play a role in clinical canine leishmaniosis? *Parasites Vectors* **2018**, *11*, 135. [[CrossRef](#)] [[PubMed](#)]
20. Homer, M.J.; Aguilar-Delfin, I.; Telford III, S.R.; Krause, P.J.; Persing, D.H. Babesiosis. *Clin. Microbiol. Rev.* **2000**, *13*, 451–469. [[CrossRef](#)] [[PubMed](#)]
21. Krawczak, F.D.S.; Reis, I.A.; Silveira, J.A.D.; Avelar, D.M.; Marcelino, A.P.; Werneck, G.L.; Labruna, M.B.; Paz, G.F. *Leishmania*, *Babesia* and *Ehrlichia* in urban pet dogs: Co-infection or cross-reaction in serological methods? *Rev. Soc. Bras. Med. Trop.* **2015**, *48*, 64–68. [[CrossRef](#)] [[PubMed](#)]
22. Swanson, S.J.; Neitzel, D.; Reed, K.D.; Belongia, E.A. Coinfections acquired from Ixodes ticks. *Clin. Microbiol. Rev.* **2006**, *19*, 708–727. [[CrossRef](#)]
23. Snelling, W.J.; Moore, J.E.; McKenna, J.P.; Lecky, D.M.; Dooley, J.S. Bacterial–protozoa interactions; an update on the role these phenomena play towards human illness. *Microbes Infect.* **2006**, *8*, 578–587. [[CrossRef](#)]
24. Eriksson, K.I.; Thelaus, J.; Andersson, A.; Ahlinder, J. Microbial interactions—Underexplored links between public health relevant bacteria and protozoa in coastal environments. *Front. Microbiol.* **2022**, *13*, 877483. [[CrossRef](#)]
25. Clarholm, M. Protozoan grazing of bacteria in soil—Impact and importance. *Microb. Ecol.* **1981**, *7*, 343–350. [[CrossRef](#)] [[PubMed](#)]
26. Song, C.; Mazzola, M.; Cheng, X.; Oetjen, J.; Alexandrov, T.; Dorresteijn, P.; Watrous, J.; van der Voort, M.; Raaijmakers, J.M. Molecular and chemical dialogues in bacteria–protozoa interactions. *Sci. Rep.* **2015**, *5*, 12837. [[CrossRef](#)]
27. Gupta, R.; Rajendran, V.; Ghosh, P.C.; Srivastava, S. Assessment of anti-plasmodial activity of non-hemolytic, non-immunogenic, non-toxic antimicrobial peptides (AMPs LR14) produced by *Lactobacillus plantarum* LR/14. *Drugs R&D* **2014**, *14*, 95–103.
28. Strle, K.; Sulka, K.B.; Pianta, A.; Crowley, J.T.; Arvikar, S.L.; Anselmo, A.; Sadreyev, R.; Steere, A.C. T-Helper 17 Cell Cytokine Responses in Lyme Disease Correlate with *Borrelia burgdorferi* Antibodies during Early Infection and with Autoantibodies Late in the Illness in Patients with Antibiotic-Refractory Lyme Arthritis. *Clin. Infect. Dis.* **2017**, *64*, 930–938. [[CrossRef](#)]
29. Saha, S.; Basu, M.; Guin, S.; Gupta, P.; Mitterstiller, A.-M.; Weiss, G.; Jana, K.; Ukil, A. *Leishmania donovani* Exploits Macrophage Heme Oxygenase-1 To Neutralize Oxidative Burst and TLR Signaling-Dependent Host Defense. *J. Immunol.* **2019**, *202*, 827–840. [[CrossRef](#)]

30. Djokic, V.; Akoolo, L.; Primus, S.; Schlachter, S.; Kelly, K.; Bhanot, P.; Parveen, N. Protozoan parasite *Babesia microti* subverts adaptive immunity and enhances Lyme disease severity. *Front. Microbiol.* **2019**, *10*, 1596. [[CrossRef](#)] [[PubMed](#)]
31. Bowser, N.H.; Anderson, N.E. Dogs (*Canis familiaris*) as sentinels for human infectious disease and application to Canadian populations: A systematic review. *Vet. Sci.* **2018**, *5*, 83. [[CrossRef](#)] [[PubMed](#)]
32. Cevidanes, A.; Di Cataldo, S.; Muñoz-San Martín, C.; Latrofa, M.S.; Hernández, C.; Cattan, P.E.; Otranto, D.; Millán, J. Co-infection patterns of vector-borne zoonotic pathogens in owned free-ranging dogs in central Chile. *Vet. Res. Commun.* **2023**, *47*, 575–585. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.