



HAL
open science

Toward Decision Support System for Lower Limb Endovascular Revascularization

M Roux, R. Spear, Pascal Haigron, Alexandre Demeure, Céline Fouard

► **To cite this version:**

M Roux, R. Spear, Pascal Haigron, Alexandre Demeure, Céline Fouard. Toward Decision Support System for Lower Limb Endovascular Revascularization. 2024 46th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Jul 2024, Orlando, United States. pp.1-4, 10.1109/embc53108.2024.10782314 . hal-04874183

HAL Id: hal-04874183

<https://hal.science/hal-04874183v1>

Submitted on 11 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Toward Decision Support System for Lower Limb Endovascular Revascularization

Roux M.¹, Spear R.², Haigron P.³, Demeure A.⁴ and Fouard C.¹

Abstract—Vascular surgeons take decisions on the revascularization technics when surgery is needed for patients with Lower Extremities Arterial Disease (LEAD). This decision requires experience, good knowledge of official recommendations and new technics and an exhaustive description of the patient status. The Case Based Reasoning (CBR) is a good candidate to clinical decision support. To assess its applicability to LEAD, this study proposes a numerical modelization of patient data in this context, application of hierarchical and heterogenous similarity measures on retrospective data. We showed that similar pathological contexts of patients induce similar revascularization technics, which confirms CBR is usable for our purpose. These experiments need to be reproduced in the future to confirm these results on prospective data.

Index Terms—Decision Support System, Case Based Reasoning, Modelization, Vascular Surgery

I. INTRODUCTION

Atherosclerosis is an arterial pathology that describes the accumulation of plaques on arterial walls. At some point, it can lead to obstruction of arteries: this phenomenon is called ischemia. This work focuses on arteries of lower limbs where the pathology is the Lower Extremities Arterial Disease (LEAD). Four different stages were identified by Leriche and Fontaine [1] depending of the severity of the symptoms: stage I is the asymptomatic stage, stage II when pain is felt while walking, stage III when pain is felt at rest and stage IV for the combination of pain at rest and trophic disorders (i.e. ischemic wounds).

To avoid amputation, vascular surgeons can revascularize the lower limb with different technics: endovascular surgery and open surgery are the two main types of revascularization that exist with different level of invasivity. Endovascular revascularization is minimally invasive surgery that consists in treating the obstruction by introducing the tools directly through the arterial lumen. Different technics exist, angioplasty and stenting for instance that can be combined. Open surgery, is more invasive as the technics usually involve opening the thigh to access the artery.

The revascularization has three goals: the wound healing, the relief of pain, and the maintenance of the ambulatory status. As patients concerned by LEAD are usually patients

over 60 years old with numerous comorbidities, the recovery and thus the invasivity are important parameters to consider.

The choice of a technic or another relies on Patient Data, Official recommendations, surgeon experience and the experience of their peers. However, these decision factors are not always available: patient data in the hospital informatic system is scattered in several databases, redundant and unstructured ; official recommendations are not always up to date with newest technics and leave space for interpretations; the experience of other surgeons is not always fully available nor explicit.

To help overcoming these limits, Decision Support tools have already proved their efficacy in other medical domains, [2]. There are two main types of algorithms : the training based algorithms and knowledge based algorithms [3]. Among training based ones, we can count neural networks, bayesian networks as example, that have already been used for clinical decision support [4]. However, when using training based methods, knowledge is acquired through training, which requires large and exhaustive databases. In addition, as the whole pathological context of patient needs to be considered, the LEAD database uses much more descriptive variables compared to the number of observations. Therefore, statistical significance cannot be reached. In this study, we focus on knowledge based methods where the knowledge is given to the system.

Among knowledge-based methods, the Case Based Reasoning (CBR) uses experience of past surgeries as the source of knowledge. It has already been used in other clinical contexts [5]–[7].

It is called 'Case Based' because it uses a certain amount of already solved cases, here past surgeries. It works in a four steps cycle: when a new case to solve enters the system, it will first retrieve the most similar case(s) among the case base thanks to similarity measures. The solution(s) of these similar cases give the suggested solution(s). The revise step aims to adapt the suggested solution(s) to the given problem, thanks to adaptation rules, to obtain the confirmed solution(s). Once a solution is confirmed, it is tested and the result obtained, the retain step will integrate this new case into the case base to enrich the system.

This algorithm mimics the human reasoning as it solves a new problem by remembering a previous similar situation and by reusing information and knowledge of that situation [8]. This functioning and its explainability make it an interesting method for our purpose.

The CBR works by retrieving the most similar cases.

This work has been funded by the French Agence Nationale de la Recherche, CAMI Labex (Grant Number: ANR-11-LABX-0004), (Grant Number MIAI@Grenoble Alpes ANR-19-P3IA-0003)

1. Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, TIMC, 38000 Grenoble, France
2. Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, CHU Grenoble Alpes, TIMC, 38000 Grenoble, France
3. Univ. Rennes, Inserm, LTSI – UMR 1099, Rennes, France
4. Laboratoire Informatique de Grenoble, Université Grenoble Alpes, France

Thus, we will need to measure similarity which is a quantity reflecting of how much two elements are alike. Numerous similarity measures already exist [9]. It can be defined as a function that quantifies either the distance i.e. the dissimilarity or the correlation i.e. the strength of the link between two elements. Helène Feuillâtre et. al. [5] defined a hierarchical and heterogeneous distance between patients in the context of aortic valve replacement.

This study aims to assess the applicability of CBR to the context of revascularization of lower limbs. After detailing the data model needed, we illustrate how it is used to measure similarity between cases.

II. METHODS

To evaluate the pertinence of CBR, we focus on the following clinical question : " For a given patient, what would be the most efficient revascularization technic to relief pain, heal wounds and maintain patient mobility ?". To answer this question thanks to CBR methods, we need to design the model of a numerical patient using raw clinical data. Thus, we first need to define how to describe cases, i.e. the attributes necessary to describe patient pathological context. Secondly, we need to define what are similar cases.

A. Definition of a Case

A case is a triplet of the following elements: the attributes describing the problem, here the list of variables that will describe the symptoms and the whole pathological context of a patient; the solution that was chosen and applied, the revascularization method the most efficient for a given patient i.e. either an endovascular technics or open surgery technics; and the result of the solution (success or fail), here the variables telling weather the revascularization goals (mentioned earlier) are fulfilled or not. In our case, the success can be defined by the technical success, which is the immediate restoration of blood flow within surgery, and the clinical success i.e. relieved pain, healing wounds and retrieved mobility. Whereas the failure is the combination of either a reintervention or amputation on the same side, perioperative death or technical failure during surgery.

B. Data Structure

After defining a case in the context of vascular surgery, we identified the variables necessary for our numerical model.

1) *Hierarchy of data model*: Selection of variables was done using the official recommendations and clinical studies that proposed clinical decision trees, gradation scales and outcome assessment regarding our pathology of interest, and with the help of an expert surgeon.

Concerning the preoperative data, the Society of Vascular Surgery proposed a therapeutic strategy for patient with LEAD : the PLAN concept [10]. It states that for a patient, three aspects have to be considered in the following order: *Patient risk estimation*, that is to say comorbidities (chronic kidney failure, bronchopneumopathies, cardiac pathologies) and risk factors [11], [12]; *Limb staging* that uses the WIfI score [13],

[14] to estimate the amputation risk of the patient situation and the benefits of revascularization; *ANatomical pattern* based on interpretation and scoring of CT scans.

Regarding the solution attributes i.e. the variables describing the revascularization technics used, we identified three levels of attributes: the type of surgery which is either endovascular or open surgery ; the second level of attributes concerns the technics or combination of technics used (angioplasty, stenting, bypass, ...); the third level of attributes are the details on the technics, such as the size of stents, diameters, coatings of ballons, and material of prosthesis for example.

The attributes of the results of the surgery are about the achievements of the revascularization goals described in the clinical context. To assess the outcome as a success or a fail, the evolution of pain, evolution of wounds and ambulatory status is needed at different delays These informations are collected during postoperative consultations.

As a result of this selection, more than 650 variables constitute our patient model in total.

2) *Type of data model*: The type of data is also important to consider as it directly concerns the consistency and exhaustivity of our model, which is crucial for the similarity measurements in CBR. Clinical data are heterogeneous as they include quantitative, binary, qualitative and ordinal data and images.

To avoid qualitative data, they were converted to lists of binary variables. For example, rather than having a variable where the possible values are the different cardiopathies, each possible values became a binary variables.

Ordinal data, are between qualitative and quantitative: they are categories representing a scale. They are often used in medical domain for diagnosis: as an example ranges of hemoglobin concentration values to give the grade of anemia. However, in order to compare patients as precisely as possible, we chose to keep the raw quantitative measurements instead of grades when it was possible.

Regarding images, CT scans are the examination helping surgeons to determine how tortuous is the arterial tree of a given patient. We chose to use variables describing the interpretations from surgeons of these images: precise localization of lesions, presence of calcifications, length, degree of stenosis among others.

3) *Missing data*: Missing data is an important characteristic of clinical data. Indeed, depending on the context, the patients arrive with different amount of information. Moreover during patient interrogation some data is considered as implicit. For example, when nothing is mentioned concerning the smoking status on a patient file, we can either assume that the patient is not a smoker, or that we do not know. To avoid that confusion, we defined a ternary format of variables: value could be True, False, or Unknown.

C. Data Collection

To build a database following the data structure described above, we develop a software to collect data. The first objective is to design this tool in a user-centered way, the first users




Fig. 1. View of a patient file on the data collection software

being the surgeons. The whole organization of the forms is based on vascular surgeons workflow. The second objective is the collection of data in a exhaustive and structured way allowing access to anonymized these clinical data. This is achieved by integrating the patient data model in the forms.

D. Similarity Measures

To retrieve contexts which are the most alike, we need to measure similarity between surgeries with distances. Among numerous distances, we decided to try the Hierarchical Weighted and Heterogenous Similarity Measure (H-WHSM) [5] as it integrates the different format of data encountered in the clinical context. Hierarchy of variables is integrated with the ponderation deduced by clinical decision trees and in our case from the PLAN concept described earlier. Ponderation was applied to preoperative variables only.

III. RESULTS

To verify the applicability of CBR to LEAD, we developed a software to collect structured, normalized and exhaustive data and performed preliminary studies on retrospective data.

A. Data collection software

We developed prospective data collection system integrated in a web application in figure 1.

This tool is currently deployed in the Centre Hospitalier Universitaire Grenoble Alpes.

B. Retrospective data

The retrospective database used for our preliminary experiments was collected for technical and legal studies between 2018 and 2020 with patients going under lower limb revascularization. As They were not intended for CBR, simplifications hypothesis of the data model had to be made.

Concerning missing data, the ternary format can only be used for prospective data collection. As quantitative measurements as were not all available, we used ordinal variables (scales and grades). Regarding images, interpretations of surgeons were not available. Thus, we used interpretations of reports made by radiologists instead.

Another element impacting our data collection was the chronicity of LEAD: even after revascularization obstruction




Fig. 2. Example1: 3D plot of the distance between patients and their solution and results

can happen again, at the same location or a different one. Hence, we considered only one surgery per patient.

With these simplifications, from 675 variables in the proposed model, we kept 75 variables: 45 are preoperative attributes, 9 are surgical solution attributes, and 17 are results attributes. These retrospective data on surgeries were used to perform preliminary studies on similarity measures.

The solution, surgical technic, could be described by multiple variables. To simplify the result display and interpretation, we defined three categories for surgery types: pen surgery, angioplasty or angioplasty and stenting.

The result of a revascularization procedure is a combination of multiple factors. We chose to define the result as binary: success or failure. The success is the non-occurrence of re-intervention, amputation or death within 3 months after surgery. The failure is the occurrence of at least one of these events.

The case base constituted includes 38 cases to perform preliminary studies.

C. Similarity measure on retrospective dataset

We used H-WHSM on preoperative attributes as similarity measure to retrieve the most similar cases to a test case. The test case is a known case not included in the case base. Among the closests, we represented the surgery type that was used for the patient and the result (success or failure.)

The figure 2 represents for a given patient the distance regarding the preoperative variables on x axis, the type of surgery applied on the y axis, and the clinical result of the surgery on the z axis. The triangle represents the patients of reference i.e. the patient for which we retrieve the most similar ones. In orange, we represent the 5 most similar patients. The test case went under endovascular surgery for Angioplasty with stenting and it was a success. The 5 most similar patients went under endovascular surgery with success as a result and among these 5 patients, 4 received an angioplasty with stenting exactly as the patient of reference. Hence, the pathological




Fig. 3. Example2: 3D plot of the distance between patients and their solution and results

context described by the attributes seems to indicate an endovascular technic.

On the second example (figure 3), the test case went under open surgery for revascularization and it was a success. The most similar patients did not go under open surgery but 3 of the 5 most similar patients went under endovascular surgery (angioplasty and/or stenting) with clinical failure. Nevertheless, 2 of the 5 most similar patients went under endovascular surgeries with a successful outcome.

IV. DISCUSSION

These preliminary results show that CBR could be applied to the context of endovascular revascularization. However, we observe disparities, in solutions and results, for some cases.

As retrospective dataset was not designed for a CBR application, simplification hypothesis were made compared to the proposed model. The exhaustivity planned in the proposed model could bring more accuracy on the patient model with a prospective database. This is why, such experiment should be reproduced with a prospective and exhaustive dataset.

The case base used here might not be representative of the whole spectrum of pathological context.

Definition of success and failure of a surgery were made binary in our preliminary results. However, in a clinical context and especially concerning chronic diseases, success or failure of a solution is not binary. We also need to consider other variables impacting the results such as the evolution of wounds and of the ambulatory status. These postoperative variables could be combined to use a success score of a revascularization intervention.

We chose the H-WHSM distance for our preliminary study, however other similarity measures should be tested to identify the one that is the most efficient on retrieving the most similar cases in our field of application.

REFERENCES

- [1] R. Fontaine, M. Kim, and R. Kieny, "Surgical treatment of peripheral circulation disorders," *Helv. Chir. Acta*, vol. 21, no. 5-6, pp. 499–533, Dec. 1954.
- [2] D. Dinevski, U. Bele, T. Sarenac, U. Rajkovic, and O. Sustersic, *Clinical Decision Support Systems*. InTech, June 2011. [Online]. Available: <http://dx.doi.org/10.5772/25399>
- [3] R. T. Sutton, D. Pincock, D. C. Baumgart, D. C. Sadowski, R. N. Fedorak, and K. I. Kroeker, "An overview of clinical decision support systems: benefits, risks, and strategies for success," *npj Digital Medicine*, vol. 3, no. 1, Feb. 2020. [Online]. Available: <http://dx.doi.org/10.1038/s41746-020-0221-y>
- [4] Z. B. Perkins, B. Yet, A. Sharrock, R. Rickard, W. Marsh, T. E. Rasmussen, and N. R. M. Tai, "Predicting the outcome of limb revascularization in patients with lower-extremity arterial trauma: Development and external validation of a supervised machine-learning algorithm to support surgical decisions," *Annals of Surgery*, vol. 272, no. 4, p. 564–572, July 2020. [Online]. Available: <http://dx.doi.org/10.1097/SLA.0000000000004132>
- [5] H. Feuillâtre, V. Auffret, M. Castro, F. Lalys, H. L. Breton, M. Garreau, and P. Haigron, "Similarity measures and attribute selection for case-based reasoning in transcatheter aortic valve implantation," *PLOS ONE*, vol. 15, no. 9, p. e0238463, Sept. 2020. [Online]. Available: <https://doi.org/10.1371/journal.pone.0238463>
- [6] J. Buyer, A. Oeser, N. Grieb, A. Dietz, T. Neumuth, and M. Stoehr, "Decision support for oropharyngeal cancer patients based on data-driven similarity metrics for medical case comparison," *Diagnostics*, vol. 12, no. 4, p. 999, Apr. 2022. [Online]. Available: <https://doi.org/10.3390/diagnostics12040999>
- [7] S. Guessoum, M. T. Laskri, and J. Lieber, "Respiadiag: A case-based reasoning system for the diagnosis of chronic obstructive pulmonary disease," *Expert Systems with Applications*, vol. 41, no. 2, pp. 267–273, 2014. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S0957417413003631>
- [8] A. Aamodt and E. Plaza, "Case-based reasoning: Foundational issues, methodological variations, and system approaches," *AI Communications*, vol. 7, no. 1, p. 39–59, 1994. [Online]. Available: <http://dx.doi.org/10.3233/AIC-1994-7104>
- [9] R. M. M., W. Rosina, and M. M. Richter, *Case-Based Reasoning : A Textbook / Michael M. Richter / Rosina O. Weber*. Berlin, Berlin, Heidelberg: Springer-Verlag Berlin Heidelberg, 2013.
- [10] M. S. Conte, A. W. Bradbury, P. Kolh, J. V. White, F. Dick, R. Fitridge, J. L. Mills, J.-B. Ricco, K. R. Suresh, M. H. Murad, and GVG Writing Group, "Global vascular guidelines on the management of chronic limb-threatening ischemia," *J Vasc Surg*, vol. 69, no. 6S, pp. 3S–125S.e40, May 2019.
- [11] J. P. Simons, A. Schanzer, J. M. Flahive, N. H. Osborne, J. L. Mills Sr., A. W. Bradbury, and M. S. Conte, "Survival prediction in patients with chronic limb-threatening ischemia who undergo infrainguinal revascularization," *Journal of Vascular Surgery*, vol. 69, no. 6, pp. 137S–151S.e3, Jun 2019. [Online]. Available: <https://doi.org/10.1016/j.jvs.2018.08.169>
- [12] M. C. Verwer, J. G. Wijnand, M. Teraa, H. Gremmels, J. P. Simons, M. S. Conte, M. C. Verhaar, and G. J. de Borst, "External validation of the vascular quality initiative prediction model for survival in no-option chronic limb-threatening ischemia patients," *Journal of Vascular Surgery*, vol. 72, no. 5, pp. 1659–1666.e1, Nov 2020. [Online]. Available: <https://doi.org/10.1016/j.jvs.2020.02.018>
- [13] L. X. Zhan, B. C. Branco, D. G. Armstrong, and J. L. Mills Sr., "The society for vascular surgery lower extremity threatened limb classification system based on wound, ischemia, and foot infection (wif) correlates with risk of major amputation and time to wound healing," *Journal of Vascular Surgery*, vol. 61, no. 4, pp. 939–944, Apr 2015. [Online]. Available: <https://doi.org/10.1016/j.jvs.2014.11.045>
- [14] D. L. Cull, G. Manos, M. C. Hartley, S. M. Taylor, E. M. Langan, J. F. Eidt, and B. L. Johnson, "An early validation of the society for vascular surgery lower extremity threatened limb classification system," *Journal of Vascular Surgery*, vol. 60, no. 6, pp. 1535–1542, Dec 2014. [Online]. Available: <https://doi.org/10.1016/j.jvs.2014.08.107>