



HAL
open science

**Starvation of the bacteria *Vibrio atlanticus* promotes
lightning group-attacks on the dinoflagellate
*Alexandrium pacificum***

Jean Luc Rolland, Estelle Masseret, Mohamed Laabir, Guillaume Tetreau,
Benjamin Gourbal, Anne Thebault, Eric Abadie, Alice Rodrigues-Stien,
Carole Veckerlé, Elodie Servanne-Meunier, et al.

► **To cite this version:**

Jean Luc Rolland, Estelle Masseret, Mohamed Laabir, Guillaume Tetreau, Benjamin Gourbal, et al..
Starvation of the bacteria *Vibrio atlanticus* promotes lightning group-attacks on the dinoflagellate
Alexandrium pacificum. 2024. hal-04879695

HAL Id: hal-04879695

<https://hal.science/hal-04879695v1>

Preprint submitted on 10 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Starvation of the bacteria *Vibrio atlanticus* promotes lightning group-attacks on the
2 dinoflagellate *Alexandrium pacificum*

3

4 Jean Luc Rolland^{1,2†*}, Estelle Masseret³, Mohamed Laabir³, Guillaume Tetreau⁴, Benjamin Gourbal⁴,
5 Anne Thebault⁵, Eric Abadie⁶, Alice Rodrigues-Stien^{7,8}, Carole Veckerlé^{7,8}, Elodie Servanne-Meunier¹,
6 Delphine Destoumieux-Garzón¹, Arnaud Lagorce^{4,9}, Raphaël Lami^{7,9}.

7

8 **Affiliations**

9 ¹IHPE, Univ. Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Montpellier, France.

10 ²MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Sète, France.

11 ³MARBEC, Univ. Montpellier, IRD, IFREMER, CNRS, Montpellier, France.

12 ⁴IHPE, Univ Montpellier, CNRS, IFREMER, Université de Perpignan Via Domitia, Perpignan, France

13 ⁵Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES),
14 ANSES-DER 27-31, avenue du Général Leclerc, 94701 Maison-Alfort, France

15 ⁶IFREMER, 9 rue Jean Bertho, 97822, Le port, la Reunion, France

16 ⁷Sorbonne Université, CNRS, LBBM USR3579, Observatoire Océanologique de Banyuls sur Mer,
17 Avenue Pierre Fabre, 66650 Banyuls-sur-Mer, France.

18 ⁸Sorbonne Université, CNRS, Bio2MAR, Observatoire Océanologique de Banyuls sur Mer, Avenue Pierre
19 Fabre, 66650 Banyuls-sur-Mer, France.

20

21 ⁹These authors contributed equally

22 †Current address

23 *Corresponding author: Jean.Luc.Rolland

24 **Email:** Jean.Luc.Rolland@ifremer.fr

25 **ABSTRACT**

26 Algae serve as a source of nutrients for bacteria in the marine environment. The interactions between
27 algae and bacteria at the phycosphere interface are known to include mutualism, commensalism,
28 competition or antagonism. Here, based on *in situ* observation and on an *in vitro* interaction study, we
29 report on a novel form of starvation-induced hunting that the cells of selected *Vibrio* species exert on the
30 dinoflagellate species. Results showed that *Vibrio atlanticus* was able to coordinate lightning group
31 attacks then kill the dinoflagellate *Alexandrium pacificum* ACT03. Briefly, the observed coordinated
32 mechanism of algal-killing consists of first, the ‘immobilization stage’ involving the secretion of algicidal
33 metabolites that disrupt the flagella of the prey. The ‘attack stage’ resembles the ‘wolf-pack attack’
34 strategy, during which *Vibrios* surrounds algal cells at high density for a brief period without invading
35 them. Finally, the ‘killing stage’ induces the lysis and degradation of the dinoflagellates.
36 By using a combination of biochemical, proteomic, genetic and fluorescence microscopy approaches, we
37 showed that this relationship is not related to the decomposition of algal organic matter, *Vibrio* quorum
38 sensing pathways, to the toxicity of the algae or to the pathogenicity of the bacterium but is conditioned by
39 nutrient stress, iron availability and link to the iron-vibrioferrin transport system of *V. atlanticus*. This is
40 the first evidence of a new mechanism that could be involved in regulating *Alexandrium* spp.
41 blooms and giving *Vibrio* a competitive advantage in obtaining nutrients from the environment.

42

43 **KEYWORDS**

44 Interaction; Bacterium; Alga; Predation; Environment; Iron related predation

45

46 INTRODUCTION

47 Harmful algal blooms (HABs) have experienced an increase in their occurrence, intensity, and
48 geographical distribution on a global scale, resulting in adverse environmental, health, and socioeconomic
49 impacts (Marampouti et al., 2021). HABs have a considerable impact on human health as a result of direct
50 exposure to volatile toxins or by toxic seafood consumption (Burkholder et al., 2018). From an ecological
51 point of view, the expansion of HABs can result in the erosion of biodiversity, because they cause massive
52 mortality of marine species and they are generally monospecific in nature (Chai et al., 2020). In coastal
53 areas, understanding the biological interactions that control toxic algal blooms is therefore a major
54 ecological challenge. Among HABs, a number of *Alexandrium* species have been placed on the list of
55 invasive Mediterranean species. Among them, *Alexandrium pacificum* is a flagellated eukaryotic
56 unicellular organism that form with *Alexandrium tamarense* and *Alexandrium fundyense* the
57 “*Alexandrium tamarense*” complex (Hadjadji et al., 2020). Since 1998, *A. pacificum* (former *A. catenella*)
58 was monitored by RePHY survey in the Thau lagoon (French Mediterranean) because it produces paralytic
59 shellfish toxins (PSTs) resulting of paralytic shellfish poisoning (PSP) syndrome.
60 Laania (Laania et al., 2013) showed that in Thau lagoon, a water temperature around 20°C for several
61 days and organic and inorganic nutrients in sufficient concentrations are parameters favoring the
62 developemnt of *A. pacificum*, whose massive blooms occur in autumn.
63 Algal blooms are seasonal events resulting in a rapid increase in the concentration of a species of algae in
64 an aquatic environment. Depending on the phytoplankton species, the tolerance range of physicochemical
65 parameters is different and influences the time of appearance of blooms (Leblad et al., 2020).
66 Interestingly, although the collapse of phytoplankton blooms has been previously attributed to viruses (Pal
67 et al., 2020), some ecological studies have suggested an important role of algicidal bacteria (Su et al.,
68 2007; Wang et al., 2010), such as some *Vibrio spp.* (Li et al., 2014; Wang et al., 2020).
69 *Vibrio* (class γ -proteobacteria) are common microorganisms in marine systems worldwide (Baker-Austin
70 et al., 2017; Mavian et al., 2020), where they are important components of the food chain, particularly in

71 biodegradation, nutrient regeneration and biogeochemical cycles (Oberbeckmann et al., 2012). *Vibrio* is
72 one of the most studied bacterial taxa due to their ubiquity in coastal marine systems and their capacity to
73 cause infections in humans and animals, leading sometimes to epizootic or zoonotic epidemics (LeRoux et
74 al., 2015; Mavian et al., 2020). *Vibrio* are extremely adaptable to their environment (Johnson, 2013). The
75 main factors influencing their occurrence and distribution in water are temperature, salinity, nutrient
76 availability (Wang et al., 2020), multiple strategies such as biofilm formation on biotic and abiotic
77 surfaces (Espinoza-Vergara et al., 2020), or interactions with a multitude of other organisms such as
78 eukaryotic predators (Drebes Dörr and Blokesch, 2020) or plankton (Lopez-Joven et al., 2018) are used by
79 *Vibrio* in the environment. There is also evidence that global climate change has increased *Vibrio*-
80 associated illnesses affecting humans and animals (Brumfield et al., 2021; Muhling et al., 2017).
81 However, the drivers and dynamics of *Vibrio* survival and propagation in the marine environment are not
82 yet fully understood.

83 A substantial number of research articles have highlighted the potential of γ -proteobacteria to exert
84 algicidal activity against dinoflagellates, supporting the hypothesis that γ -proteobacteria such as *Vibrio*
85 play a role in the control of algae blooms *in situ* (Coyne et al., 2022). However, the mechanisms behind
86 *Vibrio*-driven algal lysis in the environment remain to be elucidated. Particularly, it is unclear how in the
87 water column, algicidal compounds secreted by bacteria can concentrate around the algae to exert their
88 lytic effect.

89 Based on observations from the natural environment showing a potent relationship between *Vibrio* and
90 *Alexandrium* algae bloom events, this study aim to determine *in vitro*, the main factors implicated in this
91 relationship. Using a combination of biochemical, proteomic, genetic, and fluorescence
92 microscopy approaches, we explored the role of algal toxicity, bacterial pathogenicity and the
93 quorum sensing pathway on this relationship and showed the important role of nutrient stress and
94 the iron uptake pathway in this unique *Vibrio*/*Alexandrium* predator/prey interaction.

95

96 **METHODS**

97 **Quantification of *Alexandrium* algae and *Vibrio* bacteria in the environment.**

98 Seawater samples were collected in the Thau Lagoon (southern France, a shallow Mediterranean
99 ecosystem open to the sea (Abadie et al., n.d.), during spring and autumn 2015. Briefly, samples were
100 collected from the subsurface (-50 cm) near an oyster table at a phytoplankton surveillance site (part of the
101 REPHY network, N 43°26.058' and E 003°39.878'). Once a week during spring and autumn 2015, during
102 field sampling campaigns, 20 L of water was filtered on board through a 180 µm pore-size nylon
103 membrane. At the laboratory, according to Lopez-Joven et al. (Lopez-Joven et al., 2018) seawater was
104 fractionated into two size classes as follows: 2 L of the above filtrate was filtered through a 0.8 µm pore-
105 size polycarbonate Whatman Nuclepore membrane to obtain organisms in the 0.8–180 µm range
106 corresponding to plankton-associated *Vibrio* and living *Alexandrium* forms. Then, the filtrate from the 0.8
107 µm membrane was filtered again through a 0.2 µm pore-size polycarbonate Whatman Nuclepore
108 membrane until the membrane was saturated. *Alexandrium* cells, ranging from 25 to 40 µm, belong to the
109 microphytoplankton and are therefore retained in the 0.8–180 µm fraction. Any *Vibrio* cells potentially
110 associated with or attached to *Alexandrium* cells will also be retained in this fraction. *Vibrio* cells are
111 approximately 0.5-0.8 µm-thick. The fraction between 0.2 and 0.8 µm therefore includes the free-living
112 *Vibrio*. The bacterial population collected on 0.8-µm-pore-size filters was designated the particle-
113 associated community, and the population on 0.2-µm-pore-size filters was designated as the free-living
114 community. Membranes (in triplicate) were then conserved in 500 µL of 100% EtOH at -20°C.
115 Environmental DNA (eDNA) was extracted from the MF Millipore membrane using the Macherey-Nagel
116 NucleoSpin Tissue Kit and resuspended in 100 µL of water. The samples were then stored at -20°C after
117 eDNA quantity and purity were assessed using a NanoDrop system (NanoDrop Technologies,
118 Wilmington, DE, USA). PCR amplification reactions were done on a Roche LightCycler 480 Real-Time

119 thermocycler (qPHD platform, University of Montpellier, France).using specific primer pairs (Table 3).
120 Typically, the reactions contained 1 μL of template DNA (the DNA concentration for all samples varied
121 from 1 to 40 $\mu\text{g mL}^{-1}$), 0.5 μL of each primer (3.33 μM) and 4 μL of reaction mixture (SYBR Green
122 Master Mix) in a total volume of 6 μL . The reaction parameters were as follows: 5 minutes at 95°C (initial
123 denaturation) and 40 cycles of 10 s at 95°C (denaturation), 10 s at the corresponding hybridization
124 temperature (Table 3) and 10 s at 72°C (elongation). Melting curve profiles were generated by increasing
125 the temperature from 65°C to 95°C at 0.5°C s^{-1} . Amplification products were analysed using LightCycler
126 software (Roche Diagnostics). *Vibrio spp.* and *A. pacificum* and *A. tamarensis* were quantified by
127 constructing calibration curves based on DNA from the *V. atlanticus* LGP32 reference strain (former *V.*
128 *tasmaniensis* LGP32) and from the *A. pacificum* reference strain (ACT03: *A. catenella* strain isolated from
129 Thau in 2003) and the *A. tamarensis* reference strain (ATT07: *A. tamarensis* isolated from Thau in 2007)
130 (not shown).

131

132 **Strains and growth conditions.**

133 *Vibrio* strains. Wild-type and isogenic mutants of *Vibrio atlanticus* LGP32 (Table 1) were used in this
134 study. Deletion-mutants included $\Delta luxR$, $\Delta luxP$, $\Delta luxM$ and $\Delta pvuB$ isogenic strains. The $\Delta pvuB$ mutant
135 was constructed here by allelic exchange as described previously by Le Roux (Le Roux et al., 2007). We
136 also used *V. atlanticus* LGP32 carrying the pSW3654T-GFP plasmid (Le Roux et al., 2007), hereafter
137 referred to as *V. atlanticus* LGP32-GFP. Bacterial strains were grown at $22 \pm 1^\circ\text{C}$ in Zobell medium (0, 38
138 μM iron (III)). When needed, 25 $\mu\text{g mL}^{-1}$ chloramphenicol (Cm) was added to cultures of *V. atlanticus*
139 LGP32-GFP (Le Roux et al., 2007). Oligonucleotides used for RNA expression analysis are specified in
140 table 3.

141 *Phytoplankton strains.* Non-axenic phytoplankton species (Table 2) were grown in batch culture in
142 enriched natural sea water (ENSW, 6,55 μM iron (III)) with a salinity of 36 practical salinity units (PSU)
143 at $22 \pm 1^\circ\text{C}$ under cool white fluorescent illumination (100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and a 12 h:12 h

144 light:dark cycle (Harrison et al., 1980). The algae were used for experiments in their exponential growth
145 phase.

146 **Co-culture assay**

147 For each tested phytoplankton species, $2 \cdot 10^4$ cells harvested in their exponential growth phase were placed
148 in 20 mL of ENSW medium in a 50 mL suspension culture flask (Cellstar® PS, Greiner bio-one). After
149 incubation for 24 h at $22 \pm 1^\circ\text{C}$ under cool white fluorescent illumination, 40 μL of *Vibrio* strains grown
150 for 12, 36, 60 and 156 h in Zobell medium or in Zobell medium supplemented with FeCl_3 (6 μM iron (III))
151 or with H_3BO_3 (0.47 mM) or the corresponding culture supernatant were added to the algal culture. After
152 incubation at $22 \pm 1^\circ\text{C}$ under cool white fluorescent lights, living, non-swimming, group-attacked and
153 lysed phytoplankton cells were counted in a sedimentation chamber under an inverted microscope. The
154 number of lysed cells corresponded to algae showing disrupted membranes. Non-swimming algae were
155 not counted as lysed cells. For *Vibrio* analysis, 100 μL of a 1:10 serial dilution mixture in ENSW (from
156 10^{-2} to 10^{-10}) was plated on *Vibrio* Selective TCBS (thiosulfate-citrate-bile salts-sucrose) agar (in
157 triplicate). After incubation for 24 h at $22 \pm 1^\circ\text{C}$, the number of living *Vibrio* cells was determined by
158 counting colony-forming units (CFUs). Data are the means of three independent experiments.

159

160 **Microscope observations**

161 The dynamic of the interaction between *A. pacificum* ACT03 and *V. atlanticus* LGP32, which is an oyster
162 pathogen isolated from the French Atlantic coast (Gay et al., 2004) and present in the Thau Lagoon, south
163 of France (Lopez-Joven et al., 2018)) was surveyed. As the algal strain used in the study is not axenic,
164 means that additional bacteria, other than the *V. atlanticus* LGP32, are present in the experiments.
165 To elucidate the interaction without thoroughly accounting for the non-axenic cells, interaction was
166 observed under a Zeiss Axio upright fluorescence microscope equipped with an AxioCamMRm 2 digital
167 microscope camera using *V. atlanticus* LGP32 tagged with green fluorescent protein (GFP). Lasers were

168 used at excitation wavelength (λ_{ex}) 488 nm for GFP (emission wavelengths (λ_{em}): 505–530 nm) and λ_{ex}
169 532 nm for plankton chloroplasts (λ_{em} 560–630 nm). Images were taken sequentially to avoid cross-
170 contamination between fluorochromes. Sequences of images were merged during the *Vibrio-Alexandrium*
171 interaction using ZEN 2012 (blue edition) software. Interaction events between *Vibrio* strains and
172 phytoplankton strains were also observed under a Leica TCS SPE confocal laser scanning system connected
173 to a Leica DM 2500 upright microscope camera (Montpellier RIO Imaging Platform, University of
174 Montpellier, France).

175 **Comparative proteomic analysis**

176 *Vibrio sampling and protein extraction.* *V. atlanticus* LGP32 was grown for 60 h at 22°C in artificial
177 seawater (high nutrient stress, cond. 1) or 12 h at 22°C in Zobell media (low nutrient stress, cond. 2). After
178 10 min centrifugation at 8000 rpm, crude protein extracts of *V. atlanticus* LGP32 in each culture condition
179 (triplicates) were obtained by sonication on ice at 20% amplitude for 20 s in 200 μL of ice-cold denaturing
180 buffer (7 M urea, 2 M thiourea, 4% CHAPS in 30 mM Tris-HCl, pH 8.5) and clarified by centrifugation at
181 2000 $\times g$, 15 min, 4°C. The protein concentration of the supernatant was estimated using the 2D Quant Kit
182 (Cytiva™, MERCK) and samples were stored at -80°C until use.

183
184 *Bi-dimensional gel electrophoresis (2D gel).* Proteins extracts were individually analysed on 2D gel
185 electrophoresis (6 gels per condition each corresponding to different biological replicates). To do so, 100
186 μg of proteins from each extract was added to rehydration buffer (7 M urea, 2 M thiourea, 4% CHAPS, 65
187 mM DTT) for a total volume of 350 μL . They were then individually loaded onto 17 cm isoelectric
188 focusing strips (Bio-Rad) with a stabilized non-linear pH ranging from 3 to 10. Due to the high
189 complexity of the protein profile in the acidic part (left) of the gel pH 3–10 (Fig. S3), we conducted
190 additional ‘close-up’ analyses in gels using 17 cm isoelectric focusing strips (Bio-Rad) with a narrower,
191 stabilized pH gradient ranging from 4 to 7. Strips were rehydrated passively for 5 h at 22°C, followed by

192 active rehydration for 14 h under a 50 V current at 22°C (to help large proteins enter the strips).
193 Thereafter, isoelectrofocusing was carried out using the following programme: 50 V for 1 h, 250 V for 1
194 h, 8000 V for 1 h and a final step at 8000 V for a total of 140 000 V.h with a slow ramping voltage
195 (quadratic increasing voltage) at each step. Focused proteins were reduced by incubating the strip twice in
196 equilibration buffer (1.5 M Tris, 6 M urea, 2% SDS, 30% glycerol; bromophenol blue, pH 8.8) containing
197 DTT (130 mM) at 55°C. Then, they were alkylated by incubation with equilibration buffer containing
198 iodoacetamide (135 mM) on a rocking agitator (400 rpm) at room temperature protected from light.
199 Proteins were also separated according to their molecular weight (second dimension) on 12%
200 acrylamide/0.32% piperazine diacrylamide gels run at 25 mA per gel for 30 min followed by 75 mA per
201 gel for 8 h using a Protean II XL system (Bio-Rad). Gels were stained using an MS-compatible silver
202 staining protocol and scanned using a ChemiDoc MP Imaging System (Bio-Rad) associated with Image
203 Lab software version 4.0.1 (Bio-Rad).

204
205 *Comparative bioinformatics analysis of 2D gels.* Twelve gels (six per condition) were selected for
206 comparative analysis on PD-Quest v. 7.4.0 (Bio-Rad) to identify changes in protein abundance between
207 the proteomic profiles of *V. atlanticus* LGP32 cultured in contrasting nutrient conditions (ENSW/Zobell).
208 Spots whose mean intensity across six replicates per strain was two times higher or lower than those from
209 the other strain, with a $P < 0.01$ (Mann-Whitney U-test), were considered significantly different in terms of
210 abundance between the two conditions (quantitative difference). Differentially represented spots were then
211 excised from the gels, destained, trypsin-digested and the obtained peptides were identified by tandem
212 mass spectrometry (MS-MS) using the PISSARO platform facility (University of Rouen, France). To
213 identify protein(s) present in each spot, the obtained peptides were compared with *V. atlanticus* LGP32
214 reference genome (<https://vibrio.biocyc.org/>). The genes whose peptides matched strongly were retrieved
215 and used for an BLASTx query against non-redundant databases to determine the protein identity of the
216 best match. A gene was considered as strongly matched when at least two peptides matched the sequence

217 with a coverage of > 6%. Their theoretical isoelectric point (pI) and molecular weights were also
218 calculated using the ExPasy server (<https://www.expasy.org/>) to compare them with the location of the
219 spot on the gel. Altogether, these complementary analyses made it possible to characterize the protein
220 identity of each spot with confidence.

221

222 **Gene expression analysis**

223 *Vibrio sampling and RNA extraction.* *V. atlanticus* LGP32 was grown in Zobell media for 36 h and 60 h at
224 22°C (decline phase of growth, nutrient stress) or for 12 h at 22°C (exponential grow phase, poor nutrient
225 stress). Total RNA was isolated from *V. atlanticus* LGP32 using the standard TRIzol method (Invitrogen
226 Life Technologies SAS, Saint-Aubin, France) and then treated with DNase (Invitrogen) to eliminate
227 genomic DNA contamination. After sodium acetate precipitation, the quantity and quality of the total
228 RNA were determined using a NanoDrop spectrophotometer and agarose gel electrophoresis. Following
229 heat denaturing (70°C for 5 minutes), reverse transcription was performed using 1 µg of RNA prepared
230 with 50 ng µL⁻¹ oligo-(dT) 12–18 mer in a 20-µL reaction containing 1 mM dNTPs, 1 unit µL⁻¹
231 RNaseOUT and 200 units µL⁻¹ Moloney murine leukaemia virus reverse transcriptase (M-MLV RT) in
232 reverse transcriptase buffer, according to the manufacturer's instructions (Invitrogen Life Technologies
233 SAS, Saint-Aubin, France).

234

235 *PCR amplification.* Amplification reactions were analysed using a Roche LightCycler 480 Real-Time
236 thermocycler (Bio-Environnement platform, University of Perpignan, France). In this study, several PCR
237 primer pairs were designed using Primer3 software (optimal primer size: 20 bases; Tm: 60°C; primer
238 GC%: 50; 2GC clamp and product size range: 150–200 bp) and calibrated with *V. atlanticus* LGP32
239 genomic DNA (Table 1). To determine the qPCR efficiency of each primer pair used, standard curves
240 were generated using seven serial dilutions of genomic DNA (10⁻², 10⁻³, 10⁻⁴, 10⁻⁵, 10⁻⁵, 10⁻⁷ and 10⁻⁸) (not

241 shown); the qPCR efficiencies of the tested genes varied between 1.85 and 2.08 (Table 3). For gene
242 expression, reverse transcription was performed with 1 μ g of total RNA using random hexamers and
243 SuperScript IV reverse transcriptase (Invitrogen). The total qPCR reaction volume was 10 μ L and
244 consisted of 5 μ L of cDNA (diluted 1:5), 2.5 μ L of SensiFAST SYBR No-ROX Mix (Bioline) and 100
245 nM or 300 nM PCR primer pair (Table 3). The reaction parameters were as follows: 2 min at 95°C (initial
246 denaturation) and 40 cycles of 5 s at 95°C (denaturation), 10 s at 59°C (annealing) and 20 s at 72°C
247 (elongation). The specificity of each PCR was checked by measuring fluorescent signals during melting
248 curve analysis (PCR product heated from 65°C to 95°C continuously and slowly at 0.1°C s⁻¹). Relative
249 expression was calculated by normalization to the expression of two constitutively expressed
250 housekeeping genes, namely, 6PKF (VS_2913) and CcmC (VS_0852), using the delta-delta threshold
251 cycle ($\Delta\Delta$ Ct) method (Pfaffl, 2001).

252

253 **Detection of quorum-sensing signalling molecules.**

254 *Vibrio culture.* To detect the QS molecules (AI-2, AI-1 and CAI-1) *V. atlanticus* LGP32 was grown
255 in Zobell media for 12 h (exponential growth phase, control) and 60 h (decline phase of growth, nutrient
256 stress).

257 *AI-2 analysis.* Bioluminescence assay using the QS bioluminescent of *Vibrio campbellii* MM32
258 (*luxN::Cm, luxS::Tn5Kan*) was used to detect AI-2 molecules in culture supernatants. Briefly, *Vibrio*
259 cultures were centrifuged at 17,000 x g for 10 min, and the resulting supernatants were filtered on 0.22
260 μ m. Then 20 μ L of the filtrates were mixed with 180 μ L of *V. campbellii* MM32 diluted 1:5000 then
261 incubated at 30°C and 100 rpm. Luminescence and cell density (OD₆₂₀) were collected in triplicate and
262 analysed according to Tourneroche et al. (Tourneroche et al., 2019).

263

264 *AI-1 and CAI-1 extraction and LC-MS analysis.* Chemical analyses were conducted with a Q Exactive
265 Focus Orbitrap System coupled to an Ultimate 3000™ ultrahigh-performance liquid chromatography
266 (UHPLC) system (Thermo Fisher Scientific) according to Rodrigues et al. (Rodrigues et al., 2022).
267 Briefly, ethyl acetate (2 mL) was added into each culture (2 mL). This mixture was shaken overnight at
268 room temperature (150 rpm). The two phases were then separated and the aqueous phase was extracted
269 once again. The two obtained organic phases were pooled and the solvent was evaporated under vacuum.
270 The crude extracts were dissolved in 500 µL LC-MS grade methanol for analysis. All experiments were
271 conducted in triplicate. Analyses of extracts and standards (3 µL injected) were performed in electrospray
272 positive ionization mode in the 50–750 m/z range in centroid mode. The parameters were as follows:
273 spray voltage: 3 kV; sheath flow rate: 75; aux gas pressure: 20; capillary temperature: 350°C; heater
274 temperature: 430°C. The analysis was conducted in Full MS data-dependent MS2 mode (Discovery
275 mode). Resolution was set to 70,000 in Full MS mode, and the AGC (automatic gain control) target was
276 set to 1×10^6 . In MS2, resolution was 17,500, AGC target was set to 2×10^5 , isolation window was 0.4 m/z,
277 and normalized collision energy was stepped to 15, 30 and 40 eV. The UHPLC column was a
278 Phenomenex Luna Omega Polar C18 1.6 µm, 150 x 2.1 mm. The column temperature was set to 42°C,
279 and the flow rate was 0.4 mL min⁻¹. The solvent system was a mixture of water (A) with increasing
280 proportions of acetonitrile (B), with both solvents modified with 0.1% formic acid. The gradient was as
281 follows: 1% B 3 min before injection, then from 1 to 15 min, a gradient increase of B up to 100% (curve
282 5), followed by 100% B for 5 min. The flow was injected into the mass spectrometer starting immediately
283 after injection. All data were acquired and processed using FreeStyle 1.5 software (Thermo Fisher
284 Scientific).

285
286 *Chemicals and solvents.* N-acyl-homoserine lactones (AHL) were obtained from Cayman Chemical (Ann
287 Arbor, MI, USA). Stock solutions were obtained by dissolving standards in methanol or dichloromethane
288 (C18-AHL) at a concentration of 1 mg mL⁻¹ and stored at -80°C. Standard solutions for UHPLC- high-

289 resolution tandem mass spectrometry (HRMS) analyses were prepared by diluting each individual
290 standard solution with methanol in order to obtain a concentration range from 2000 to 20 ng mL⁻¹. LC-MS
291 grade methanol, acetonitrile and formic acid were purchased from Biosolve (Biosolve Chimie, Dieuze,
292 France), analytical-grade ethyl acetate was obtained from Sigma-Aldrich. Pure water was obtained from
293 Elga Purelab Flex System (Veolia LabWater STI, Antony, France).

294

295 **Nature of lytic compounds secreted by *V. atlanticus* LGP32.**

296 To determine the nature of the lytic compounds secreted, *V. atlanticus* LGP32 grown for 60 h in Zobell
297 media at 22 °C was filtered through a 10 kDa membrane (Amicon[®] Ultra-4 filter unit). The eluate
298 containing molecules with MW below 10 KDa was then incubated in a water bath at 100°C for 30 min.
299 Boiled filtrates (0.1 mL) were subsequently used to inoculate *A. pacificum* (ACT03 strain) cultures, then
300 lytic activity was observed under the Leica TCS SPE confocal laser scanning system. Zobell media with
301 the same treatment was used as control.

302

303 **Statistical analysis.***Environmental data.* Statistical analyses were performed using R 3.6.3 software. The
304 relationship between *Alexandrium* and *Vibrio* was explored separately in spring and autumn. We used a
305 generalized linear model specifying a Gaussian family. For spring, the dataset for salinity and temperature
306 was complete (14 periods of observation). One influential period was detected and removed from the
307 dataset whenever no dead *Alexandrium* cells were observed. The effects of explanatory variables such as
308 log₁₀ (Vibrio+1), salinity and temperature were centred, reduced and tested as fixed effects with a linear
309 relationship. Model selection was performed using the Akaike information criterion corrected for small
310 sample size (AIC_c). Models were considered different whenever the difference between their AIC_c value
311 and the lowest AIC_c value (Δ AIC_c) was lower than 2 (Burnham and Anderson, 2002). *Alexandrium*

312 distribution and model residuals were checked for normal distribution assumptions (QQ plot and Shapiro-
313 Wilk test). For autumn, the dataset was complete for 10 periods of observation. Salinity and temperature
314 were missing for three periods. We explored the relationship between *Alexandrium* and *Vibrio* alone using
315 the method detailed above for spring.

316 *In vitro* data. Statistical analyses were performed using one-way ANOVA (analysed by pair) followed by
317 Tukey's test (Statistica 10.0 software, StatSoft, Maison-Alfort, France). *P <0.05, **P <0.01, ***P
318 <0.001

319 RESULTS

320 **Concomitant occurrence of *A. pacificum* ACT03, *A. tamarense* ATT07 and free-living *Vibrio* spp. in**
321 **the Thau lagoon.** In the spring and autumn of 2015, in the Thau Lagoon (Fig. 1A), we detected
322 *Alexandrium* algae (*A. pacificum* ACT03 and *A. tamarense* ATT07, both alive and in degraded cell forms)
323 and free-living *Vibrio*, but no plankton-associated *Vibrio* were observed (Fig. 1B, Data S1). Using model
324 selection based on AICc, we found no significant relationship between *Alexandrium* (*A. pacificum*
325 ACT03, *A. tamarense*) and *Vibrio* Spp. abundances in autumn. This result is consistent with the difficulty
326 that *Vibrio* has in growing at temperatures below 20°C and with the complex interacting factors driving
327 bloom dynamics (Laanaia et al., 2013). Interestingly, in spring 2015, the mean densities of *Alexandrium*
328 and of free-living *Vibrio* were positively correlated. The lowest AICc was obtained with the model
329 explaining degraded form of *Alexandrium* density based on the free *Vibrio* density (Fig. 1C). Given that,
330 this model is not so different from the model with only the intercept, but better than any other linear
331 combination with other potentially interfering drivers, such as temperature and salinity (Fig. 1D), we
332 searched for evidence of a relationship between *Vibrio* and *Alexandrium* by studying their interaction *in*
333 *vitro*.

334

335 ***Vibrio atlanticus* LGP32 feeds on *Alexandrium pacificum* ACT03.**

336 To investigate whether *Alexandrium* interacts with *Vibrio*, we incubated in Enriched Natural SeaWater
337 (ENSW) *A. pacificum* ACT03 with *V. atlanticus* LGP32 previously grown for 12 hours in Zobell media.
338 In interaction with *V. atlanticus* LGP32, *A. pacificum* ACT03 cell abundance decreased significantly from
339 2.10×10^4 cells mL⁻¹ to 1.07×10^4 cells mL⁻¹ after 48 h (Fig. 2A), while the *V. atlanticus* LGP32
340 concentration grew significantly after 26 h of interaction, reaching a maximum peak density of 7.67×10^7
341 CFU mL⁻¹ at 34 h (Fig. 2B). In the control experiment where *A. pacificum* ACT03 was cultured alone in
342 ENSW, the algal concentration remained stable over time (Fig. 2A) and no bacteria were on the
343 corresponding TCBS plates (*Vibrio* selective medium). In the control where *V. atlanticus* LGP32 was
344 cultured alone in ENSW, the concentration of the bacteria tended to increase during the first 24 h (from
345 2.0×10^5 to 7.0×10^5 CFU mL⁻¹) and then decreased at 48h, below the initial starting concentration (Fig.
346 2B). These results show that the interaction between *V. atlanticus* LGP32 and *A. pacificum* ACT03 leads
347 to a decline in the algal population and promotes the growth of *V. atlanticus* LGP32. This suggests that *V.*
348 *atlanticus* LGP32 is able to feed on *A. pacificum* ACT03.

349 ***V. atlanticus* LGP32 performs group-attacks on *A. pacificum* ACT03.**

350 Epifluorescence microscopy observation of GFP-labelled *V. atlanticus* LGP32 (previously grow in Zobell
351 medium) in interaction showed that *A. pacificum* ACT03 cells that had lost their motility were attacked
352 individually by groups of *V. atlanticus* LGP32 before being lysed (Movie 1). Attacks were extremely
353 rapid, with empty thecae (algal envelopes) observed in the medium after less than 60 s (Movie 2). During
354 the attack, *V. atlanticus* LGP32 did not invade the algal cell but remained clustered on the cell surface
355 (Fig. 3A2).

356 **Group attack of *A. pacificum* ACT03 is activated by *V. atlanticus* LGP32 starvation.**

357 In 2002, Martin hypothesized that nutritional stress induces bacteria to lyse algae (Martin, 2002). To test
358 this hypothesis, we monitored *V. atlanticus* LGP32 behaviour in response to starvation (Fig. 3). We

359 observed that *V. atlanticus* LGP32 in exponentially growth phase (12 h of culture in Zobell medium) did
360 not interact with *A. pacificum* ACT03 cells for the first hour of contact (Fig. 3A, B, C). In contrast, *V.*
361 *atlanticus* LGP32 in the decline phase (36 h of culture in Zobell medium) induced a significant decrease in
362 the number of motile algae cells by 8.9% after 15 min and by 43.3% after 60 min (Fig. 3A). This
363 phenomenon corresponded to the degradation and/or disruption of algal flagella (Movie 3). The flagella no
364 longer functioned correctly, which caused erratic swimming of the algae (Video 3, left cell). This was
365 followed by a complete cessation of swimming. When the flagellum detached from the algae (Video 3,
366 right cell), the attack occurred. With starved *V. atlanticus* LGP32 (60 h of culture in Zobell medium),
367 algae immobilization was fast and significant (91.4% in 15 min, Fig. 3A), and algae were attacked
368 individually being targeted by groups of *V. atlanticus* LGP32 (Movie 4). The percentage of cells attacked
369 and killed peaked at 30% after 15–30 min of contact (Fig. 3B) and then decreased. After 1 h, attacks had
370 stopped with approximately 40% of the algal cells still alive (Fig. 3C). Although it remains unclear
371 whether the attacks occur during a specific phase of growth, it is evident that the cells are already
372 weakened before attack as they have all lost their flagella. An old-starved culture of *V. atlanticus* LGP32
373 (126 h) significantly immobilized *A. pacificum* ACT03 cells within a few minutes, with lysis occurring
374 immediately (Fig. 3A, C), making it impossible to detect attacks by *V. atlanticus* LGP32 (Fig. 3B). The
375 lysis phase corresponded to initial vesicle formation followed by the bursting of *A. pacificum* ACT03 cells
376 (Movie 5) and was induced by the old-starved culture supernatant of *V. atlanticus* LGP32 (Fig. S1).
377 We next tested whether this lytic effect was mediated by thermostable molecule (s) secreted by *Vibrio*.
378 The culture supernatant of starved culture of *V. atlanticus* LGP32 (36 h) filtered through a 10 kDa
379 membrane and then incubated at 100°C for 30 min still possessed its lytic properties, indicating that the
380 algicidal compounds produced by *V. atlanticus* LGP32 are small thermostable molecules unlikely to be
381 lytic enzymes, or lysins able to digest the algae cell. However, these experimental observations clearly
382 show the key role of nutrient limitation in triggering the ‘group-attack’ behaviour and the secretion of lytic
383 compounds of *V. atlanticus* LGP32.

384 **Group attack occurs on *A. pacificum* ACT03 in exponential phase of growth.**

385 Here, we wondered whether the live/dead status of algae is important for *V. atlanticus* LGP32-mediated
386 group attacks targeting. To this aim, *A. pacificum* ACT03 in exponential growth phase was first exposed
387 for 30 min to supernatant from 60 hours starved *V. atlanticus* LGP32 Zobell media that induced 25% lysis
388 of *A. pacificum* ACT03 cells and next to the corresponding *V. atlanticus* LGP32 cells. Group attacks were
389 observed on non-degraded *A. pacificum* ACT03 cells, but not on lysed cells. This result is similar to what
390 is observed on the Movie 1 with flash attacks only on immobilized, but not degraded *A. pacificum* ACT03
391 cells (red), and not on lysed cells (green). In addition, no group attacks occurred on cells from an old *A.*
392 *pacificum* ACT03 culture (1-month culture). Together, with the very short duration of group attacks
393 (Movie 1), these results indicate that *V. atlanticus* LGP32 group-attacked exponentially growing cells of
394 *A. pacificum* ACT03, but not decomposing algae, suggesting that this behaviour is not just an
395 opportunistic response of heterotrophic bacteria to organic substrates.

396

397 **Group attack is independent of quorum sensing.**

398 Considering the coordinated action of *V. atlanticus* LGP32 in attacks on *A. pacificum* ACT03, we tested
399 whether the group-attack process depended on the key physiological mechanism that regulates many
400 functions in marine microbial cells, quorum sensing (QS) (Lami, 2019; Papenfort and Bassler, 2016), a
401 type of cell-cell communication. Although QS is a cell-density-dependent mechanism, our results showed
402 no attack from a 12 h culture of *V. atlanticus* LGP32 up to a concentration of $4 \cdot 10^6$ *Vibrio* mL⁻¹ (Fig 4A).
403 Attacks were only observed with *V. atlanticus* LGP32 from a 60 h culture at low concentration of $5 \cdot 10^3$
404 *Vibrio* mL⁻¹ to the highest concentration tested of $5 \cdot 10^5$ *Vibrio* mL⁻¹ (Fig 4A), consistent with the
405 hypothesis that the attacks were independent of *Vibrio* density.

406 The analysis of the expression of genes involved in the known QS pathways in *Vibrio* cell (Fig. S3A),
407 highlighted that only the AI-2 pathway was induced during nutrient stress of *V. atlanticus* LGP32, because
408 only the expression of the AI synthase (*LuxS*) and its receptor (*LuxP*) increased significantly (Fig. 4B;
409 ANOVA $p < 0.05$). This was confirmed by a QS bioluminescence assay, which showed a AI-2 molecules
410 (unquantified) in the Zobell culture supernatant of *V. atlanticus* LGP32 after 60 h of culture but not after
411 12 h of culture and not in the ENSW supernatant of *V. atlanticus* LGP32 after 12 or 60 h of culture.
412 UHPLC-HRMS/MS provided no evidence of detectable HAI-1) and CAI-1 in any experiments.
413 By targeted mutagenesis of key genes involved in QS pathways $\Delta luxM$ (HAI-1 production), $\Delta luxS$ (AI-2
414 production) and $\Delta luxR$ (high-density QS master regulator) did not lead to any change in the group-attack
415 behaviour of *V. atlanticus* LGP32 (Fig. 4C). Taken together these results showed that group-attack by *V.*
416 *atlanticus* LGP32 is not coordinated by QS.

417 **Group attack related to the availability of iron.**

418 The comparative analysis of the proteome of *V. atlanticus* LGP32 incubated 60 h in artificial seawater
419 (ENSW) versus *V. atlanticus* LGP32 grown 12 h in Zobell nutrient-rich medium revealed 10 proteins
420 modulated by nutrient stress (Fig. S2). The two most down-regulated proteins correspond to be β -
421 ketoacyl-(acyl-carrier-protein) synthase II (-22-fold in ENSW compared to Zobell), a key regulator of
422 bacterial fatty acid synthesis, and the dihydroorotase (-6.6-fold in ENSW compared to Zobell), an enzyme
423 essential for pyrimidine biosynthesis and thus bacterial proliferation and growth. The low expression of
424 these proteins in ENSW is consistent with *V. atlanticus* LGP32 nutritional starvation. The most up-
425 regulated protein in starved *V. atlanticus* LGP32, with an increase of more than 6-fold, was glucosamine-
426 6-phosphate deaminase, an enzyme involved in bacterial energy metabolism probably necessary for its
427 survival. Among the other up-regulated proteins, one was an iron siderophore-binding protein (Spot 4413,
428 Fig. S2A) corresponding to the vibrioferrin outer membrane receptor PvuB, whose gene is part of the *pvu*
429 operons involved in iron transport (Fig. S3B). Interestingly, the corresponding gene *pvuB* as well as the

430 vibrioferrin membrane receptor gene *pvuA2* (Fig. S3B) were both significantly induced in *Vibrio* under
431 nutrient stress (Fig. 4B; ANOVA $p < 0.01$) but not the one involved in the vibrioferrin biosynthesis, *pvsA*
432 (Fig. 4B). Remarkably, among the 10 proteins identified by proteomic analysis only *V. atlanticus* LGP32
433 mutant lacking *pvuB* failed to attack *A. pacificum* ACT03 (Fig. 4C; ANOVA $p < 0.001$). In the absence of
434 the *pvuB* gene, *V. atlanticus* LGP32 was unable to attack in group *A. pacificum* ACT03. In addition, *V.*
435 *atlanticus* LGP32 cells that had been washed with ENSW to remove their culture supernatant metabolites
436 also failed to attack *A. pacificum* ACT03 (Fig. 4C; ANOVA $p < 0.001$), which is congruent with the
437 hypothesis that attacks depend on the *V. atlanticus* LGP32 vibrioferrin transport system. Finally, group-
438 attacks increased significantly, when FeCl_3 was added to the *Vibrio* culture medium (Fig. 4D) but not with
439 H_3BO_4 (Fig. 4D). Taken together, those results are consistent with the hypothesis that attacks are regulated
440 by iron.

441 **Group attack is a *Vibrio spp.* behavior specific to *Alexandrium spp.***

442 To evaluate the dinoflagellates specificity of the group-attack behaviour, a selection of *Vibrio spp.* was co-
443 cultured with a selection of dinoflagellate strains commonly found in the Mediterranean Sea. The results
444 showed that, among the *Vibrio spp.* tested (pathogenic or not) all, under nutrient stress, were able to
445 secrete algicidal compounds, immobilize, ‘group-attack’ and lyse *A. pacificum* ACT03 cells (Table 1) and
446 no tropism linked to their pathogenesis for fish or invertebrates was observed (Table 1). Among the
447 sixteen dinoflagellates species tested, all eight *Alexandrium spp.* (non-toxic and paralytic shellfish toxin
448 (PST) producers) and *Gymnodinium catenatum* (PST producer) were immobilized, ‘group-attacked’ and
449 lysed by *V. atlanticus* LGP32 (Table 2), but no tropism linked to PSTs was revealed (Table 2).

450

451 **DISCUSSION**

452 Predation is a widespread mode of interaction for survival in the natural world (Finke and Denno, 2004;
453 Sinclair et al., 2003). Among predators, predatory bacteria are found in a wide variety of environments,

454 and like bacteriophages and predatory protists, they have been reported to prey exclusively on other
455 bacteria (Johnke et al., 2014; Perez et al., 2016). Considering predator as a free organism that feeds at the
456 expense of another, this study is the first evidence of the capacity of some *Vibrio* to develop a predatory
457 strategy against an alga. This behaviour differs from parasitism, because the survival of *Vibrio* is not
458 exclusively dependent on the algae.

459 The strategy developed by *Vibrio* to kill algae may be reminiscent of strategies previously described in the
460 prokaryotes (Johnke et al., 2014). As shown in Movie 1, the interaction between *V. atlanticus* LGP32 and
461 *A. pacificum* ACT03 proceeds in three stages (Fig. 5). The first stage, the '**immobilization stage**', recalls
462 the strategy used by *Streptomyces* to immobilize its prey (Kumbhar et al., 2014) based on the secretion of
463 algicidal metabolites that disrupt the flagella. The second stage, the '**attack stage**', resembles the 'wolf-
464 pack attack' strategy described for *Myxococcus xanthus* and *Lysobacter* (Martin, 2002; Perez et al.,
465 2014). *V. atlanticus* LGP32 also surrounds *A. pacificum* ACT03 cells at high density for a very short time,
466 but does not invade the algal cell. This phenomenon is comparable to that of bacteria clustering around
467 lysed ciliate cells (Blackburn et al., 1998). The third stage, the '**killing stage**', is similar to that of epibiotic
468 bacterial predators, which induces the lysis of bacteria from the outside (Rashidan and Bird, 2001).

469 Overall, these observations show that *V. atlanticus* LGP32 is able of wolf-pack hunting behaviour.
470 Results showed that quorum sensing does not control algal predation by *Vibrio*, in contrast to iron
471 acquisition systems, which play a major role in regulating this phenotype, as we demonstrated genetically.
472 Indeed, our quorum sensing mutants exhibited the same predatory phenotype as the wild-type *Vibrio*
473 *atlanticus* LGP32, whereas mutants impaired in iron uptake completely lost this phenotype. However,
474 quorum sensing and iron acquisition are sometimes interconnected in *Vibrio* (McRose et al., 2018). For
475 instance, in *Vibrio vulnificus*, the production of vulnibactin (a siderophore) is controlled by AI-2 (Kim and
476 Shin, 2011). Whether the production of AI-2 by *V. atlanticus* LGP32, induced during group attacks of *A.*
477 *pacificum* ACT03, promotes the production of vibrioferrin (Lami, 2019) remains to be investigated.

478 We further show that the group-attack behaviour of *V. atlanticus* LGP32 depends on iron availability. Iron
479 is an essential element for growth in most organisms, including phytoplankton (Martin and Fitzwater,
480 1988) and bacteria (Neilands, 1981), and its concentration in seawater is known to be very low with
481 measurements in the open ocean surface at 0.2 nM and in deep waters at 0.6 nM (Millero, 1998).
482 Moreover, Its low solubility in seawater limits its availability (Bruland et al., 1994; Wu and Luther, 1994).
483 To acquire iron, bacteria have developed systems based on the secretion (and subsequent uptake) of iron-
484 chelating siderophores to obtain this element from the environment (Amin et al., 2009a). Therefore, many
485 *Vibrio spp.* produce a siderophore known as vibrioferrin, which is synthesized and secreted by proteins
486 encoded by the *pvsABCDE* gene cluster (Fig. S3). Boron being known to be a regulator or capable of
487 being transported by vibrioferrin (Romano et al., 2013; Weerasinghe et al., 2013) we tested its potential
488 involvement in the interaction but no effect was evidenced here. For iron-vibrioferrin uptake, *Vibrio*
489 *parahaemolyticus* uses a membrane siderophore receptor, called PvuA, which is coupled to an inner
490 membrane ATP-binding cassette (ABC). This ABC transporter system comprised of proteins encoded by
491 the *pvuABCDE* gene cluster (Fig. S3) is required for transporting the siderophore across the inner
492 membrane (Tanabe et al., 2003). Siderophores are not only iron carriers but also important regulators of
493 virulence (Miethke and Marahiel, 2007) and mediators of bacterial interaction with phytoplankton (Amin
494 et al., 2009b; Kramer et al., 2020). We showed here a pivotal role of iron in the interaction between *V.*
495 *atlanticus* LGP32 and *A. pacificum* ACT03. This mirrors the mutualistic interaction observed between
496 *Gymnodinium catenatum* and *Marinobacter* (Amin et al., 2009b). In fact, in natural settings, the co-
497 occurrence of *Marinobacter* and *G. catenatum* is suggested to depend on a mutually beneficial utilization
498 of iron and carbon resources (Bolch et al., 2011). As in the present study, iron seems to play a key role in
499 the interaction. Indeed, the labile iron released through the photolysis of ferric chelates with vibrioferrin
500 providing a crucial iron source for phytoplankton, which need substantial amounts of iron to support
501 carbon fixation through photosynthesis (Amin et al., 2009a; Yang et al., 2021). This fixed carbon, in turn,
502 sustains the growth of both the phytoplankton and their bacterial counterparts (Amin et al., 2009b; Kramer

503 et al., 2020). Future study could demonstrate further the role of vibrioferrin in group attack, by adding
504 iron-saturated vibrioferrin to algae-*Vibrio* co-cultures.

505 In the natural environment, associations between bacteria and algae have already been observed (Lopez-
506 Joven et al., 2018; Miller et al., 2005; Rosales et al., 2022; Xu et al., 2022). Although we do not have
507 definitive proof of a predator/prey interaction *in situ* due to confounding environmental variables,
508 environmental data acquired in the Thau Lagoon showed a correlation between *Alexandrium* and *Vibrio*
509 occurrence, suggesting that such interaction could occur in nature. Evolution of such a predatory behavior
510 would provide an important ecological advantage to *Vibrio* to obtain nutrients in environment, where
511 *Alexandrium spp.* and *Gymnodinium catenatum* form blooms.

512 With more than 30 species distributed all over the world (Anderson et al., 2012; Hallegraeff et al., 2021),
513 *Alexandrium spp.* and *Gymnodinium spp.* considered as invasive species by the Delivering Alien Invasive
514 Species Inventories for Europe (<http://www.europe-aliens.org>)), could play an unexpected and important
515 role in maintaining, structuring and regulating *Vibrio* populations in the ecosystem. In turn, *Vibrio* could
516 contribute to the regulation and control of their blooms.

517 To conclude, this study reveals the capacity of some *Vibrio spp.* to be facultative predators that hunt
518 specific algae. In the current context of climate change favourable to their development monitoring the
519 invasive algae, *Alexandrium spp.* and *Gymnodinium catenatum* should be considered not only for their
520 potent harmful effect on humans and animals, but also as a potential source of nutrients for the expansion
521 of *Vibrio*, particularly pathogenic species (Lemire et al., 2015).

522

523 **DATA AVAILABILITY**

524 All data on environmental study are included in the article and/or supporting information files. All other
525 study data are available on request from the corresponding author.

526

527 **REFERENCES**

- 528 Abadie, E., Amzil, Z., Belin, C., Comps, M.A., Elzière-Papayanni, P., Lassus, P., le Bec, C., Baut, M.L., Nezan,
529 E., Poggi, R., n.d. Contamination de l'étang de Thau par *Alexandrium tamarense* (No. Rapport
530 884). Ifremer.
- 531 Abadie, E., Kaci, L., Berteaux, T., Hess, P., Sechet, V., Masseret, E., Rolland, J.L., Laabir, M., 2015. Effect of
532 Nitrate, Ammonium and Urea on Growth and Pinnatoxin G Production of *Vulcanodinium*
533 *rugosum*. *Mar. Drugs* 13, 5642–5656. <https://doi.org/10.3390/md13095642>
- 534 Amin, S.A., Green, D.H., Hart, M.C., Kuepper, F.C., Sunda, W.G., Carrano, C.J., 2009a. Photolysis of iron-
535 siderophore chelates promotes bacterial-algal mutualism. *Proc. Natl. Acad. Sci. U. S. A.* 106,
536 17071–17076. <https://doi.org/10.1073/pnas.0905512106>
- 537 Amin, S.A., Green, D.H., Kuepper, F.C., Carrano, C.J., 2009b. Vibrioferrin, an Unusual Marine Siderophore:
538 Iron Binding, Photochemistry, and Biological Implications. *Inorg. Chem.* 48, 11451–11458.
539 <https://doi.org/10.1021/ic9016883>
- 540 Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012. The
541 globally distributed genus *Alexandrium*: Multifaceted roles in marine ecosystems and impacts on
542 human health. *Harmful Algae* 14, 10–35. <https://doi.org/10.1016/j.hal.2011.10.012>
- 543 Baker-Austin, C., Trinanes, J., Gonzalez-Escalona, N., Martinez-Urtaza, J., 2017. Non-Cholera *Vibrios*: The
544 Microbial Barometer of Climate Change. *Trends Microbiol.* 25, 76–84.
545 <https://doi.org/10.1016/j.tim.2016.09.008>
- 546 Ben-Gharbia, H., Yahia, O.K.-D., Amzil, Z., Chomerat, N., Abadie, E., Masseret, E., Sibat, M., Triki, H.Z.,
547 Nouri, H., Laabir, M., 2016. Toxicity and Growth Assessments of Three Thermophilic Benthic
548 Dinoflagellates (*Ostreopsis cf. ovata*, *Prorocentrum lima* and *Coolia monotis*) Developing in the
549 Southern Mediterranean Basin. *Toxins* 8, 297. <https://doi.org/10.3390/toxins8100297>
- 550 Blackburn, N., Fenchel, T., Mitchell, J., 1998. Microscale nutrient patches in planktonic habitats shown by
551 chemotactic bacteria. *Science* 282, 2254–2256. <https://doi.org/10.1126/science.282.5397.2254>
- 552 Bolch, C.J.S., Subramanian, T.A., Green, D.H., 2011. THE TOXIC DINOFLAGELLATE *GYMNODINIUM*
553 *CATENATUM* (DINOPHYCEAE) REQUIRES MARINE BACTERIA FOR GROWTH¹. *J. Phycol.* 47, 1009–
554 1022. <https://doi.org/10.1111/j.1529-8817.2011.01043.x>
- 555 Bruland, K., Orians, K., Cowen, J., 1994. Reactive Trace-Metals in the Stratified Central North Pacific.
556 *Geochim. Cosmochim. Acta* 58, 3171–3182. [https://doi.org/10.1016/0016-7037\(94\)90044-2](https://doi.org/10.1016/0016-7037(94)90044-2)
- 557 Brumfield, K.D., Usmani, M., Chen, K.M., Gangwar, M., Jutla, A.S., Huq, A., Colwell, R.R., 2021.
558 Environmental parameters associated with incidence and transmission of pathogenic *Vibrio spp.*
559 *Environ. Microbiol.* 23, 7314–7340. <https://doi.org/10.1111/1462-2920.15716>
- 560 Burkholder, J.M., Shumway, S.E., Glibert, P.M., 2018. Food Web and Ecosystem Impacts of Harmful
561 Algae, in: *Harmful Algal Blooms*. John Wiley & Sons, Ltd, pp. 243–336.
562 <https://doi.org/10.1002/9781118994672.ch7>
- 563 Burnham, K.P., Anderson, D.R. (Eds.), 2002. Summary, in: *Model Selection and Multimodel Inference: A*
564 *Practical Information-Theoretic Approach*. Springer, New York, NY, pp. 437–454.
565 https://doi.org/10.1007/978-0-387-22456-5_8
- 566 Chai, Z.Y., Wang, H., Deng, Y., Hu, Z., Tang, Y.Z., 2020. Harmful algal blooms significantly reduce the
567 resource use efficiency in a coastal plankton community. *Sci. Total Environ.* 704, 135381.
568 <https://doi.org/10.1016/j.scitotenv.2019.135381>
- 569 Coyne, K.J., Wang, Y., Johnson, G., 2022. Algicidal Bacteria: A Review of Current Knowledge and
570 Applications to Control Harmful Algal Blooms. *Front. Microbiol.* 13, 871177.
571 <https://doi.org/10.3389/fmicb.2022.871177>

- 572 Drebes Dörr, N.C., Blokesch, M., 2020. Interbacterial competition and anti-predatory behaviour of
573 environmental *Vibrio cholerae* strains. *Environ. Microbiol.* 22, 4485–4504.
574 <https://doi.org/10.1111/1462-2920.15224>
- 575 Espinoza-Vergara, G., Hoque, M.M., McDougald, D., Noorian, P., 2020. The Impact of Protozoan
576 Predation on the Pathogenicity of *Vibrio cholerae*. *Front. Microbiol.* 11, 17.
577 <https://doi.org/10.3389/fmicb.2020.00017>
- 578 Fertouna-Bellakhal, M., Dhib, A., Fathalli, A., Bellakhal, M., Chomérat, N., Masseret, E., Laabir, M., Turki,
579 S., Aleya, L., 2015. *Alexandrium pacificum* Litaker sp. nov (Group IV): Resting cyst distribution and
580 toxin profile of vegetative cells in Bizerte Lagoon (Tunisia, Southern Mediterranean Sea). *Harmful*
581 *Algae* 48, 69–82. <https://doi.org/10.1016/j.hal.2015.07.007>
- 582 Finke, D.L., Denno, R.F., 2004. Predator diversity dampens trophic cascades. *Nature* 429, 407–410.
583 <https://doi.org/10.1038/nature02554>
- 584 Gay, M., Renault, T., Pons, A.M., Le Roux, F., 2004. Two *Vibrio splendidus* related strains collaborate to
585 kill *Crassostrea gigas*: taxonomy and host alterations. *Dis. Aquat. Organ.* 62, 65–74.
586 <https://doi.org/10.3354/dao062065>
- 587 Genovesi, B., Shin-Grzebyk, M.-S., Grzebyk, D., Laabir, M., Gagnaire, P.-A., Vaquer, A., Pastoureaud, A.,
588 Lasserre, B., Collos, Y., Berrebi, P., Masseret, E., 2011. Assessment of cryptic species diversity
589 within blooms and cyst bank of the *Alexandrium tamarense* complex (Dinophyceae) in a
590 Mediterranean lagoon facilitated by semi-multiplex PCR. *J. Plankton Res.* 33, 405–414.
591 <https://doi.org/10.1093/plankt/fbq127>
- 592 Hadjadj, I., Laabir, M., Frihi, H., Collos, Y., Shao, Z.J., Berrebi, P., Abadie, E., Amzil, Z., Chomérat, N.,
593 Rolland, J.L., Rieuvilleneuve, F., Masseret, E., 2020. Unsuspected intraspecific variability in the
594 toxin production, growth and morphology of the dinoflagellate *Alexandrium pacificum* R.W.
595 Litaker (Group IV) blooming in a South Western Mediterranean marine ecosystem, Annaba Bay
596 (Algeria). *Toxicon* 180, 79–88. <https://doi.org/10.1016/j.toxicon.2020.04.005>
- 597 Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M.-Y.D., Bresnan, E., Chinain, M., Enevoldsen, H.,
598 Iwataki, M., Karlson, B., McKenzie, C.H., Sunesen, I., Pitcher, G.C., Provoost, P., Richardson, A.,
599 Schweibold, L., Tester, P.A., Trainer, V.L., Yñiguez, A.T., Zingone, A., 2021. Perceived global
600 increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts.
601 *Commun. Earth Environ.* 2, 117. <https://doi.org/10.1038/s43247-021-00178-8>
- 602 Harrison, P., Waters, R., Taylor, F., 1980. A Broad-Spectrum Artificial Seawater Medium for Coastal and
603 Open Ocean Phytoplankton. *J. Phycol.* 16, 28–35. [https://doi.org/10.1111/j.1529-](https://doi.org/10.1111/j.1529-8817.1980.tb00724.x)
604 [8817.1980.tb00724.x](https://doi.org/10.1111/j.1529-8817.1980.tb00724.x)
- 605 Johnke, J., Cohen, Y., de Leeuw, M., Kushmaro, A., Jurkevitch, E., Chatzinotas, A., 2014. Multiple micro-
606 predators controlling bacterial communities in the environment. *Curr. Opin. Biotechnol.* 27, 185–
607 190. <https://doi.org/10.1016/j.copbio.2014.02.003>
- 608 Johnson, C.N., 2013. Fitness Factors in Vibrios: a Mini-review. *Microb. Ecol.* 65, 826–851.
609 <https://doi.org/10.1007/s00248-012-0168-x>
- 610 Kim, C.M., Shin, S.H., 2011. Modulation of Iron-Uptake Systems by a Mutation of *luxS* Encoding an
611 Autoinducer-2 Synthase in *Vibrio vulnificus*. *Biol. Pharm. Bull.* 34, 632–637.
612 <https://doi.org/10.1248/bpb.34.632>
- 613 Kitatsukamoto, K., Oyaizu, H., Nanba, K., Simidu, U., 1993. Phylogenetic-Relationships of Marine-
614 Bacteria, Mainly Members of the Family Vibrionaceae, Determined on the Basis of 16s
615 Ribosomal-Rna Sequences. *Int. J. Syst. Bacteriol.* 43, 8–19. [https://doi.org/10.1099/00207713-](https://doi.org/10.1099/00207713-43-1-8)
616 [43-1-8](https://doi.org/10.1099/00207713-43-1-8)

- 617 Kramer, J., Oezkaya, O., Kuemmerli, R., 2020. Bacterial siderophores in community and host interactions.
618 Nat. Rev. Microbiol. 18, 152–163. <https://doi.org/10.1038/s41579-019-0284-4>
- 619 Kumbhar, C., Mudliar, P., Bhatia, L., Kshirsagar, A., Watve, M., 2014. Widespread predatory abilities in
620 the genus *Streptomyces*. Arch. Microbiol. 196, 235–248. [https://doi.org/10.1007/s00203-014-](https://doi.org/10.1007/s00203-014-0961-7)
621 0961-7
- 622 Laabir, M., Jauzein, C., Genovesi, B., Masseret, E., Grzebyk, D., Cecchi, P., Vaquer, A., Perrin, Y., Collos, Y.,
623 2011. Influence of temperature, salinity and irradiance on the growth and cell yield of the
624 harmful red tide dinoflagellate *Alexandrium catenella* colonizing Mediterranean waters. J.
625 Plankton Res. 33, 1550–1563. <https://doi.org/10.1093/plankt/fbr050>
- 626 Laanaia, N., Vaquer, A., Fiandrino, A., Genovesi, B., Pastoureaud, A., Cecchi, P., Collos, Y., 2013. Wind and
627 temperature controls on *Alexandrium* blooms (2000-2007) in Thau lagoon (Western
628 Mediterranean). Harmful Algae 28, 31–36. <https://doi.org/10.1016/j.hal.2013.05.016>
- 629 Labreuche, Y., Le Roux, F., Henry, J., Zatylny, C., Huvet, A., Lambert, C., Soudant, P., Mazel, D., Nicolas, J.-
630 L., 2010. *Vibrio aestuarianus* zinc metalloprotease causes lethality in the Pacific oyster
631 *Crassostrea gigas* and impairs the host cellular immune defenses. Fish Shellfish Immunol. 29,
632 753–758. <https://doi.org/10.1016/j.fsi.2010.07.007>
- 633 Lami, R., 2019. Quorum Sensing in Marine Biofilms and Environments, in: Quorum Sensing, Academic
634 Press. Elsevier, pp. 55–96. <https://doi.org/10.1016/B978-0-12-814905-8.00003-4>
- 635 Le Roux, F., Binesse, J., Saulnier, D., Mazel, D., 2007. Construction of a *Vibrio splendidus* Mutant Lacking
636 the Metalloprotease Gene *vsm* by Use of a Novel Counterselectable Suicide Vector. Appl.
637 Environ. Microbiol. 73, 777–784. <https://doi.org/10.1128/AEM.02147-06>
- 638 Leblad, B.R., Amnhir, R., Reqia, S., Sitel, F., Daoudi, M., Marhraoui, M., Abdellah, M.K.O., Veron, B., Er-
639 Raioui, H., Laabir, M., 2020. Seasonal variations of phytoplankton assemblages in relation to
640 environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species.
641 Harmful Algae 96, 101819. <https://doi.org/10.1016/j.hal.2020.101819>
- 642 Lemire, A., Goudenege, D., Versigny, T., Petton, B., Calteau, A., Labreuche, Y., Le Roux, F., 2015.
643 Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. Isme J.
644 9, 1523–1531. <https://doi.org/10.1038/ismej.2014.233>
- 645 LeRoux, F., Wegner, K.M., Baker-Austin, C., Vezzulli, L., Osorio, C.R., Amaro, C., Ritchie, J.M., Defoirdt, T.,
646 Destoumieux-Garzon, D., Blokesch, M., Mazel, D., Jacq, A., Cava, F., Gram, L., Wendling, C.C.,
647 Strauch, E., Kirschner, A., Huehn, S., 2015. The emergence of *Vibrio* pathogens in Europe:
648 ecology, evolution, and pathogenesis (Paris, 11-12th March 2015). Front. Microbiol. 6, 830.
649 <https://doi.org/10.3389/fmicb.2015.00830>
- 650 Li, D., Zhang, H., Fu, L., An, X., Zhang, B., Li, Y., Chen, Z., Zheng, W., Yi, L., Zheng, T., 2014. A Novel
651 Algicide: Evidence of the Effect of a Fatty Acid Compound from the Marine Bacterium, *Vibrio* sp
652 BS02 on the Harmful Dinoflagellate, *Alexandrium tamarense*. Plos One 9, e91201.
653 <https://doi.org/10.1371/journal.pone.0091201>
- 654 Liu, P.-C., Lee, K.-K., Yii, K.-C., Kou, G.-H., Chen, S.-N., 1996. News & Notes: Isolation of *Vibrio harveyi*
655 from Diseased Kuruma Prawns *Penaeus japonicus*. Curr. Microbiol. 33, 129–132.
656 <https://doi.org/10.1007/s002849900087>
- 657 Lopez-Joven, C., Rolland, J.-L., Haffner, P., Caro, A., Roques, C., Carre, C., Travers, M.-A., Abadie, E.,
658 Laabir, M., Bonnet, D., Destoumieux-Garzon, D., 2018. Oyster Farming, Temperature, and
659 Plankton Influence the Dynamics of Pathogenic *Vibrios* in the Thau Lagoon. Front. Microbiol. 9,
660 2530. <https://doi.org/10.3389/fmicb.2018.02530>

- 661 Mandel, M.J., Stabb, E.V., Ruby, E.G., 2008. Comparative genomics-based investigation of resequencing
662 targets in *Vibrio fischeri*: Focus on point miscalls and artefactual expansions. *Bmc Genomics* 9,
663 138. <https://doi.org/10.1186/1471-2164-9-138>
- 664 Marampouti, C., Buma, A.G.J., De Boer, M.K., 2021. Mediterranean alien harmful algal blooms: origins
665 and impacts. *Environ. Sci. Pollut. Res.* 28, 3837–3851. [https://doi.org/10.1007/s11356-020-](https://doi.org/10.1007/s11356-020-10383-1)
666 10383-1
- 667 Martin, J., Fitzwater, S., 1988. Iron-Deficiency Limits Phytoplankton Growth in the Northeast Pacific
668 Subarctic. *Nature* 331, 341–343. <https://doi.org/10.1038/331341a0>
- 669 Martin, M.O., 2002. Predatory prokaryotes: An emerging research opportunity. *J. Mol. Microbiol.*
670 *Biotechnol.* 4, 467–477.
- 671 Mavian, C., Paisie, T.K., Alam, M.T., Browne, C., Beau De Rochars, V.M., Nembrini, S., Cash, M.N., Nelson,
672 E.J., Azarian, T., Ali, A., Morris, J.G., Salemi, M., 2020. Toxigenic *Vibrio cholerae* evolution and
673 establishment of reservoirs in aquatic ecosystems. *Proc. Natl. Acad. Sci.* 117, 7897–7904.
674 <https://doi.org/10.1073/pnas.1918763117>
- 675 McRose, D.L., Baars, O., Seyedsayamdost, M.R., Morel, F.M.M., 2018. Quorum sensing and iron regulate
676 a two-for-one siderophore gene cluster in *Vibrio harveyi*. *Proc. Natl. Acad. Sci.* 115, 7581–7586.
677 <https://doi.org/10.1073/pnas.1805791115>
- 678 Miethke, M., Marahiel, M.A., 2007. Siderophore-Based Iron Acquisition and Pathogen Control. *Microbiol.*
679 *Mol. Biol. Rev.* 71, 413–451. <https://doi.org/10.1128/MMBR.00012-07>
- 680 Miller, S.D., Haddock, S.H.D., Elvidge, C.D., Lee, T.F., 2005. Detection of a bioluminescent milky sea from
681 space. *Proc. Natl. Acad. Sci. U. S. A.* 102, 14181–14184.
682 <https://doi.org/10.1073/pnas.0507253102>
- 683 Millero, F.J., 1998. Solubility of Fe(III) in seawater. *Earth Planet. Sci. Lett.* 154, 323–329.
684 [https://doi.org/10.1016/S0012-821X\(97\)00179-9](https://doi.org/10.1016/S0012-821X(97)00179-9)
- 685 Muhling, B.A., Jacobs, J., Stock, C.A., Gaitan, C.F., Saba, V.S., 2017. Projections of the future occurrence,
686 distribution, and seasonality of three *Vibrio* species in the Chesapeake Bay under a high-emission
687 climate change scenario. *Geohealth* 1, 278–296. <https://doi.org/10.1002/2017GH000089>
- 688 Neilands, J., 1981. Iron-Absorption and Transport in Microorganisms. *Annu. Rev. Nutr.* 1, 27–46.
689 <https://doi.org/10.1146/annurev.nu.01.070181.000331>
- 690 Oberbeckmann, S., Fuchs, B.M., Meiners, M., Wichels, A., Wiltshire, K.H., Gerdt, G., 2012. Seasonal
691 Dynamics and Modeling of a *Vibrio* Community in Coastal Waters of the North Sea. *Microb. Ecol.*
692 63, 543–551. <https://doi.org/10.1007/s00248-011-9990-9>
- 693 Pal, M., Yesankar, P.J., Dwivedi, A., Qureshi, A., 2020. Biotic control of harmful algal blooms (HABs): A
694 brief review. *J. Environ. Manage.* 268, 110687. <https://doi.org/10.1016/j.jenvman.2020.110687>
- 695 Papenfort, K., Bassler, B.L., 2016. Quorum sensing signal-response systems in Gram-negative bacteria.
696 *Nat. Rev. Microbiol.* 14, 576–588. <https://doi.org/10.1038/nrmicro.2016.89>
- 697 Perez, J., Jimenez-Zurdo, J.I., Martinez-Abarca, F., Millan, V., Shimkets, L.J., Munoz-Dorado, J., 2014.
698 Rhizobial galactoglucan determines the predatory pattern of *Myxococcus xanthus* and protects
699 *Sinorhizobium meliloti* from predation. *Environ. Microbiol.* 16, 2341–2350.
700 <https://doi.org/10.1111/1462-2920.12477>
- 701 Perez, J., Moraleda-Munoz, A., Javier Marcos-Torres, F., Munoz-Dorado, J., 2016. Bacterial predation: 75
702 years and counting! *Environ. Microbiol.* 18, 766–779. <https://doi.org/10.1111/1462-2920.13171>
- 703 Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic*
704 *Acids Res.* 29, e45. <https://doi.org/10.1093/nar/29.9.e45>
- 705 Rashidan, K.K., Bird, D.F., 2001. Role of predatory bacteria in the termination of a cyanobacterial bloom.
706 *Microb. Ecol.* 41, 97–105.

- 707 Rodrigues, A.M.S., Lami, R., Escoubeyrou, K., Intertaglia, L., Mazurek, C., Doberva, M., Pérez-Ferrer, P.,
708 Stien, D., 2022. Straightforward *N*-Acyl Homoserine Lactone Discovery and Annotation by LC-
709 MS/MS-based Molecular Networking. *J. Proteome Res.* 21, 635–642.
710 <https://doi.org/10.1021/acs.jproteome.1c00849>
- 711 Rolland, J.-L., Pelletier, K., Masseret, E., Rieuvilleneuve, F., Savar, V., Santini, A., Amzil, Z., Laabir, M.,
712 2012. Paralytic Toxins Accumulation and Tissue Expression of alpha-Amylase and Lipase Genes in
713 the Pacific Oyster *Crassostrea gigas* Fed with the Neurotoxic Dinoflagellate *Alexandrium*
714 *catenella*. *Mar. Drugs* 10, 2519–2534. <https://doi.org/10.3390/md10112519>
- 715 Romano, A., Trimble, L., Hobusch, A.R., Schroeder, K.J., Amin, S.A., Hartnett, A.D., Barker, R.A., Crumbliss,
716 A.L., Carrano, C.J., 2013. Regulation of iron transport related genes by boron in the marine
717 bacterium *Marinobacter algicola* DG893. *Metallomics* 5, 1025.
718 <https://doi.org/10.1039/c3mt00068k>
- 719 Rosales, D., Ellett, A., Jacobs, J., Ozbay, G., Parveen, S., Pitula, J., 2022. Investigating the Relationship
720 between Nitrate, Total Dissolved Nitrogen, and Phosphate with Abundance of Pathogenic Vibrios
721 and Harmful Algal Blooms in Rehoboth Bay, Delaware. *Appl. Environ. Microbiol.* 88, e00356-22.
722 <https://doi.org/10.1128/aem.00356-22>
- 723 Sinclair, A.R.E., Mduma, S., Brashares, J.S., 2003. Patterns of predation in a diverse predator-prey system.
724 *Nature* 425, 288–290. <https://doi.org/10.1038/nature01934>
- 725 Su, R.Q., Yang, X.R., Zheng, T.L., Tian, Y., Jiao, N.Z., Cai, L.Z., Hong, H.S., 2007. Isolation and
726 characterization of a marine algicidal bacterium against the toxic dinoflagellate *Alexandrium*
727 *tamarensis*. *Harmful Algae* 6, 799–810. <https://doi.org/10.1016/j.hal.2007.04.004>
- 728 Tanabe, T., Funahashi, T., Nakao, H., Miyoshi, S.-I., Shinoda, S., Yamamoto, S., 2003. Identification and
729 Characterization of Genes Required for Biosynthesis and Transport of the Siderophore
730 Vibrioferriin in *Vibrio parahaemolyticus*. *J. Bacteriol.* 185, 6938–6949.
731 <https://doi.org/10.1128/JB.185.23.6938-6949.2003>
- 732 Thompson, F.L., Thompson, C.C., Swings, J., 2003. *Vibrio tasmaniensis* sp. nov., isolated from Atlantic
733 Salmon (*Salmo salar* L.). *Syst. Appl. Microbiol.* 26, 65–69.
734 <https://doi.org/10.1078/072320203322337326>
- 735 Tourneroche, A., Lami, R., Hubas, C., Blanchet, E., Vallet, M., Escoubeyrou, K., Paris, A., Prado, S., 2019.
736 Bacterial–Fungal Interactions in the Kelp Endomicrobiota Drive Autoinducer-2 Quorum Sensing.
737 *Front. Microbiol.* 10, 1693. <https://doi.org/10.3389/fmicb.2019.01693>
- 738 Vanhove, A.S., Rubio, T.P., Nguyen, A.N., Lemire, A., Roche, D., Nicod, J., Vergnes, A., Poirier, A.C.,
739 Disconzi, E., Bachere, E., Le Roux, F., Jacq, A., Charriere, G.M., Destoumieux-Garzon, D., 2016.
740 Copper homeostasis at the host vibrio interface: lessons from intracellular vibrio transcriptomics.
741 *Environ. Microbiol.* 18, 875–888. <https://doi.org/10.1111/1462-2920.13083>
- 742 Wang, X., Li, Z., Su, J., Tian, Y., Ning, X., Hong, H., Zheng, T., 2010. Lysis of a red-tide causing alga,
743 *Alexandrium tamarensis*, caused by bacteria from its phycosphere. *Biol. Control* 52, 123–130.
744 <https://doi.org/10.1016/j.biocontrol.2009.10.004>
- 745 Wang, Y., Li, S., Liu, G., Li, X., Yang, Q., Xu, Y., Hu, Z., Chen, C.-Y., Chang, J.-S., 2020. Continuous
746 production of algicidal compounds against *Akashiwo sanguinea* via a *Vibrio* sp. co-culture.
747 *Bioresour. Technol.* 295, 122246. <https://doi.org/10.1016/j.biortech.2019.122246>
- 748 Weerasinghe, A.J., Amin, S.A., Barker, R.A., Othman, T., Romano, A.N., Parker Siburt, C.J., Tisnado, J.,
749 Lambert, L.A., Huxford, T., Carrano, C.J., Crumbliss, A.L., 2013. Borate as a Synergistic Anion for
750 *Marinobacter algicola* Ferric Binding Protein, FbpA: A Role for Boron in Iron Transport in Marine
751 Life. *J. Am. Chem. Soc.* 135, 14504–14507. <https://doi.org/10.1021/ja406609s>

- 752 Wu, J., Luther, G., 1994. Size-Fractionated Iron Concentrations in the Water Column of the Western
753 North-Atlantic Ocean. *Limnol. Oceanogr.* 39, 1119–1129.
754 <https://doi.org/10.4319/lo.1994.39.5.1119>
- 755 Xu, Q., Wang, P., Huangleng, J., Su, H., Chen, P., Chen, X., Zhao, H., Kang, Z., Tang, J., Jiang, G., Li, Z., Zou,
756 S., Dong, K., Huang, Y., Li, N., 2022. Co-occurrence of chromophytic phytoplankton and the *Vibrio*
757 community during *Phaeocystis globosa* blooms in the Beibu Gulf. *Sci. Total Environ.* 805, 150303.
758 <https://doi.org/10.1016/j.scitotenv.2021.150303>
- 759 Yang, Q., Feng, Q., Zhang, B., Gao, J., Sheng, Z., Xue, Q., Zhang, X., 2021. *Marinobacter alexandrii* sp. nov.,
760 a novel yellow-pigmented and algae growth-promoting bacterium isolated from marine
761 phycosphere microbiota. *Antonie Van Leeuwenhoek Int. J. Gen. Mol. Microbiol.* 114, 709–718.
762 <https://doi.org/10.1007/s10482-021-01551-5>
763


764 **ACKNOWLEDGMENT**

765 The authors thank F. Le Roux and Y. Labruche for their gift of *Vibrio* mutants; E. Garcés, S. Turki, H.
766 Frehi, A. Bouquet and W. Medhioub for their contribution in obtaining Mediterranean phytoplankton
767 strains; K. Escoubeyrou and D. Stien for advice on HRMS/MS data; V. Diakou-Verdin and E. Jublanc
768 from the Montpellier RIO Imaging Platform for access to microscope facilities (www.mri.cnrs.fr); P. Clair
769 from the Montpellier qPHD platform and the Technoviv platform of the UPVD for access to PCR
770 facilities; N. Brunet, A. Lang, A. Payelleville, T. Milhau and M. Leroy for their technical assistance; The
771 IFREMER LER-LR laboratory for access to the Thau Lagoon; the BIO2MAR platform at the
772 Observatoire Océanologique de Banyuls for providing access to UHPLC-HRMS/MS facilities.

773 **FUNDING**

774 European VIVALDI grant 678589, French EC2CO ROSEOCOM grant and IFREMER ALGOVIR grant.


775 FIGURES, TABLES and VIDEOS



776


777 **Figure 1. Dynamics of *Alexandrium* and *Vibrio* in the environment. (A)** Location of the monitoring
778 station in the Thau Lagoon (southern France). **(B)** Mean abundance (DNA equiv.) of *Vibrio* spp. (16S) and
779 *Alexandrium* spp. (*A. pacificum* ACT03 + *A. tamarense* ATT07). *Vibrio* cells (black line with diamond
780 dot) and degraded *Alexandrium* cells (grey line with round dot) were evidence in the 0.2–0.8 μm fraction
781 (free *Vibrio* fraction) in spring and autumn 2015. Living *Alexandrium* cells (grey line with round dot) but no
782 plankton-associated *Vibrio* spp. (black line with diamond dot) were evidence in the 0.8–180 μm in spring
783 and autumn. **(C)** Result of Akaike information criterion (AICc) models tested to explain the mean value of
784 degraded *Alexandrium* cells (dead cells) in spring. **(D)** Wald test of the AICc model attributing the mean
785 value of degraded cells of *Alexandrium* in spring to free *Vibrio*.

786



787
 788 **Figure 2. Incubation of *Vibrio atlanticus* LGP32 and *Alexandrium pacificum* ACT03 in enriched**
 789 **natural seawater (ENSW). (A) *A. pacificum* ACT03 cultured alone (grey bar) and incubated with *V.***
 790 ***atlanticus* LGP32 (black bar) in ENSW. (B) *V. atlanticus* LGP32 cultured alone (grey bar) and incubated**
 791 **with *A. pacificum* ACT03 (black bar) in ENSW. (C) Images of the interaction of *V. atlanticus* LGP32-**
 792 **GFP with *A. pacificum* ACT03 taken at 8h00 and 28h00 of co-culture. *V. atlanticus* LGP32 (small green**
 793 **cells), living *A. pacificum* ACT03 cells (large red cells) and degraded *A. pacificum* ACT03 cells (large**
 794 **green cells). *P < 0.01 (analysis by pairs).**


795



796

797 **Figure 3. Role of *Vibrio atlanticus* LGP32 starvation in the interspecific interaction process.**
 798 Experiments were conducted by incubating *A. pacificum* ACT03 with *V. atlanticus* LGP32 previously
 799 grown for 12, 36, 60 and 126 h in Zobell medium. **(A)** Percentage of motile *A. pacificum* ACT03. **(B)** *A.*
 800 *pacificum* ACT03 group-attached by *V. atlanticus* LGP32 and **(C)** *A. pacificum* ACT03 lysis after 0, 15,
 801 30, 45 and 60 min of interaction. Corresponding pictures showing **(A1)** Black arrows indicate unhooked
 802 and degrade flagellum from *A. pacificum* ACT03 flagellum, **(B1)** Image of the interaction of *V. atlanticus*
 803 LGP32-GFP with *A. pacificum* ACT03 taken at 8h00 of co-culture. *V. atlanticus* LGP32 (small green
 804 cells), living *A. pacificum* ACT03 (large red cells) and dead *A. pacificum* ACT03 (large green cell). **(C1)**
 805 Black arrow 1 indicate vesicle formation on *A. pacificum* ACT03 cell and black arrow 2 indicate exploded
 806 *A. pacificum* ACT03 cell. ***P < 0.001.


807



808


809 **Figure 4. Role of quorum sensing and the vibrioferrin iron uptake pathway in the interaction**
810 **process.** (A) Effect of *V. atlanticus* LGP32 cell density on the predation process. Experiments examined
811 the impact of *V. atlanticus* LGP32 cell density on its ability to attack *A. pacificum* ACT03. *A. pacificum*
812 ACT03 cells (2.10^4 cells) were incubated with *V. atlanticus* LGP32 grown for 60 hours in Zobell medium
813 at concentrations ranging from 5×10^3 to 5×10^5 cells mL⁻¹ (represented by black bars). For comparison, *A.*
814 *pacificum* ACT03 was also incubated with *V. atlanticus* LGP32 grown for only 12 hours in Zobell
815 medium at concentrations ranging from 5×10^3 to 4×10^6 cells mL⁻¹ (represented by grey bars). The term
816 “Att” indicates cells of *A. pacificum* ACT03 that were subjected to group attacks by *V. atlanticus* LGP32.
817 (B) CqsS, luxM, luxN, luxS, and luxP quorum sensing and PvsA, PvuB and PvuA2 vibrioferrin pathway
818 genes expression in *V. atlanticus* LGP32 grown for 12, 36 and 60 h in Zobell medium. (C) Effect of *V.*
819 *atlanticus* LGP32 mutants on the group-attacked process. Experiments were conducted by incubating *A.*
820 *pacificum* ACT03 with *V. atlanticus* LGP32, *V. atlanticus* LGP32 tagged with GFP, *V. atlanticus* LGP32
821 washed with ENSW or *V. atlanticus* LGP32 mutant ΔPvuB, ΔluxM, ΔluxR and ΔluxS previously grown
822 60 h in Zobell media (control), The percentage of *A. pacificum* ACT03 group-attacked was determined
823 during the first 30 min of exposure. (D) Effect of *V. atlanticus* LGP32 cultures media composition on the
824 group-attacked process. Experiments were conducted by incubating *A. pacificum* ACT03 with *V.*
825 *atlanticus* LGP32 grown 60 h in Zobell media supplemented with H₃BO₄ or FeCl₃. The results were
826 compared with an exposure to *V. atlanticus* LGP32 grown 60 h in Zobell media. **P <0.01, ***P <0.001
827

33



828
829 **Figure 5. Schematic representation of a putative strategy developed by *Vibrio* spp. to feed on**
830 ***Alexandrium* spp. and *G. catenatum* in the environment. (A) *Vibrio* in the environment when subjected**
831 **to starvation secrete non-protein lytic compounds. (B) Some of these lytic compounds degrade the**
832 **flagella, immobilizing the alga (immobilization stage). (C) Then *Vibrio* swims and clusters around its prey**
833 **coordinating group attacks (group-attack stage). (D) Lytic compounds released by *Vibrio* where able to**
834 **concentrate around the algae cells, thereby lysing the algae (killing stage). (E) Feeding on the released**
835 **nutrients, *Vibrio* multiply and then spread in the environment. Yellow clouds: Lytic compound release by**
836 ***Vibrio*, Grey clouds: Algal nutrients released upon lysis.**
837

838



839

840

841

842

843

844


845

846

847

Figure S1. Time and dose-dependent effects of the *V. atlanticus* LGP32 culture supernatant on *A. pacificum* ACT03 motility. (A) A time dependence experiment was conducted by incubating *A. pacificum* ACT03 for 1, 5 or 20 h with 1/1000 v/v (1 μ L/mL) of culture supernatant from *V. atlanticus* LGP32 previously grown for 60 h in Zobell culture media. **(B)** A dose dependence experiment was conducted by incubating *A. pacificum* ACT03 for 1 h with 1/1000 to 1/20 v/v (1-50 μ L/mL) of culture supernatant from *V. atlanticus* previously grown for 60 h in Zobell media. The percentage of motile *A. pacificum* ACT03 was determined after 1 hours of exposure. *P < 0.05, ***P < 0.001


848



849

850 **Figure S2. *Vibrio atlanticus* LGP32 proteome analysis following nutrient stress. (A)** Example of 2D
 851 gel, the numbers in white on the gel 4-7 correspond to the number and position of the protein spots
 852 analyzed. **(B)** Proteins identified by LC-MS/MS as differentially represented in the 2D gel comparative
 853 approach following nutrient stress. ND: Not determined; ENSW (artificial seawater).
 854

855



856

857 **Figure S3. Quorum sensing and the vibrioferrin iron uptake pathway in *Vibrio*.** (A) Putative
 858 quorum sensing (QS) pathways at low and high cell density in *Vibrio* according to Lami et
 859 al.(Lami, 2019). (B) Genetic organization of the vibrioferrin utilization gene cluster on *V.*
 860 *atlanticus* LGP32 chromosome 2. The Pvu and Pvs operons are involved in the secretion and the
 861 transport of ferric vibrioferrin and biosynthesis of vibrioferrin, respectively. Arrows indicate the
 862 transcriptional directions of the genes. VSH1126, VSH1137 and VSH1129 corresponding to
 863 PvuB, PvuA2 and PvsA genes respectively.

864

865

866 **Table 1.** Ability of *Vibrio spp.* and strains to secrete algicidal compounds and attack *Alexandrium*
 867 *pacificum* LGP32 in groups. NT: not determined.

| Vibrio species | Strains | Virulence for fish or invertebrates | References | Secretion of algicidal compounds (This study) | Group attack (This study) |
|-----------------------------------|-----------------------|--|---------------------------|--|----------------------------------|
| <i>Vibrio atlanticus</i> LGP32 | WT | + | (Gay et al., 2004) | + | + |
| // | WT + pSW3654T-GFP | + | (Le Roux et al., 2007) | + | + |
| // | ΔLuxM | ND | Ifremer Institute, France | + | + |
| // | ΔLuxS | ND | // | + | + |
| // | ΔLuxR | ND | // | + | + |
| // | ΔPvuB | ND | This work | + | - |
| <i>Vibrio tasmaniensis</i> | J5-9 | + | (Lemire et al., 2015) | + | + |
| // | LMG20012 ^T | - | (Thompson et al., 2003) | + | + |
| <i>Vibrio crassostreae</i> | J2.9 | + | (Lemire et al., 2015) | + | + |
| // | J2-8 | - | // | + | + |
| <i>Vibrio fischeri</i> | ES114 | ND | (Mandel et al., 2008) | + | + |
| <i>Vibrio harveyi</i> | ATCC14126 | + | (Liu et al., 1996) | + | + |
| <i>Vibrio aestuarianus</i> | janv-32 | + | (Labreuche et al., 2010) | + | + |

868

869 **Table 2.** Ability of *Vibrio Atlanticus* LGP32 to degrade flagella, attack in group and lysed the targeted
 870 dinoflagellates *spp.* commonly found in the Mediterranean Sea. ND not determined.

| Dinoflagellates species | Strains | Toxicity for human | References | Degraded flagella (This study) | Attacked in groups (This study) | Lysed cells (This study) |
|-------------------------------|------------------------------|--------------------|--|--------------------------------|---------------------------------|--------------------------|
| <i>Alexandrium pacificum</i> | ACT03, Thau, France | + | (Laabir et al., 2011) | + | + | + |
| <i>Alexandrium catenella</i> | Bizerte, Tunisia | + | (Fertouna-Bellakhal et al., 2015) | + | + | + |
| // | F3-9F, Tarragona, Spain | ND | // | + | + | + |
| // | C10-5, Annaba, Algeria | + | (Hadjadji et al., 2020) | + | + | + |
| <i>Alexandrium tamarense</i> | ATT07, Thau, France | - | (Rolland et al., 2012) | + | + | + |
| <i>Alexandrium spp.</i> | Golf of Tunis, Tunisia | ND | Algal collection university of Montpellier, France | + | + | + |
| // | Bizerte, Tunisia | ND | // | + | + | + |
| // | Mediterranean coast, Morocco | ND | // | + | + | + |
| <i>Prorocentrum lima</i> | PLBZT14, Bizerte, Tunisia | + | (Ben-Gharbia et al., 2016) | - | - | - |
| <i>Coolia monotis</i> | CMBZT14, Bizerte, Tunisia | ND | // | - | - | - |
| <i>Vulcanodinium rugosum</i> | IFR-VRU-01, Ingril, France | + | (Abadie et al., 2015) | - | - | - |
| <i>Karenia selliformis</i> | Golf of Gabes, Tunisia | NT | Algal collection university of Montpellier, France | - | - | - |
| <i>Scripsiella trochoidea</i> | Mellah Lagoon, Algeria | - | // | - | - | - |
| <i>Gyrodinium impudicum</i> | Golf of Tunis, Tunisia | - | // | - | - | - |
| <i>Amphidium carterae</i> | SAMS, Scotland | NT | SAMS laboratory, Scotland | - | - | - |
| <i>Gymnodinium catenatum</i> | M'diq Bay, Morocco | + | (Leblad et al., 2020) | + | + | + |

871

872 **Table 3.** Oligonucleotide sequences of primers used for RNA expression analysis.

| Species | genes | Primers Sequences | T _m (°C) | Efficiency | References |
|--------------------------------------|---|--|---------------------|------------|------------------------------|
| <i>Alexandrium pacificum</i> (ACT03) | 18S – 28S rRNA ITS region | TGATATTGTGGGCAACTGTAA AACATCTGTTAGCTCACGGAA | 54 | | (Genovesi et al., 2011) |
| <i>Alexandrium tamarense</i> (ATT07) | 18S – 28S rRNA ITS region | TGGTAATTCTTCATTGATTACAATG AACATCTGTTAGCTCACGGAA | 54 | | // |
| <i>Vibrios spp.</i> | 16S | CGGTGAAATGCGTAGAGAT TTACTAGCGATTCCGAGTTC | 62 | | (Kitatsukamoto et al., 1993) |
| <i>Vibrio atlanticus</i> LGP32 | LuxN (VS_II0260) | CACCTTGCTAGTATCATCGC ATCGAGTTAGCAAGAGCAC | 60 | 1,92 | This work |
| // | LuxM (VS_II0261) | TCCACTTATCACAAACAGG ACTGTACTTCCATTTGTCTG | 60 | 1,91 | // |
| // | LuxP (VS_II0355) | AAGTTCAGGATGAACCTATC CAAAGAGATACTTTGCTGAG | 60 | 1,89 | // |
| // | LuxS (VS_2562) | ACTCTCGAGCACCTATACG GAAGGCGTACCAATCAAGC | 60 | 1,85 | // |
| // | CqsS (VS_1725) | GACATCTATTGATGTTATGC TCACCCACTTCACGTAACCTG | 60 | 1,91 | // |
| // | PvsA (VS_II0355), Vibrio ferrin biosynthesis protein | CAGAGCAAGAGCTAGAACC TCGTTGAGAACCTGACGAG | 59 | 1,91 | // |
| // | PvuB (VS_II1126), ABC transporter vibrio ferrin uptake FecB | TAGTGCAACCATGGGAATCG TAAACCGTACGTAGACGCTC | 57 | 2,01 | // |
| // | PvuA2 (VS_II1127), Vibrio ferrin receptor FecA | GGAGCTACAAGCATTTCGTTT TTCGTTCATATGGTCGCTTCG | 57 | 2,08 | // |
| // | Housekeeping gene 1, CcmC (VS_0852) | ATTGCCGCCTTTATCGGTTT CAAGCACCCACATTGGTTT | 60 | | (Vanhove et al., 2016) |
| // | Housekeeping gene 2, 6-phosphofructokinase (VS_2913) | GCCGTCACTGTGGTGACCTT TGCTTCTTGCCTTTCGCAAT | 60 | | // |

873

874 **VIDEOS**

875 **Video 1. Dynamics of *Vibrio atlanticus* LGP32-*Alexandrium pacificum* ACT03 interaction.** GFP-
876 tagged *V. atlanticus* (small green cells); living *A. pacificum* (large red cells); degraded *A. pacificum* (large
877 green cells) filmed under an epifluorescence microscope.

878

879 **Video 2. Second-by-second timing of *Vibrio Atlanticus* LGP32 attacking *Alexandrium pacificum***
880 **ACT03.** GFP-tagged *V. atlanticus* (small green cells); *A. pacificum* living cell (large red cells) filmed
881 under an epifluorescence microscope.

882

883 **Video 3. Degradation and disruption of *Alexandrium pacificum* ACT03 flagella.** Effect of *Vibrio*
884 supernatant on the first stage of the interaction filmed under a confocal microscope.

885

886 **Video 4. Group attacks of *Vibrio atlanticus* LGP32 on target *Alexandrium pacificum* ACT03.** *Vibrio*,
887 filmed under a confocal microscope, attacks in groups one immobilized *Alexandrium* cell then moves on
888 to attack — still as a group — another cell without touching the other whole cells, suggesting active
889 communication between *Vibrio* cells. *V. atlanticus* LGP32 (small cells); *A. pacificum* ACT03 (large cells).

890

891 **Video 5. Vesicle formation and bursting of an *Alexandrium pacificum* ACT03.** Direct effect of *Vibrio*
892 supernatant on *Alexandrium* after 126 h of culture filmed under a confocal microscope