

# *Electronic Supplementary Information*

## **Spatio-temporal investigation of the crosslinking and the mechanical properties in a UV-curable rubber bearing photo-dimerizable moieties**

Colanges, S.;<sup>1, 2, 3</sup> Ballu, K.;<sup>1, 2</sup> Lamouroux, A.;<sup>1, 2, 3</sup> Pecastaings, G.;<sup>2</sup> Ly, I.;<sup>2</sup> Le Coz, C.;<sup>1</sup> Tassaing, T.;<sup>4</sup> Brunet, T.;<sup>3</sup> Grau, E.;<sup>1</sup> Cramail, H.;<sup>1</sup> Mondain-Monval, O.;<sup>2,\*</sup> Vidil, T.<sup>1,\*</sup>

<sup>1</sup> University of Bordeaux – CNRS – Bordeaux INP, LCPO, 16 avenue Pey-Berland 33600 Pessac, France

<sup>2</sup> University of Bordeaux – CNRS – Centre de Recherche Paul Pascal, Pessac, France

<sup>3</sup> University of Bordeaux – CNRS – Bordeaux INP, Institut de Mécanique et d'Ingénierie, Talence, France

<sup>4</sup> University of Bordeaux – CNRS – Bordeaux INP, Institut des Sciences Moléculaires, Talence, France


### **Table of Content**

I.	<i>Characterization of the thio-cinnamate ester, TC</i>	S2
II.	<i><sup>1</sup>H NMR characterization of TC-PBut for X = 0.25</i>	S2
III.	<i>Calculation of x<sub>funct</sub>, x<sub>unreact</sub>, and x<sub>cycle</sub></i>	S3
IV.	<i>FTIR characterization of TC-PBut for X = 0.25</i>	S5
V.	<i>Size Exclusion Chromatography analysis of PBut, and TC-PBut for X = 0.25 and 0.50</i>	S5
VI.	<i>TGA analysis of PBut, and TC-PBut for X = 0.25</i>	S7
VII.	<i>DSC analysis of PBut, and TC-PBut for X = 0.25 and 0.50</i>	S7
VIII.	<i>Temperature profile in the UV oven</i>	S8
IX.	<i>DSC thermograms of the photo-crosslinked samples</i>	S8
X.	<i>DMA testing of TC-PBut-15 and TC-PBut-120: frequency sweeps</i>	S12
XI.	<i>Reduced Young's modulus maps obtained in AFM</i>	S14

## I. Characterization of the thio-cinnamate ester, TC




**Scheme S1:** Synthesis pathway for the thio-cinnamate ester, TC




**Figure S1:**  $^1\text{H}$  NMR spectrum of TC (400 MHz,  $\text{CDCl}_3$ )

## II. $^1\text{H}$ NMR characterization of TC-PBut for $X = 0.25$



**Figure S2:**  $^1\text{H}$  NMR spectrum of TC-PBut,  $X = 0.25$  (400 MHz,  $\text{CDCl}_3$ )

### III. Calculation of $x_{\text{funct}}$ , $x_{\text{unreact}}$ , and $x_{\text{cycle}}$



**Figure S3:**  $^1\text{H}$  NMR spectrum of TC-PBut,  $X = 0.5$  (400 MHz,  $\text{CDCl}_3$ )

It is possible to calculate  $x_{\text{funct}}$ ,  $x_{\text{unreact}}$ , and  $x_{\text{cycle}}$  from the integrals,  $A_1$ ,  $A_2$ ,  $A_2'$  and  $A_3$  from the  $\text{RCH}_2\text{S}-$  methylene protons (2), the  $\text{H}_2\text{C}=\text{CH}-$  alkenic protons (f), the  $-\text{HC}=\text{CH}-$  and  $\text{H}_2\text{C}=\text{CH}-$  alkenic protons (b, e), and the aliphatic protons of chemical shifts below 2.2 ppm, respectively. In terms of the indices defined in **Figure S3**:

$$S1 \left\{ \begin{array}{l} A_1 = 2p + 2q \\ A_2 = 2n \\ A_{2'} = 2m + n \\ A_3 = 4m + 3n + 5p + 4q + 7q' \\ q = q' \end{array} \right.$$

Moreover, the direct mathematical definitions of  $x_{\text{funct}}$  and  $x_{\text{unreact}}$  are given by:

$$S2 \left\{ \begin{array}{l} x_{\text{funct}} = \frac{p + q}{m + n + p + q + q'} \\ x_{\text{unreact}} = \frac{m + n}{m + n + p + q + q'} \end{array} \right.$$

From the system of equations S1, we have:

$$S1 \left\{ \begin{array}{l} p + q = A_1/2 \\ n = A_2/2 \\ m = A_{2'}/2 - A_2/4 \\ A_3 = 2A_{2'}, -A_2 + 3A_{2'}/2 + 5(p + q) + 6q \\ q = q' \end{array} \right.$$

$$\Leftrightarrow S1 \left\{ \begin{array}{l} p + q = A_1/2 \\ n = A_2/2 \\ m = A_{2'}/2 - A_2/4 \\ A_3 = 2A_{2'} + A_2/2 + 5A_1/2 + 6q \\ q = q' \end{array} \right.$$

$$\Leftrightarrow S1 \left\{ \begin{array}{l} p + q = A_1/2 \\ n = A_2/2 \\ m = A_{2'}/2 - A_2/4 \\ q = A_3/6 - A_{2'}/3 - A_2/12 - 5A_1/12 \\ q = q' \end{array} \right.$$

By reinjecting these equations in S2, it comes:

$$S2 \left\{ \begin{array}{l} x_{funct} = \frac{A_1/2}{A_{2'}/2 - A_2/4 + A_2/2 + A_1/2 + A_3/6 - A_{2'}/3 - A_2/12 - 5A_1/12} \\ x_{unreact} = \frac{A_{2'}/2 - A_2/4 + A_2/2}{A_{2'}/2 - A_2/4 + A_2/2 + A_1/2 + A_3/6 - A_{2'}/3 - A_2/12 - 5A_1/12} \end{array} \right.$$

$$\Leftrightarrow S2 \left\{ \begin{array}{l} x_{funct} = \frac{A_1/2}{A_1/12 + A_2/6 + A_{2'}/6 + A_3/6} \\ x_{unreact} = \frac{A_{2'}/2 + A_2/4}{A_1/12 + A_2/6 + A_{2'}/6 + A_3/6} \end{array} \right.$$

$$\Leftrightarrow S2 \left\{ \begin{array}{l} x_{funct} = \frac{6A_1}{A_1 + 2A_2 + 2A_{2'} + 2A_3} \\ x_{unreact} = \frac{6A_{2'}}{A_1 + 2A_2 + 2A_{2'} + 2A_3} \end{array} \right.$$

And


$$x_{cycle} = 1 - x_{funct} - x_{unreact}$$

The following Table gather the values measured for the different integrals of the NMR spectra (the peak 3 was used as a reference), and the corresponding values of  $x_{funct}$ ,  $x_{unreact}$ , and  $x_{cycle}$ , using the above equations:

$X$	$A_1$	$A_2$	$A_{2'}$	$A_3$	$x_{funct}$ (%)	$x_{unreact}$ (%)	$x_{cycle}$ (%)
0.5	1.96	1.32	1.33	14.38	33.0	33.5	33.5
0.25	1.98	5.38	4.19	24.10	17.1	59.5	23.3


**Table S1 :** Integral values measured for the NMR spectra of **TC-PBut**, X = 0.25 and X = 0.5

#### IV. FTIR characterization of TC-PBut for $X = 0.25$




**Figure S4:** FTIR spectrum of **TC-PBut**,  $X = 0.25$


#### V. Size Exclusion Chromatography analysis of PBut, and TC-PBut for $X = 0.25$ and $0.50$



**Figure S5:** SEC elugram of PBut (THF, polystyrene calibration)



**Figure S6:** SEC elugram of TC-PBut,  $X = 0.25$  (THF, polystyrene calibration)



**Figure S7:** SEC elugram of TC-PBut,  $X = 0.5$  (THF, polystyrene calibration)

## VI. TGA analysis of PBut, and TC-PBut for $X = 0.25$




Figure S8: TGA analysis of PBut and TC-PBut,  $X = 0.25$  ( $10 \text{ } ^\circ\text{C min}^{-1}$ ,  $\text{N}_2$ )

## VII. DSC analysis of PBut, and TC-PBut for $X = 0.25$ and $0.50$






Figure S9: DSC thermograms of PBut and TC-PBut,  $X = 0.5, 0.25$  ( $10 \text{ } ^\circ\text{C min}^{-1}$ )

## VIII. Temperature profile in the UV oven




**Figure S10:** Temperature profile inside the UV oven (KX-314-01, Novachem®,  $\lambda = 365$  nm,  $1000 \text{ mW cm}^{-2}$ ). The data are collected with a thermocouple, inside a sample photo-crosslinked at 14 cm from the LEDs.


## IX. DSC thermograms of the photo-crosslinked samples




**Figure S11:** DSC thermograms ( $10 \text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-10,  $X = 0.5$ , exposed to UV ( $1000 \text{ mW cm}^{-2}$ , 14 cm) for 5 min on both sides.




**Figure S12:** DSC thermograms ( $10\text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-15,  $X = 0.5$ , exposed to UV ( $1000\text{ mW cm}^{-2}$ , 14 cm) for 7.5 min on both sides.




**Figure S13:** DSC thermograms ( $10\text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-20,  $X = 0.5$ , exposed to UV ( $1000\text{ mW cm}^{-2}$ , 14 cm) for 10 min on both sides.



**Figure S14:** DSC thermograms ( $10\text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-30,  $X = 0.5$ , exposed to UV ( $1000\text{ mW cm}^{-2}$ , 14 cm) for 15 min on both sides.




**Figure S15:** DSC thermograms ( $10\text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-120,  $X = 0.5$ , exposed to UV ( $1000\text{ mW cm}^{-2}$ , 14 cm) for 60 min on both sides.




**Figure S16:** DSC thermograms ( $10 \text{ }^{\circ}\text{C min}^{-1}$ ) of TC-PBut-240,  $X = 0.5$ , exposed to UV ( $1000 \text{ mW cm}^{-2}$ , 14 cm) for 120 min on both sides.

## X. DMA testing of TC-PBut-15 and TC-PBut-120: frequency sweeps



**Figure S17:** Frequency sweep analysis of TC-PBut-15 at various temperatures ( $\varepsilon = 0.005\%$ )



**Figure S18:** Frequency sweep analysis of TC-PBut-120 at various temperatures ( $\varepsilon = 0.005\%$ )

$a_T$	Temperature (°C)
8.40E-04	40
0.0188351	30
0.780899	20
49.2714	10
3105.67	0
172149	-10
1.09E+07	-20
6.18E+08	-30
2.24E+10	-40
1.21E+11	-50

**Table S2 :** Shift factors,  $a_T$ , for the construction of the master curve of TC-PBut-15

Coefficient WLF	
c1	5.06037e15
c2	3.02189e16 K
Tref	9.89238 °C
R <sup>2</sup>	0.996135

**Table S3 :** WLF coefficient for the construction of the master curve of TC-PBut-15

$a_T$	Température (°C)
1.98E-05	50
4.97E-04	40
0.020458	30
1.28951	20
81.2802	10
3958.08	0
231955	-10
1.37E+07	-20
8.19E+08	-30
5.04E+10	-40


**Table S4 :** Shift factors,  $a_T$ , for the construction of the master curve of TC-PBut-120

Coefficient WLF	
c1	195.195
c2	1164.71 K
Tref	29.9924 °C
R <sup>2</sup>	0.999399


**Table S5 :** WLF coefficient for the construction of the master curve of TC-PBut-120

## XI. Reduced Young's modulus maps obtained in AFM


The figures below describe the evolution of the Young's modulus over a surface of approximately  $1 \mu\text{m}^2$  for each sample. The modulus value is linked to the color scale. They all suggest local inhomogeneities in cross-linking, with modulus values varying over a wide range, which account for the uncertainties in the values reported in Figure 11 of the manuscript. The values reported in Figure 11 are spatial averages taken over the entire image, performed using the Nanoscope Analysis software.




**Figure S19:** Reduced Young's modulus maps for TC-PBut-10,  $z = 100 \mu\text{m}$




**Figure S20:** Reduced Young's modulus maps for TC-PBut-20,  $z = 100 \mu\text{m}$




**Figure S21:** Reduced Young's modulus maps for TC-PBut-120,  $z = 100 \mu\text{m}$



**Figure S22:** Reduced Young's modulus maps for TC-PBut-240,  $z = 100 \mu\text{m}$



**Figure S23:** Reduced Young's modulus maps for TC-PBut-120,  $z = 1.5$  mm



**Figure S24:** Reduced Young's modulus maps for TC-PBut-240,  $z = 1.5$  mm