



Make visible the invisible: Optimised development of an environmental DNA metabarcoding tool to characterise trematode parasitic communities

Philippe Douchet, Jérôme Boissier, Stephen Mulero, Hubert Ferté, Margot Doberva, Jean-François Allienne, Eve Toulza, Kévin Bethune, Olivier Rey

► To cite this version:

Philippe Douchet, Jérôme Boissier, Stephen Mulero, Hubert Ferté, Margot Doberva, et al.. Make visible the invisible: Optimised development of an environmental DNA metabarcoding tool to characterise trematode parasitic communities. Environmental and Agronomical Genomics 2021 Symposium EAGS, Oct 2021, Tours, France. . hal-04888136

HAL Id: hal-04888136

<https://hal.science/hal-04888136v1>

Submitted on 15 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Make visible the invisible: Optimised development of an environmental DNA metabarcoding tool to characterise trematode parasitic communities

One world – One health – One ocean




Philippe Douchet, Jérôme Boissier, Stephen Mulero, Hubert Ferté, Margot Doberva, Jean François Allienne, Eve Toulza, Kévin Bethune, Olivier Rey

IHPE, Univ Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Montpellier, France

Contacts: philippe.douchet@univ-perp.fr
olivier.rey@univ-perp.fr

Trematoda → -Ubiquitous over all marine and freshwater ecosystems
-44.000 estimated species (14% described)¹
-Key organisms in ecosystem functioning²
-100 species involved in human and livestock diseases
-Important number of cryptic species³
-Difficult to tract in natural environments

→ Need to develop new tools for a better assessment and understanding of the structure and dynamics of trematode diversity



In silico primers design and evaluation of the trematode metabarcode




Figure 1. Percentage of sequences amplified by virtual PCR (a) per phylum and (b) per class of Platyhelminths when allowing up to 3 mismatches between each primer and the targeted sequences. *: Fisher's exact test p-value < 0.001




Figure 2. Partial phenetic tree illustrating the resolution of the trematode metabarcode within the Schistosomatidae family. The numbers on each branch indicate the percentage difference in nucleotides between amplicons of the different trematode species

In vitro validation of the metabarcode efficiency to detect and identify trematodes



In natura comparative analysis between a classical trematode species characterisation and our eDNA-based monitoring method




Figure 3. Pictures of the four sites sampled in this study (Occitanie region; southern France)




Figure 4. Number of trematode species characterised in field by classical sampling (in grey), by eDNA monitoring (in blue), and by both over the four sites sampled in this study

This new eDNA-based metabarcoding tool open new perspectives for fundamental and applied studies in conservation, sanitary survey, and field ecology

Références: 1- Carlson et al., Proceedings of the Royal Society B: Biological Sciences 287, n° 1939 (2020)
2- Dougherty et al., Conservation Biology 30, n° 4 (2016)
3- Poulin, Biology Letters 7, n° 2 (2011)