



## Poster Functionalized membrane domains: an ancestral feature of Archaea

Maxime Tourte, Marta Salvador-Castell, P. Schaeffer, Vincent Grossi,  
Philippe Oger

### ► To cite this version:

Maxime Tourte, Marta Salvador-Castell, P. Schaeffer, Vincent Grossi, Philippe Oger. Poster Functionalized membrane domains: an ancestral feature of Archaea. Bilayers at the ILL, Dec 2019, Grenoble, France. hal-04895218

HAL Id: hal-04895218

<https://hal.science/hal-04895218v1>

Submitted on 17 Jan 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Bilayers at the ILL

ID de Contribution: 36

Type: Poster

### Functionalized membrane domains: an ancestral feature of Archaea

mercredi 11 décembre 2019 19:45 (15 minutes)

The current tree of life is divided into three domains: Bacteria, Eukarya and Archaea. While bacterial and eukaryotic lipids are constituted of sn-3 glycerol on which straight fatty acids are ester-bound, archaeal lipids contain ramified isoprenoid hydrocarbons that are linked through ether bounds on a sn-1 glycerol. As a result, plasma membranes are spatially organized into domains of distinct functions in Bacteria and Eukarya, but membrane functionalization remains a debated question in Archaea. However, the recent reinvestigation of the lipid content of the hyperthermophilic and piezophilic archaeon *Thermococcus barophilus* led to the elaboration of a novel membrane ultrastructure in which monolayered and apolar hydrocarbon-containing bilayered domains may be delineated[1]. To estimate the ubiquity of this novel membrane ultrastructure in and out of the order Thermococcales and all the organizational possibilities it implies, we reassessed the lipid composition of all the Thermococcales type species and all archaea. We show that almost all archaea can synthesize di- and tetraether lipids, which reflect their lifestyle more than their phylogeny and support the universal existence of a functionalized membrane in Archaea. Our findings establish functionalized membrane domains as a universal feature of Archaea, and thus of all living organisms, opening new avenues to understand the membrane physiology and adaptation since the origin of cellular life.

1. Cario A, Grossi V, Schaeffer P, Oger PM. Membrane homeoviscous adaptation in the piezo-hyperthermophilic archaeon *Thermococcus barophilus*. *Front Microbiol*. 2015; 6: 1152.

**Auteur principal:** TOURTE, Maxime (Microbiology, Adaptation and Pathogeny Lab CNRS UMR5240 Université de Lyon)

**Co-auteurs:** SALVADOR CASTELL, Marta (Microbiology, Adaptation and Pathogeny Lab CNRS UMR5240 Université de Lyon); SCHAEFFER, Philippe (Institut de Chimie de Strasbourg CNRS UMR 7177 Université de Strasbourg, France); GROSSI, Vincent (Laboratoire de Géologie de Lyon, Terre, Planètes, Environnement CNRS UMR 5176 Université de Lyon, France); OGER, Philippe (Microbiology, Adaptation and Pathogeny Lab CNRS UMR5240 Université de Lyon)

**Orateur:** TOURTE, Maxime (Microbiology, Adaptation and Pathogeny Lab CNRS UMR5240 Université de Lyon)

**Classification de Session:** Wine & cheese poster session