



Open Questions and Future Directions in Titan Science

Conor Nixon, Natalie Carrasco, Christophe Sotin

► To cite this version:

Conor Nixon, Natalie Carrasco, Christophe Sotin. Open Questions and Future Directions in Titan Science. Rosaly M.C. Lopes, Charles Elachi, Ingo Mueller-Wodarg, Anezina Solomonidou. Titan after Cassini-Huygens, Elsevier, In press, 9780323991612. 10.48550/arXiv.2501.00971 . hal-04901013

HAL Id: hal-04901013

<https://hal.science/hal-04901013v1>

Submitted on 20 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Chapter 15: Open Questions and Future Directions in Titan Science

Conor A. Nixon^a, Nathalie Carrasco^{b,c}, Christophe Sotin^d

^a*Planetary Systems Laboratory, NASA Goddard Space Flight Center, Greenbelt, 20771, MD, USA*

^b*Universite Paris-Saclay, UVSQ, CNRS, LATMOS, Guyancourt, 78280, France*

^c*ENS Paris-Saclay, 4 Av. des Sciences, Gif-sur-Yvette, 91190, France*

^d*Laboratoire de Planétologie et Géodynamique, UMR 6112, CNRS, Université de Nantes, 2 rue de la Houssinière, Nantes, 44322, France*

1. Introduction: Current Sources of Knowledge

In the preceding 14 chapters, a detailed description of the current state of knowledge about Titan in the era of Cassini-Huygens has been laid out. In this chapter, we finally take a step back from the details and look at the big picture: how did we arrive at this current state of knowledge? What are the large topical areas that remain mysterious? And how can we attempt to answer these questions via future investigations?

First let us recap at the high level how our constraints on Titan have been collected. There are primarily four techniques through which knowledge has been gained:

Telescopic observations: Titan was first discovered by means of telescope by Christiaan Huygens in 1655, and until the post-WW2 space age, this remained the only direct means of gathering data about Titan. Seminal telescopic findings about Titan include the discovery of the first atmospheric gas, methane, by Gerard Kuiper in 1944 (Kuiper, 1944); the discovery of carbon monoxide (Lutz et al., 1983) which helped to resolve questions about the atmospheric bulk composition and density; the active radar measurements of surface albedo that showed a lack of global ocean (Muhleman et al., 1990); the first imaging of the surface (Smith et al., 1996); the discovery of clouds (Griffith et al., 1998). Further descriptions of telescopic observations of Titan throughout the pre-Cassini and Cassini era can be found elsewhere

in this volume: see especially Chapter 7 for details on observations of the atmosphere and Chapter 11 for surface observations.

Laboratory Work: Laboratory research has filled many gaps in knowledge left by remote sensing and the limited in-situ investigations performed to date. This especially includes the investigations of analog haze materials ('tholins') (see review by Cable et al., 2011) which has provided significant insights into the atmospheric chemistry, the nature of haze materials, and the potential for alteration of these grains especially in hydrolysis reactions that can lead to amino acids (Khare et al., 1986; Neish et al., 2009). Moreover, laboratory gas and ice spectroscopy provided important insights to decipher the observations made during the Cassini space mission (Sung et al., 2013; Sung et al., 2018, 2020; Bernath and Fernando, 2021; Bernath et al., 2023; Anderson et al., 2018). Reaction rates and pathway measurements (kinetics) have informed our knowledge of chemical processes taking place in Titan's atmosphere (Balucani et al., 2010; Kaiser et al., 2015; Dutuit et al., 2013; Gans et al., 2013; Carrasco et al., 2022). Clathrate and co-crystal formation investigations (Cable et al., 2021b), and aeolian, pluvial and fluvial erosion experiments (e.g. Litwin et al., 2012; Burr et al., 2015a,b; Yu et al., 2017b,a, 2018, 2020; Burr et al., 2020; Maue et al., 2022; MacKenzie et al., 2023; Hirai et al., 2023) deeply improved our understanding of the observed features at Titan's surface.

Modeling: Some problems must be tackled computationally, notably dynamical models of solar system evolution and accretion scenarios that provide insights into Titan's origins in the context of the Saturn system (Mousis et al., 2002a,b, and Chapter 4). However, models specializing in particular areas of Titan science have proved immensely valuable at explaining observations and posing further questions. These include: upper atmosphere sputtering and escape models (Lammer and Bauer, 1993; Lammer et al., 1998; Mandt et al., 2012; Johnson, 2009; Gu et al., 2019; Shematovich et al., 2003; Snowden and Higgins, 2021; Johnson, 1994; Krasnopolsky, 2016; Tseng et al., 2008, and Chapter 6); bulk atmosphere climate and dynamical models including GCMs (Global Circulation Models) (Del Genio et al., 1993; Houdin et al., 1995; Mitchell, 2008; Friedson et al., 2009; Lebonnois et al., 2012; Tokano, 2013; Lora et al., 2015, 2019; Chatain et al., 2022, and Chapter 8); mesoscale and regional models (Tokano, 2009; Charnay et al., 2014a; Mitchell et al., 2011; Rafkin and Soto, 2020; Spiga et al., 2020; Chatain et al., 2022, and Chapter 8); photochemical models (Yung et al., 1984; Wilson and Atreya, 2003; Lav-

vas et al., 2008a,b; Krasnopolsky, 2009, 2010; Vuitton et al., 2019; Loison et al., 2015; Dobrijevic et al., 2014, and Chapter 7); microphysical models of cloud droplet and haze formation (Barth and Toon, 2004, 2006; Burgalat et al., 2014; Larson et al., 2015, and Chapters 7 and 8); surface models including erosion, dune formation, river branching and more (Tokano, 2008; Barnes et al., 2015; Charnay et al., 2014a; Burr et al., 2013a; Birch et al., 2023, and Chapters 9–11); crustal tectonic models (Mitri et al., 2008; Liu et al., 2016b; Burkhard et al., 2022; Schurmeier et al., 2023, and Chapter 13); impact models (Griffith and Zahnle, 1995; Artemieva and Lunine, 2003; Kress and McKay, 2004; Korycansky and Zahnle, 2011; Zahnle et al., 2014; Crósta et al., 2021, and Chapter 3); interior structure models (Grasset et al., 2000; Tobie et al., 2006; Mitri and Showman, 2008; Tobie et al., 2009; Sohl et al., 2014; Journaux et al., 2020) and oceanographic models (Tyler, 2008, 2014; Soderlund, 2019; Vincent et al., 2022; Soderlund et al., 2023, and Chapter 12).

Spacecraft Missions: The missions that have reached the Saturn system include Pioneer 11 (1979), Voyager 1 (1980) and Voyager 2 (1981); and Cassini-Huygens (2004–2017) (Kohlhase and Penzo, 1977; Matson, 2002; Lebreton, 2002). These missions have built on each other and each of the generations of spacecraft has successively provided much greater detail than the previous. Of these, only Voyager 1 and Cassini-Huygens have been actively targeted to visit Titan, and while Voyager 1 provided an exciting first look at Titan, sufficient to stimulate excitement for a follow-on mission, it was Cassini-Huygens that first revealed Titan in detail. The results from Cassini-Huygens have been extensively described in this book (see Chapter 2 for mission overview).

At the end of this chapter we will return to explore how these four techniques may be used to answer the large, high-level open questions in Titan science. First, we will describe the principal open questions themselves.

2. Open Questions

We have attempted to distill the very large number of possible future inquiries of Titan into a relatively concise list of twenty high level questions,¹

¹Inspired by David Hilbert’s 23 unsolved mathematical problems of 1900 (republished in Hilbert, 2000).

each of which would necessarily entail a multitude of more specific investigations and studies. While this list may not encompass all possible open questions, and is divided into topics according to our preference and not in any way uniquely, we believe that it does however span a wide range of the most intriguing topics about Titan, and may form some sort of guide especially for those embarking into Titan studies for the first time. See Fig. 1 and Table 1.




Figure 1: Open questions in Titan science.

Table 1: Key Open Questions in Titan Science

Area	Question Statement	Current Evidence	Data Needed
1 <i>Formation</i>	What were the building blocks of Titan?	Isotopic ratios; elemental ratios	Isotopic ratios; crustal clathrates; noble gases
2 <i>Formation</i>	How does Titan fit into the dynamical history of the Saturn system?	Size; density; orbital elements	Dynamical modeling
3 <i>Formation</i>	When and how did Titan's atmosphere form?	Isotopic ratios; elemental ratios	Isotopic ratios; crustal clathrates; noble gases
4 <i>Interior</i>	What is the internal state of differentiation including the core?	Moment of inertia (MOI)	Higher order MOI
5 <i>Interior</i>	Is Titan's deep water ocean habitable?	Modeling	Gravity; MOI; sub-surface radar
6 <i>Interior</i>	Does the crust contain significant amounts of clathrates?	Modeling; NIR spectroscopy	Gravity; MOI; sub-surface radar
7 <i>Interior/Surface</i>	Is there active geotectonic activity on Titan at the present day?	Surface imaging and NIR spectroscopy	Imaging at higher resolution; sub-surface radar
8 <i>Surface</i>	Are Titan's lakes and seas connected through the sub-surface?	Lake shapes and morphologies; depths	Improved bathymetry; GPR
9 <i>Surface</i>	What is the composition of Titan's seas and lakes?	Radar and NIR data	Improved remote sensing data; in situ sampling
10 <i>Surface</i>	How are molecules produced in the atmosphere altered on the surface?	Modeling; NIR spectroscopy	Surface sampling; laboratory investigations
11 <i>Surface</i>	What is the origin and composition of Titan's dunes?	Modeling	Surface sampling; modeling
12 <i>Surface</i>	When were the river channels carved and the low latitude seas filled?	Imaging/radar data; modeling	Global topographic mapping; surface sampling
13 <i>Surface</i>	Is there exposed water (or other) ice on the surface?	Imaging & spectroscopy	In situ sampling; improved remote sensing
14 <i>Atmosphere</i>	What types of clouds exist in the troposphere and stratosphere?	Imaging/spectroscopic data; modeling	Improved spectral imaging; micophysical modeling
15 <i>Atmosphere</i>	How often does it rain on Titan?	Imaging/spectroscopic data; modeling	Surface/over atmosphere in situ measurements
16 <i>Atmosphere</i>	What is the availability of P and S compounds for chemistry?	Spectroscopic and MS data	Improved remote sensing and in situ measurements
17 <i>Atmosphere</i>	What is the composition of Titan's aerosols and how do they evolve?	Lab experiments; modeling	Modeling and in situ experiments
18 <i>Atmosphere</i>	What is the cause of the tilted atmospheric rotation?	Modeling	Modeling and IR atmospheric mapping
19 <i>Atmosphere</i>	Are important prebiological molecules produced in the atmosphere?	Spectroscopic and MS data	Spectroscopic and in situ measurements
20 <i>Atmosphere</i>	Is Titan's atmosphere stable over time?	MS measurements; modeling; interior MOI	MS measurements; modeling; interior MOI

Abbreviations: MOI = moment of inertia. MS = mass spectroscopy. NIR = near infrared. GPR = Ground penetrating radar.

2.1. Questions about Titan's formation (1-3)

1) What were the building blocks of Titan?

Our current sparse constraints regarding Titan's interior permit multiple interpretations of its core composition. The moment of inertia ($C/MR^2 \sim 0.34$) from gravity measurements by Cassini (Iess et al., 2010; Iess et al., 2012; Durante et al., 2019) can be modeled in several ways, including a bimodal model with a mixture of low density and high-density components (Néri et al., 2020), or a more uniform composition lacking in significant amounts of iron (Vance et al., 2018).

We must also explain the presence and bulk composition of the atmosphere, mostly made of nitrogen and methane. The current favored paradigm for the origin of nitrogen is photolysis of ammonia (Mandt et al., 2014), while the presence of methane is still a subject of intense debate (see Q3), with multiple possibilities including trapping as primordial clathrate (Mousis et al., 2015; Tobie et al., 2006) and generation in the interior (Glein, 2015; Miller et al., 2019).

Other atmospheric quantities can in principle provide constraint on building blocks, such as the ratios of stable isotopes (D/H, $^{12}\text{C}/^{13}\text{C}$, $^{14}\text{N}/^{15}\text{N}$, $^{16}\text{O}/^{18}\text{O}$, etc) (Pinto et al., 1986; Wong et al., 2002; Jennings et al., 2008; Mandt et al., 2009; Nixon et al., 2012; Mandt et al., 2012), stable noble gas abundances (^{20}Ne and ^{22}Ne , ^{36}Ar and ^{38}Ar) and the abundances of radiolytic elements (e.g. ^{40}Ar) (Glein, 2015; Miller et al., 2019), as described in Chapter 3. If unaltered from formation, or with adequate knowledge of alteration processes, the present-day ratios and abundances constrain the values at time of formation through comparison to ratios in other primitive materials such as comets and meteorites (Mandt et al., 2014; Glein, 2015; Miller et al., 2019).

However, making inferences about building blocks from present-day isotopic ratios in the atmosphere is problematic, since it requires assumptions about how the atmosphere has evolved (Owen and Niemann, 2009; Nixon et al., 2012; Mandt et al., 2012; Wong et al., 2015). Similarly, making interpretations from noble gas abundances is reliant on assumptions about the amount of internal differentiation and outgassing.

More robust isotopic ratios could be derived from measurements of crustal material, with the D/H in water - likely the main component of the crust - being particularly informative (Coustenis et al., 2009). Measurements from Dragonfly DraMS may be helpful here (Barnes et al., 2021; Grubisic et al., 2021). In addition, better measurements of the gravity field will yield more

precise values for the higher degrees of MoI ($J_2\dots$) and the Love numbers ($h_2, k_2\dots$) which can be used to make inferences about the distribution of mass and the rigidity of the ice shell (Iess et al., 2010; Iess et al., 2012).

With such future measurements, we will be better positioned to find answers to our questions about the primordial building blocks of Titan.

2) How does Titan fit into the dynamical history of the Saturn system?

The Saturn system appears less regular than the Jovian system with its four large satellites and multiple smaller, apparently captured bodies. In contrast, Saturn's moon system is dominated by the giant Titan, with the other mid-sized moons much less massive in comparison, adding in total to less than 5% of Titan's mass.

Recent numerical work has pointed to a potentially young age for the Saturnian moons interior to Titan (i.e. closer to Saturn) (Ćuk et al., 2016; Lainey et al., 2017; Wisdom et al., 2022; Teodoro et al., 2023), although this has been disputed based on surface age estimates (Wong et al., 2023). In addition, the dynamical ages depend on the initial value of Saturn's tidal quality factor, which is a poorly known parameter. Neveu and Rhoden (2019) argue for a large initial value for this parameter in order to explain the observations of each moon, which leads to a formation age similar to that of Saturn, with the exception of Mimas, which may have formed later.

Also, Saturn possesses the only icy ring system, which appears unstable over time (Kempf et al., 2023) and may have been formed relatively late by disruption of an inner moon (Dubinski, 2019). These strands of evidence point to possible evolution of the Saturn system at recent times (Crida et al., 2019), and the role of Titan in this drama remains unclear (see Chapter 4).

Factual evidence to constrain models comes from the masses and orbital parameters of the satellites, the mass and composition of the rings, and the age of the moons as implied from surface crater counts. Improvements to our models can come from more sophisticated modeling, comparison to other satellite systems (including exoplanet moon systems) and especially better understanding of cratering histories and rates in the outer solar system, which remain significantly uncertain at the present time (Neish and Lorenz, 2012; Nesvorný et al., 2023).

3) When and how did Titan's atmosphere form?

The Voyager 1 flyby of Titan in 1980 provided the first accurate measurements of the atmospheric composition, confirming suspicions that N₂

was the dominant gas, and finding CH₄ to be the next most abundant constituent. Subsequent photochemical modeling (Strobel, 1982; Yung et al., 1984) revealed a surprising corollary: the current depletion rate of methane means that the present atmospheric inventory of CH₄ will be entirely lost in a timescale of a few 10's of Myr: much less than the age of the solar system. The age may be prolonged if methane is being actively resupplied to the atmosphere, such as by clathrate displacement (Chapter 3 Choukroun and Sotin, 2012).

Therefore, either Titan's methane is young and recent, or else we are observing the extinction of its last remnants at the end of a long history. However, the lack of sufficient hydrocarbon products on the surface makes the latter hypothesis difficult to sustain (Lorenz et al., 2008; Wilson and Atreya, 2009). As an alternative, it has been suggested that the atmospheric methane may have originated in a relatively recent event of crustal destabilization (Tobie et al., 2006).

Attempts have been made to estimate the age of methane's presence in the atmosphere through isotopic evolution by Raleigh distillation (Chapter 3 Mandt et al., 2012; Nixon et al., 2012). However, the significant error bars on present-day methane isotopic ratios, combined with uncertainties about the original D/H and ¹²C/¹³C ratios make it difficult to place quantitative values on the methane age. Another possible means of calculating the atmospheric age is to add up the surface hydrocarbon deposits and compute how long these would take to accumulate from atmospheric photochemistry, yielding upper bounds of 630 Myr (Sotin et al., 2012) and 730 Myr (Rodriguez et al., 2011).

More precise values for the present day ¹²C/¹³C, D/H and other isotopic ratios are required, and in addition some improved knowledge of their initial values, perhaps through crustal measurements, could be used to better infer the atmospheric age. Another route is a better knowledge of the amounts of hydrocarbons resulting, including both on the surface and any sequestered in the subsurface. Substantial progress on the surface hydrocarbon inventory has been made (Lorenz et al., 2008; Sotin et al., 2012; Rodriguez et al., 2014), but knowledge of the sub-surface inventory remains speculative. Future ice-penetrating radar and/or seismic measurements may help to determine the porosity and density of the crust (Mitri et al., 2021; Rodriguez et al., 2022), providing better constraint on its composition.

2.2. Questions about Titan's interior (4-6)

4) What is the internal state of differentiation including the core?

Strong evidence for differentiation within Titan comes from gravity measurements, which provide values for the Moment of Inertia (MoI) and the degree 2 tidal Love number. The MoI suggests that Titan is differentiated into a hydrosphere and a refractory core. However, Titan does not appear to have a iron-rich core like Ganymede, whose MoI value suggests a higher degree of differentiation.

Analyses of gravity data during and shortly after the mission resulted in estimated values for Titan's tidal Love number k_2 of ~ 0.6 (Iess et al., 2010; Iess et al., 2012; Durante et al., 2019), although with a large error range. Such a high value of k_2 implied not only the presence of an ocean that decouples the icy crust from the solid interior (Mitri et al., 2014b), but moreover that the ocean is much denser than pure water.

A more recent reanalysis of the gravity data (Goossens et al., 2024), including a more elaborate treatment of tidal effects, has challenged these earlier findings with a much lower estimate for k_2 of 0.375 ± 0.06 . This lower value implies a much lower ocean density of $1091 \pm 107 \text{ kg m}^{-3}$, compared to the previous estimate of 1300 kg m^{-3} . A major question, which also affects the habitability of the ocean, concerns the presence and properties of a high pressure water ice layer (or layers) at the base of the ocean (Journaux et al., 2017, 2020; Kalousová and Sotin, 2020a), affecting the exchange of core materials with the ocean (Miller et al., 2019). See also Q5.

Future measurements that can better elucidate the interior layering include further gravity data, measurements of the surface deformation during the orbit around Saturn, and seismic observations. Note that we cannot try to measure an induced magnetic field at Titan to learn about its ocean, as we can for the Galilean moons of Jupiter. This is because Saturn's magnetic field is perfectly aligned with its spin axis, which prevents the formation of an induced field that would be created by the presence of a conductive fluid (salty ocean) in a time-varying magnetic field.

5) Is Titan's deep water ocean sterile?

That Titan is an ocean world - in the sense of having a deep interior water ocean - is evidenced by measurements of the degree 2 tidal Love number k_2 (Iess et al., 2012; Durante et al., 2019; Goossens et al., 2024). However, what is much less certain is the geochemical state of the ocean and whether it

contains sufficient organics and minerals to permit life to exist (see review by Journaux et al., 2020, and also Chapter 14 of this book).

Several recent papers have examined the possible interaction of the ocean with the surface and the core. Kalousová and Sotin (2020b) show that the presence of clathrates (Q6) will thin the crustal ice shell, allowing greater chance of convection. The analysis of Fortes et al. (2007) suggests that ammonium sulfate released from the core will likewise increase the chances of crustal ice convection.

Work by Miller et al. (2019) suggests that the core of Titan is warm and rich in refractory organics, which decompose to release gases that percolate through the high pressure (HP) ice layer to reach the ocean and eventually to the atmosphere. Journaux et al. (2017) considered the effect of salt dissolution on the mobility of the high-pressure ice layer, showing that convection is possible but not certain. More recently Kalousová and Sotin (2020a) further modeled convection within the HP ice layer, concluding that convection is likely.

It is important that one or both of these interactions (core-ocean, and surface-ocean) is viable, to permit significant nutrients to enter the ocean to allow the sustenance of life (Fortes, 2000). A pessimistic estimate of the surface contribution of organics to the ocean was recently obtained by Neish et al. (2024), although many uncertainties remain.

Improved constraint on the ocean-core and ocean-crustal boundaries may be achieved by more accurate measurements of noble gases in the atmosphere, as well as interior measurements (gravity field) and modeling of the thermochemical evolution of Titan's refractory core, including use of improved data on high-pressure water ice phases.

6) Does the crust contain significant amounts of clathrates?

Clathrates - minerals wherein a crystalline cage of one substance traps an atom or molecule of a different substance - are well known on Earth and are suspected to play an important role in the formation and volatile history of solar system bodies (Mousis et al., 2015). Methane and ethane clathrates are stable at Titan's surface conditions (Lewis, 1971). Their presence and prevalence in Titan's crust, especially methane clathrate hydrates, has been a key puzzle piece required for our understanding of Titan's interior evolution (Lunine and Stevenson, 1987; Tobie et al., 2006). Clathrates have been proposed to play a dizzying array of roles, including:

- **Methane outgassing:** Several models have proposed that methane - originally trapped in clathrates - may be later released to replenish the atmospheric methane inventory (Lunine and Stevenson, 1987), ameliorating the problem of CH₄ depletion in short geological timescales. Any injection of ammonia will accelerate the destabilization process (Choukroun et al., 2010), or alternatively methane may be displaced by ethane produced in the atmosphere by photochemistry (Choukroun and Sotin, 2012).
- **Trapping of noble gases:** Studies suggest that noble gases such as Kr and Xe may be efficiently sequestered in surface clathrates, explaining the lack of detection of these gases in the atmosphere (Thomas et al., 2007, 2008; Mousis et al., 2011). However, photoionized xenon is also suggested to be more efficiently trapped into the organic haze at high altitudes than other noble gases (Hébrard and Marty, 2014).
- **Crustal thinning:** The insulating effect of a methane clathrate layer at the top of the icy crust has been proposed to lead to a thinner ice shell that is more likely to allow surface-ocean interaction and exchange (Kalousová and Sotin, 2020b)
- **Sequestration of lake materials:** The relatively rapid formation of clathrates, even under cryogenic conditions, implies that significant sequestration of liquids such as ethane likely occurs in the beds of Titan's seas and lakes (Vu et al., 2020a).

The presence or absence of clathrates has a significant impact on the structure of Titan's interior layering (Lunine and Stevenson, 1987; Choukroun et al., 2010; Mousis et al., 2014; Davies et al., 2016; Marusiak et al., 2022, and Chapters 10–14), with implications for its overall history.

For these reasons (and others), determining the prevalence, distribution and types of clathrate present in the crust is a subject of intense importance for understanding Titan. Seismic measurements by Dragonfly may provide a means of direct measurement, since methane clathrate in the crust will attenuate surface waves to a greater degree than pure water ice (Marusiak et al., 2022). Radar backscatter measurements are also sensitive to the dielectric properties of the surface material, and can be effectively used to constrain composition (Thompson and Squyres, 1990; Lorenz, 1998; Lorenz et al., 2003; Wye et al., 2007; Paillou et al., 2008; Le Gall et al., 2016).

7) Is there active geotectonic activity on Titan at the present day?

Ongoing tectonism in Titan's crust, with surface manifestations such as 'cryovolcanoes', was postulated in the era following the Voyager 1 encounter (Kargel, 1991, 1992; Lorenz, 1995), primarily based on the need to explain the long-term presence of methane in the atmosphere (see Q3, Q20). However, the higher density of liquid water relative to water ice makes cryovolcanic outflows very unlikely unless overpressure is present, for example due to the global cooling of the icy moon in the absence of a high-pressure ice layer.

A surface feature seen by VIMS early in the Cassini mission was initially hypothesized to be a signature of lobate flows (Sotin et al., 2005). However, later observations with RADAR showed that the morphology of the feature was quite bland and indistinguishable from the surroundings at centimeter wavelengths (Hayes et al., 2008).

The best candidate cryovolcanic region at the end of the mission is Sotra Patera, which consists of a prominent depression (possible cryovolcanic caldera) called Sotra Patera, a high mountain Doom Mons (Fig. 2(a)), and a lobate flow, Mohini Fluctus (Lopes et al., 2013; Lopes, 2017; Lopes et al., 2019; Solomonidou et al., 2016; Wood and Radebaugh, 2020). However, the low-resolution of the RADAR topography makes its exact nature uncertain, and no geophysical model has been proposed to explain how it may have arisen. A review of the evidence for cryovolcanic surface features is given in Lopes et al. (2013).

Cryovolcanoes aside, evidence for other types of tectonism is stronger. Cassini SAR imagery has many topographical prominences interpreted as mountain chains, contractional ridges, and similar (Barnes et al., 2007; Radebaugh et al., 2007; Mitri et al., 2010; Lopes et al., 2010; Jaumann et al., 2010; Cook-Hallett et al., 2015; Liu et al., 2016a; Lalich et al., 2022). At the southwestern boundary of the mysterious Xanadu region, evidence for several types of faulting (normal, thrust and strike-slip faults) has been proposed (Matteoni et al., 2020). Reviews of the evidence for various types of tectonism can be found in the literature (e.g. Lopes et al., 2007; Jaumann et al., 2010; Lopes et al., 2019), as well as in Chapters 9 and 13 of this volume.

However, the fact that these features mostly appear heavily eroded and there is no evidence of present-day changes makes the timing of their origin uncertain and leaves open the question of whether present day crustal activity is occurring (Solomonidou et al., 2013). These ambiguities have led some authors to pronounce Titan geophysically dead (Moore and Pappalardo, 2011). However, in view of the low spatial resolution and incomplete nature of the

present surface mapping of Titan, reaching a definitive conclusion seems premature.

The principal impediment to gathering the needed high-quality imagery and spectroscopy of the surface, and searching for temporal changes in surface morphology is the presence of the thick atmosphere, which causes several difficulties. First, its haze absorbs and scatters at visible and many near-IR wavelengths, making it difficult to image at shorter wavelengths. In the mid-infrared, emission is dominated by the warm atmosphere, while at longer RADAR wavelengths the problem becomes simply the decreased spatial resolution. Moreover, the atmosphere prevents close flybys at altitudes less than \sim 1000 km which would allow closer examination.

While Dragonfly will provide a more detailed picture of a small region of Titan's surface, mostly covered in dunes, it will not provide a high quality global map that appears to be required as the next step to unraveling the global geophysics of Titan's crust. A mapping polar orbiter equipped with multi-spectral passive and active surface imaging capability, as well as the ability to conduct sub-surface sounding (e.g. ice-penetrating radar, IPR) would allow us to much more clearly discern whether Titan is an active world at the present day, or not. A long-duration balloon mission could provide regional coverage at high resolution (Coustenis et al., 2009; Tobie et al., 2014).

2.3. Questions about Titan's surface (8-13)

8) Are Titan's lakes and seas connected through the sub-surface?

The general hydrology of Titan has been described previously in this book (see Chapter 10), so we focus here on the particular question of whether there is subsurface connection between the exposed liquid tracts.

Altimetry measurements by Cassini's RADAR indicate that the surfaces of the major seas (Kraken, Ligeia and Punga) fall on a uniform equipotential surface - interpreted as evidence that they could be connected at a subsurface level (Hayes et al., 2017). The same work found that apparently dried up lakebeds within the same drainage basins are typically found at higher elevations, adding further weight of evidence for a consistent methane 'water table'. However, it appears that Ligeia and Kraken are already connected by a network of surface channels that could help to homogenize their levels (Lorenz, 2014), and the other apparent similarities in sea surface levels may be coincidental.




Figure 2: Views of Titan’s surface. (a) Cassini topographic reconstruction of Doom Mons from Cassini radar SARtopo data - NASA image PIA13695; (b) Cassini colorized radar map of Titan’s northern hemisphere seas - NASA image PIA17655; (c) Cassini radar image of Titan dunes in the Belet ‘sand sea’ - NASA image PIA15225; (d) Huygens DISR descent image of low-latitude fluvial networks - NASA image PIA07236. Image credits: NASA/JPL-Caltech/ESA/ASI/USGS/University of Arizona.

Understanding the distribution of liquids on Titan's surface and in the subsurface has important ramifications not just for regional lacustrine studies, but for understanding of the overall climate of Titan, and its seasonal and secular trends (Tokano, 2019). Hydraulic connectivity of Titan's lakes and seas affects our understanding of their distribution, and the creation of surface morphological features through rainfall such as playas, arroyos, labyrinths and karsts (Burr et al., 2013b; Perron et al., 2006; Malaska et al., 2020; Solomonidou et al., 2020).

On the global scale, the ability of the atmospheric methane to be buffered by and replenished from the subsurface is one of the most important open topics influencing our understanding of the long-term stability of the atmosphere (Q20). Methane may be released from clathrate in the crust and seep into the methane 'water table' and eventually to the surface, providing a resupply to the atmosphere to counter methane loss (Choukroun et al., 2010; Choukroun and Sotin, 2012; Mousis et al., 2014; Davies et al., 2016).

The subsurface hydraulic connectivity can be investigated in future by improved radar or lidar altimetry over the lakes and seas (Lorenz and Waite, 2008), and especially monitored over seasonal timescales to search for any predicted changes (Mitri et al., 2007; Tokano, 2009, 2019). Measurements of the subsurface by long-wavelength radar and seismic probing may also be productive. Finally, a direct measurement of the sea surface height and seabed topography can be made by a floating probe or submarine (Stofan et al., 2013; Oleson et al., 2015)

9) What is the composition of Titan seas and lakes?

The distribution of contemporary polar seas (Fig. 2b) is not uniform, with only one significant liquid body being seen in the southern hemisphere at the present day (Ontario Lacus) (Turtle et al., 2009, and Chapter 10). In addition, Ontario's composition appears to be substantially different from the large northern seas, being apparently composed mostly of ethane (Brown et al., 2008), significantly less volatile than the methane-rich northern seas (Mastrogiuseppe et al., 2019). Lorenz (2014) has further suggested than the increased rainfall at more northerly Ligeia leads to a more methane rich composition ($\geq 80\%$) than Kraken ($\sim 60\%$).

Theoretical work by Cordier and others (Cordier et al., 2009, 2012; Tan et al., 2013; Malaska and Hodyss, 2014; Cordier et al., 2021) has sought to quantify the expected composition of the lakes including both major and minor constituents, based on predictions of condensation rates of organics

from the atmosphere.

The lake/sea composition is important from a habitability standpoint, since it has been hypothesized that prebiotically relevant organic structures can form in the lakes (Stevenson et al., 2015; Rahm et al., 2016) from molecules such as HCN and acrylonitrile, which are known to exist on Titan (Palmer et al., 2017; Thelen et al., 2019).

Our knowledge of the bulk composition of the seas may be refined in future by improved spectroscopy in the near-IR and radar loss tangent measurements at multiple wavelengths, along with continuing developments in lab experiments and theory. However, to truly determine the fractions of minor constituents dissolved in the lakes will ultimately require direct measurements through a probe sent to sample the lakes (e.g. Stofan et al., 2013; Rodriguez et al., 2022).

10) How are molecules produced in Titan's atmosphere altered on the surface?

Titan's atmosphere produces a wide variety of organic molecules (Yung et al., 1984; Waite et al., 2005, 2007; Vuitton et al., 2019) which condense in the lower stratosphere and accumulate on the surface. Simulations in the laboratory have shown that these complex organics include many molecules known to be fundamental ingredients for biology on Earth (see review by Cable et al., 2011).

An important limitation to the variety of chemicals produced in the atmosphere was thought to be the lack of oxygen involvement in the photochemistry, however lab experiments have shown that carbon monoxide can take part in reactions (Hörst et al., 2012), leading to amino acids.

Moreover, molecules produced in the atmosphere may also be altered once on the surface, e.g. by hydrolysis during episodic melting induced by impacts (Neish et al., 2008, 2009, 2010; Poch et al., 2012; Cleaves II et al., 2014; Crósta et al., 2021) - see also Chapter 13. Laboratory simulations have shown that hydrolysis of pure $H_xC_yN_z$ substances also leads to amino acids (Khare et al., 1986).

Some evidence for surface alteration exists from the detection of benzene on Titan's surface through near-IR spectroscopy with Cassini VIMS (Clark et al., 2010). This was a surprising result since the same study found no evidence of hydrocarbon ices that are expected to be more abundant, such as acetylene. A possible explanation was offered that acetylene may be modified (cyclized) into benzene under certain conditions, especially in the presence

of catalysts, for example during an impact.

Besides small molecules of astrobiological relevance, it is possible that the vast equatorial dunefields of Titan consist of atmospheric residues, perhaps reprocessed on the surface as suggested by Barnes et al. (2008) (see next section).

The Dragonfly mission, targeted to land on Titan's low-latitude dunes, is ideally suited to investigate the surface composition, and the relation between atmospheric condensates and surface molecules and particulates. Additional laboratory work will be important to verify and interpret the Dragonfly data.

11) What is the origin and composition of Titan's dunes?

Titan's equatorial dune fields (or 'sand seas', Fig 2c) were discovered during the earliest parts of the Cassini mission by RADAR imaging (Lorenz et al., 2006) and subsequently investigated in detail. These are known to cover $\sim 17\%$ of the planet, mostly at latitudes between $\pm 30^\circ$, with a combined mass of $\sim 230,000$ GT (Rodriguez et al., 2014; Lopes et al., 2020).

Multiple hypotheses exist for the formation of dune material (Barnes et al., 2015), including the 'top down' processes of sintering and flocculation, and the 'bottom up' processes of lithification and evaporite formation. However which process or processes dominate the particle formation remains as yet unknown. Once created, the particles form the large-scale, mostly longitudinal dune structures due to prevailing surface winds (Tokano, 2008).

Key questions related to the dunes include: are they formed through 'top down' or 'bottom up processes'? How long did they take to reach their present configuration, and is this stable over time? (Rubin and Hesp, 2009; Savage et al., 2014; Lucas et al., 2014; Ewing et al., 2015; Malaska et al., 2016). Modeling (Tokano, 2008) has indicated that the dunes require eastward flow to develop, however to attain atmosphere superrotation westward flow seems required. This poses a conundrum that might be resolved by a seasonal reversal of winds (Tokano, 2008). However, further investigation is warranted.

Again, Dragonfly will provide significant answers as to the dune properties, while improved remote sensing is required to search for time variations. Further laboratory work will also give comparative clues on the processes impacting the grains that comprise Titan's dunes.

12) When were the river channels carved and the low latitude seas filled?

When the Huygens spacecraft descended to Titan's surface, it was finally

revealed at the smallest scales (Tomasko et al., 2005), showing descent images of apparently empty river drainage networks leading to a dried-up seabed (Soderblom et al., 2007; Langhans et al., 2012) (Fig 2d). A consensus is now emerging that Titan appears to be drying out over time, with formerly extensive low-latitude oceans of methane retreating polewards (Larsson and McKay, 2013). However it is important to know the timescale in which this has occurred, since it has important implications for our understanding of Titan’s past history, and the unique nature of its atmosphere.

Given present-day rates of methane loss coupled with Titan’s known topography, the equatorial seas are predicted to have been connected to polar seas 300 Myr ago, and a global ocean to have existed 600 Myr ago (Larsson and McKay, 2013). However, studies of river erosive timescales to form present topography are much less constraining, with estimates ranging from 60 Kyr to 470 Myr (Cartwright et al., 2011). In addition, material accumulated in alluvial fans may be dominated by infrequent, severe storms that quickly empty again, rather than more consistent river flow (Faulk et al., 2017), complicating the estimates of fluvial network creation timescales.

To more fully understand the past history of liquids on Titan’s surface, we must increase our knowledge in several areas. Firstly, an accurate and complete high-resolution topographic map is required, since topography is a major controlling factor of lake/sea location (Lora et al., 2022) and fluvial network morphology (Miller et al., 2021). Temporal monitoring of shorelines is also necessary, to understand when and how changes are taking place at the present day, as seems to be the case at Ontario Lacus (Turtle et al., 2011c). Also, the atmosphere cannot be neglected here. In particular, having better measurements of the global methane humidity (Lellouch et al., 2014; Ádámkovics et al., 2007, 2016) and its changes on seasonal timescales is important, to better constrain climate and meteorological models (e.g. Friedson et al., 2009; Lebonnois et al., 2012; Mitchell and Lora, 2016; Lora et al., 2019).

13) Is there exposed water (or other) ice on the surface?

Detection of exposed ices on Titan’s surface would be an important finding (see also Chapter 11). Since much of the surface is covered by organics (e.g. Lopes et al., 2020) that apparently accumulate from the atmosphere (Clark et al., 2010; Rodriguez et al., 2014; Yu et al., 2017b, 2018), any exposed ices could be indicative of recent endogenic or exogenic activity, i.e. volcanism or impacts. In particular, any indication that Titan is active at

the present day would have important implications for the sustenance of its atmosphere (Q7).

Early in the Cassini mission, excitement was generated by a reported detection of exposed carbon dioxide ice by the VIMS instrument (Barnes et al., 2005), although this was later disputed (Clark et al., 2010). However, the study of Clark et al. (2010) argued for a more robust spectral detection of benzene: somewhat surprisingly since benzene is a less abundant hydrocarbon in the atmosphere than others such as acetylene, which was not detected. Later, Solomonidou et al. (2018) proposed a latitudinal variation of Titan's surface composition, including exposed water ice as a major constituent at latitudes beyond $\pm 30^\circ$ N and S. At this time, identification of specific surface ices has remained highly challenging and controversial.

The Dragonfly mission promises to yield considerable insights into the surface composition, through IR and mass spectroscopy, seismology, gamma ray-neutron probe, and other means. In particular, it will make a detailed survey of the large impact crater Selk (Barnes et al., 2021; Lorenz et al., 2021), where there may be regions of exposed ice (Solomonidou et al., 2020). However, completing a global picture of exposed ice on Titan's surface will require a dedicated orbiter (Mitri et al., 2014a; Rodriguez et al., 2022) or perhaps some type of long-range aerorover, such as a balloon (Lorenz, 2008; Coustenis et al., 2009) or airplane (Barnes et al., 2012).

2.4. Questions about Titan's atmosphere (14-20)

14) What types of clouds exist in the troposphere and stratosphere?

Clouds on Titan (see Fig. 3b, 3c, Chapter 8) have been observed both directly, by imaging, and inferred indirectly by spectroscopy and modeling. The first evidence of large-scale clouds was obtained by brightening in the near infrared spectrum noticed by Griffith et al. (1998, 2000), and subsequently imaged using ground-based telescopes with adaptive optics (Brown et al., 2002; Brown et al., 2009; Brown et al., 2010; Schaller et al., 2006a,b; Bouchez and Brown, 2004; Roe et al., 2004). These were inferred to be clouds of methane in the upper troposphere.

Cassini imaging and spectroscopy revealed yet more clouds, including not only further instances of tropospheric clouds (Rodriguez et al., 2009; Turtle et al., 2009, 2011b,a; Turtle et al., 2018), but also stratospheric ice clouds: a polar ethane cloud in late northern winter Griffith et al. (2006), north polar cyanoacetylene and dicyanoacetylene ice clouds (Anderson et al., 2010, 2016),




Figure 3: Views of Titan's atmosphere. (a) Distant Titan from Cassini ISS (12/16/2011), showing the ubiquitous golden haze (NASA/JPL-Caltech/SSI/Kevin M. Gill); (b) Ethane cloud over Titan's north pole from Cassini VIMS (NASA/JPL/University of Arizona/LPGNantes) - NASA image PIA09171; (c) Titan south polar ice cloud of HCN, June 27, 2012 (NASA/JPL-Caltech/SSI); (d) Titan limb haze showing detached haze layer Sept. 24, 2010 (NASA/JPL-Caltech/SSI/Jason Major).

a south polar hydrogen cyanide cloud (de Kok et al., 2014), and a benzene ice cloud (Vinatier et al., 2018). The existence of many condensable species in Titan’s atmosphere implies that cloud composition can be more complex still, likely with layering of ices accumulated at different altitudes (Anderson et al., 2016).

A new front in cloud research has appeared in recent years due to the laboratory investigation of organic co-crystals, co-condensed phases of one or more molecules (Cable et al., 2014, 2018, 2019, 2020, 2021b; Czaplinski et al., 2023; Maynard-Casely et al., 2016; Vu et al., 2014, 2020b; Thakur and Remsing, 2023; Francis et al., 2023; Ennis et al., 2020). Understanding of these exotic substances is helping to inform our understanding of stratospheric cloud formation and growth and the materials that may be on Titan’s surface.

Current knowledge of Titan’s cloud composition and occurrence is spotty, due to a limited amount of high-resolution imagery and spectroscopy during Titan close flybys, combined with sporadic monitoring from the Earth. Given the complexity of cloud scenarios on Titan, a much more sustained and regular monitoring is necessary, together with cloud particle growth experiments in Earth laboratories.

15) How often does it rain on Titan?

Although we can be certain about clouds in Titan’s lower atmosphere, rainfall is more difficult to observe directly, and models include some amount of rainfall that evaporates before reaching the ground (known as ‘virgae’) (Barth and Rafkin, 2007; Graves et al., 2008). However, indirect evidence for rain exists through surface darkening observed after the passage of a large storm front, followed by a lightening of the surface observed some weeks later, apparently due to re-evaporation (Turtle et al., 2011b).

Better knowledge of the actual incidence and intensity of precipitation events is valuable however, since it has important consequences for pluvial and fluvial erosion (Lorenz and Lunine, 1996; Jaumann et al., 2008; Lorenz et al., 2008; Le Gall et al., 2010; Cartwright et al., 2011; Langhans et al., 2012; Black et al., 2012; Tewelde et al., 2013; Burr et al., 2013a; Aharonson et al., 2014; Moore et al., 2014; Howard et al., 2016; Neish et al., 2016; Faulk et al., 2017; Radebaugh et al., 2018; Matteoni et al., 2020; Miller et al., 2021; Lewis-Merrill et al., 2022), and the rate of replenishment of liquid bodies (lakes and seas) (Sotin et al., 2012). In addition, knowing the latitudinal and temporal pattern of rainfall can help to explain the observed dry equatorial regions and

the hemispheric asymmetry in lakes and seas (Aharonson et al., 2009; Lora et al., 2015; Tokano, 2019; Lora et al., 2022; Mitchell, 2008; Mitchell et al., 2009; Mitchell and Lora, 2016; Schneider et al., 2012; Birch et al., 2018).

Collecting such information will not be easy, requiring a sustained observational campaign best conducted from Titan orbit, as well as potentially observations from the surface, and ideally in-situ sampling, for example by high-altitude balloons (Lorenz, 2008; Coustenis et al., 2009). Further modeling work will also help to determine whether observed rainfall can be adequately explained.

16) What is the availability of P and S compounds for chemistry?

Six elements are considered the most vital for life on Earth: H, C, N, O, P and S ('CHNOPS'), together constituting 98% of living matter (Cockell et al., 2016). To date four of the six have been found on Titan (H, C, N, O), for example in hydrogen cyanide and carbon monoxide, while all six have been found or strongly suspected in Enceladus plume material (Waite et al., 2009; Cable et al., 2021a; Postberg et al., 2023). Therefore, determining whether there is available phosphorus and sulfur on Titan assumes an astrobiological significance (Raulin et al., 2012), as well as a geochemical one (Fortes et al., 2007; Pasek et al., 2011). Searches have been made for atmospheric hydrogen sulfide, carbon sulfide and phosphine with null result so far (Nixon et al., 2013; Teanby et al., 2018).

Attempts to detect P and S in the atmosphere can continue to be made by astronomical observations and remote sensing, e.g. with ALMA, IRTF etc; while the Dragonfly mission in the 2030s will be able to sample both the atmosphere and surface with excellent ability through its mass spectrometer to detect any P and S compounds.

17) What is the composition of Titan's aerosols?

Understanding Titan's organic particulate aerosols - which give rise to its famous golden appearance (Fig. 3a) and multiple haze layers (Fig. 3d) - is important for many reasons (see also Chapters 6 and 7). First, these are critical in the radiative balance of Titan's atmosphere, including regulation of its 'greenhouse effect' (McKay et al., 1991). Second, the chemical complexity of the aerosols is vital for understanding the astrobiological potential of Titan's atmosphere (Khare et al., 1986; Neish et al., 2010; Hörlst et al., 2012; Kawai et al., 2013; Cleaves II et al., 2014; He and Smith, 2014; Kawai et al., 2019; Gautier et al., 2014). Third, they serve as condensation nucleii

for cloud particles and droplets (McKay et al., 2001; Hörst and Tolbert, 2013; Burgalat et al., 2014). And fourth, the aerosols undoubtedly sediment onto the surface, where they may form a significant component of the equatorial dunes (Barnes et al., 2008, 2015; Yu et al., 2017b).

Since the 1980s there has been a sustained effort to replicate Titan's atmospheric haze particles through laboratory experiments, in which vacuum vessels filled with varying mixtures of nitrogen, methane and other gases are reacted together by energetic stimulation with UV or electric current (see review by Cable et al., 2011). Their fractal nature has been elucidated by modeling and SEM techniques (Cabane et al., 1993; Rannou et al., 1995, 2003, 1999; Bar-Nun et al., 2008; Schulz et al., 2021).

During their atmospheric descent, aerosols are moreover predicted to chemically and physically evolve (Lavvas et al., 2011; Carrasco et al., 2018; Perrin et al., 2021). In the ionosphere, aerosols have been shown to further influence their atmospheric environment (Chatain et al., 2023), with an impact on the electron populations detected by the Cassini RPWS/LP instrument (Chatain et al., 2021).

A direct attempt was made to sample aerosol material with Huygens' ACP instrument (Aerosol Collector Pyrolyzer, Israël et al., 2003), which was designed to pass pyrolysis products to the mass spectrometer (GCMS - Gas Chromatograph and Mass Spectrometer, Niemann et al., 2002). The results showed significant amounts of hydrogen cyanide and ammonia (Israël et al., 2005; Coll et al., 2013).

Sampling and elucidating the aerosol composition and physical properties is a major goal for future missions (Mitri et al., 2021; Rodriguez et al., 2022), since the aerosols are one of Titan's most distinguishing features. The DraMS instrument on Dragonfly will be able to measure the composition of surface organics from the dune fields (Trainer et al., 2021), however direct sampling of atmospheric aerosols remains a future goal. In addition, instruments other than MS such as microscopy can play a role in illuminating the physical properties. Further laboratory work can expand our concept of the aerosol parameter space, and also the reactions that they can undergo as nanoparticles in the ionosphere until they sediment as submillimetric particles at Titan's surface.

18) What is the cause of the tilted atmospheric rotation?

Early results from mapping of Titan's atmospheric temperature contours by Cassini CIRS revealed an unexpected phenomenon: the axis of the atmo-

spheric rotation pole was offset by some \sim 3.5° from the solid body rotation axis (Achterberg et al., 2008). Later studies confirmed this finding (Roman et al., 2009; Teanby et al., 2010; Cordiner et al., 2019; Sharkey et al., 2020; Kutsop et al., 2022). To date, an explanation for this phenomenon has remained elusive (Nixon et al., 2018), and further climate simulations are required to better understand how it occurs and whether it is changing over time.

19) Are there important biological precursor molecules produced in the atmosphere?

The production of hydrocarbons and nitriles in Titan's atmosphere was predicted by photochemical models as far back as the 1970s (Strobel, 1974), validated soon thereafter by mid-infrared spectroscopy from Voyager 1, which revealed a rich tapestry of molecular vibrations throughout the mid-infrared (Hanel et al., 1981; Maguire et al., 1981; Kunde et al., 1981). Since then, the known molecular inventory has been expanded repeatedly, e.g. by ISO (Coustenis et al., 1998, 2003), and most revealingly by direct sampling with Cassini INMS and CAPS (Niemann et al., 2005; Waite et al., 2005, 2007; Vuitton et al., 2007; Niemann et al., 2010) - see also Chapter 7.

Laboratory investigations have shown that simulated Titan photochemical products, produced by reactions starting with nitrogen and methane reagents, can be hydrolyzed to produce amino acids in a manner similar to a Miller-Urey experiment (Khare et al., 1986; Neish et al., 2010; Cleaves II et al., 2014). Other experiments showed that such amino acids can even be produced directly in the atmosphere when oxygen is included in the reactions by admission of carbon monoxide (Hörst et al., 2012).

The combination of the detected complex organic chemistry along with the laboratory simulations showing the potential for prebiotic molecules to arise has naturally led to the question of whether such prebiotic molecules also occur on Titan. Despite some of Cassini's instruments (especially CAPS) being able to detect the presence of large organic molecules, the instruments did not have the ability determine the molecular composition or structure.

In future, the search for evidence of biological precursor molecules on Titan can proceed on two fronts. First, astronomical measurements will continue to push the boundaries of the known atmospheric inventory through infrared and sub-millimeter spectroscopy (e.g. Cordiner et al., 2015; Palmer et al., 2017; Lombardo et al., 2019a; Thelen et al., 2020; Nixon et al., 2020). This will be enabled by upgraded capabilities at existing observatories and

entirely new telescopes on the ground and in space (see Section 3 of this chapter), backed up by additional laboratory investigations of gases and ices at Titan relevant temperatures. In addition, in situ measurements are needed with a more sophisticated chemical analyzer: this will be achieved by the DraMS instrument onboard Dragonfly (Barnes et al., 2021; Trainer et al., 2021).

20) Is Titan’s atmosphere stable over time?

Question 3 above described the uncertainties regarding the origin of Titan’s atmosphere: a closely linked question is about its fate. Much depends on whether methane is being resupplied to the atmosphere or not (Chapter 13), as well as its rate of photochemical depletion (see Chapters 6 and 7). Proposed mechanisms of endogenous resupply include: destabilization of crustal clathrates (Tobie et al., 2006); displacement of surficial clathrates (Choukroun and Sotin, 2012); and geochemical production in the interior from CO₂ through serpentinization (Glein, 2015), although Fortes (2012) suggests that deep interior methane will be captured in clathrate and not escape to the atmosphere.

On the other hand, if methane is being irreversibly depleted, then eventually its current methane greenhouse will disappear, being replaced by a much cooler climate under a nitrogen greenhouse (Charnay et al., 2014b), or the atmosphere may collapse altogether into a ‘snowball’ state (Lorenz et al., 1997; Wong et al., 2015).

Would such a condition persist forever? Impacts may work to reverse the snowball state by vaporizing enough methane into the atmosphere to reconstitute something resembling the warmer state we see today; alternatively Titan under a future red giant Sun will have a dramatically different climate Lorenz et al. (1997).

Predicting the future state of the atmosphere by climate modeling requires most critically a firmer understanding of whether methane is being resupplied to the atmosphere by some mechanism (e.g. Tobie et al., 2006; Glein, 2015) or not. Closing this loophole in our knowledge in turn will hinge on a combination of investigative efforts: determining if the atmosphere has already changed over time, perhaps by sensitive isotopic measurements and modeling, and searching for methane resupply by cryovolcanism (surface changes, Solomonidou et al., 2016), or buffering by the solid (clathrates, Davies et al., 2016) or liquid surface (lakes and seas, Hayes, 2016).

Instrumentation that could help guide us to these goals includes: ice

penetrating RADAR; visible and near-IR imagery and spectroscopy (spectro-imaging), mass spectroscopy for improved measurements of isotopic ratios and noble gas abundances (Coustenis et al., 2009; Tobie et al., 2014; Mitri et al., 2021; Sulaiman et al., 2021; Rodriguez et al., 2022). Therefore, further orbital and *in situ* investigation is necessary.

3. Future Directions

We conclude this chapter - and book - by mentioning again some of the means by which the questions raised in this chapter may be answered in future. The categorial divisions here echo the four main techniques introduced at the start of the chapter (astronomy, laboratory work, modeling, missions), which work synergistically together to collect and interpret data (Fig. 4). In addition, we propose the increased use of a fifth type of investigation: terrestrial field analog studies.




Figure 4: Synergies between data collection (astronomy, missions) and interpretation (modeling, laboratory investigations). None of these techniques works in isolation, but serve to reinforce each other to derive new knowledge.




Figure 5: Future means of probing Titan. (a) NASA’s James Webb Space Telescope launched 12/15/2021 (image: NASA); (b) NASA’s Dragonfly mission to Titan scheduled for launch in 2028 (NASA/APL); (c) Artists impression of the completed 39m Extremely Large Telescope (ELT) being built by the European Southern Observatory (ESO), scheduled for first light ~2029 (Swinburne Astronomy Productions/ESO); (d) Atacama Large Millimeter/submillimeter Array (ALMA).

As mentioned at the opening of this chapter, ground and (later) space-based telescopes have played an important role in unraveling Titan's mysteries.² This has included spectroscopy to determine its gases, imaging with adaptive optics to view the surface and clouds, and synthetic aperture interferometry for imaging and spectroscopy at longer wavelengths.

At the time of writing, the James Webb Space Telescope (JWST, Fig. 5(a)) is the newest of the large, general purpose space telescopes to come on-line, and has already begun to make execute a planned range of spectroscopic and imaging observations of Titan (Nixon et al., 2016a). On the ground, many optical/IR telescopes are routinely used to make observations of Titan, such as cloud tracking (Griffith et al., 2000; Brown et al., 2002; Roe et al., 2002; De Pater et al., 2006; Schaller et al., 2006a,b; Ádámkovics et al., 2007; Schaller et al., 2009; Brown et al., 2009; Ádámkovics et al., 2016; Ádámkovics and de Pater, 2017), spectroscopy (Roe et al., 2003; Lellouch et al., 2003; Ádámkovics et al., 2004; Penteado et al., 2005; Penteado and Griffith, 2010; Ádámkovics et al., 2016; Lombardo et al., 2019b) and occultation observations (Hubbard et al., 1990; Sicardy et al., 1990, 1999, 2006; Zalucha et al., 2007), including large telescopes with adaptive optics such as Keck, the Very Large Telescope (VLT), Gemini and others. In future, even larger telescopes currently planned such as the Giant Magellan Telescope (GMT, 25 m, Fanson et al., 2018, 2020, 2022) the Thirty Meter Telescope (TMT, Nelson and Sanders, 2008; Sanders, 2013; Skidmore et al., 2015) and the Extremely Large Telescope (ELT, 39 m, Gilmozzi and Spyromilio, 2007; Tamai and Spyromilio, 2014) (see Fig. 5(c)) will greatly improve our ability to investigate Titan's atmosphere and surface from Earth.

Large surface interferometric arrays such as the Karl Jansky Very Large Array (VLA), the Very Long Baseline Array (VLBA), the Atacama Large Millimeter-submillimeter Array (ALMA, Fig. 5(d)), and the Northern Extended Millimeter Array (NOEMA) are making important observations at long wavelengths (Muhleman et al., 1990; Butler and Gurwell, 2004; Witasse et al., 2006; Cordiner et al., 2014; Serigano et al., 2016; Molter et al., 2016; Lai et al., 2017; Palmer et al., 2017; Thelen et al., 2018; Lellouch et al., 2019; Nixon et al., 2020), and are poised to be upgraded in coming decades with wider receiver bands, improved sensitivity and longer baselines.

²Indeed, without the invention of the telescope we not know of its existence at all.

Laboratory Work

Data derived from observations and measurements of Titan must be interpreted, and observations and instruments must be carefully designed to maximise science. Laboratory work in many varieties has proven essential in this regard, as mentioned earlier in this chapter. In future, many more laboratory investigations are necessary including: high-resolution gas phase spectroscopy of molecules; spectroscopy of ices, especially mixed phases; creation and analysis of analogs (tholins) under a wide variety of conditions; erosional experiments at realistic Titan conditions; gas reaction and gas-particle reaction kinetics; and many others besides. An important area of current work is achieving consistency between results obtained at different laboratories, which has begun to motivate intercomparison efforts (Li et al., 2022).

Modeling

Modeling has always played a crucial role in our understanding of Titan, including photochemical models of the atmosphere; interior structure models; formation and dynamical evolution models; cloud models and GCMs; and others. With ever-increasing computing power, our capability to run models at higher levels of fidelity is always increasing (e.g. reducing grid size in GCMs, running photochemical models with larger networks and over more timesteps). In tandem, better constraints are needed for these models, including reaction rates, measured temperature and wind fields, seasonal observations of gas distributions, and more. Given the very large of possible reactions that can be included in photochemical models, work to define the most important pathways by Monte Carlo approaches is especially critical (Hébrard et al., 2005, 2006, 2007), to allow focusing of laboratory and theoretical work on the most influential reactions.

Missions

Despite all the advances that can be made on Earth and using remote observations, the ultimate treasure trove of Titan knowledge is Titan itself. Hence, there is often no substitute for in situ measurements, covering everything from atmospheric temperature, pressure and gas composition, to surface morphology, land and sea composition, and much more. To this end, Titan scientists are excited about the upcoming Dragonfly mission (Fig. 5(b)) which will make a host of in situ measurements of Titan's dune fields and the environs of Selk crater. Even more ambitious missions have been proposed in-

cluding orbiters, balloons, airplanes, boats and submarines (Coustenis et al., 2009; Stofan et al., 2013; Barnes et al., 2012; Mitri et al., 2014a; Tobie et al., 2014; Oleson et al., 2015; Nixon et al., 2016b, 2019; Rodriguez et al., 2022; Barnes et al., 2022).

Terrestrial Field Analog Studies

Historically, terrestrial field studies of planetary analog environments have proven highly beneficial for both interpreting returned data (especially lunar and martian imagery) but also for testing instrumentation and the training of astronauts. To date terrestrial field studies targeting Titan analog sites have been limited and primarily focused on dune analogs (Radebaugh et al., 2010). However, the types of useful terrestrial analog environments for Titan may in fact be much larger (e.g. ices, rivers, lakes, seas, karst, polar stratospheric clouds, to name a few), providing fertile new avenues for research.

Acknowledgements

The authors wish to thank the editors of the book, Dr Rosaly Lopes and Dr Anezina Solomonidou, along with three referees - Dr Ralph Lorenz and two anonymous reviewers - for helpful feedback provided during the preparation of this chapter.

References

- Achterberg, R. K., Conrath, B. J., Gierasch, P. J., Flasar, F. M., Nixon, C. A., Oct. 2008. Observation of a tilt of Titan's middle-atmospheric superrotation. *Icarus* 197, 549–555.
- Ádámkovics, M., de Pater, I., 2017. Observations of the global haze redistribution on Titan from 2006 to 2015 with OSIRIS at Keck. *Icarus* 290, 134–149.
- Ádámkovics, M., de Pater, I., Roe, H., Gibbard, S., Griffith, C., 2004. Spatially-resolved spectroscopy at 1.6 μm of Titan's atmosphere and surface. *Geophysical Research Letters* 31 (17).
- Ádámkovics, M., Mitchell, J. L., Hayes, A. G., Rojo, P. M., Corlies, P., Barnes, J. W., Ivanov, V. D., Brown, R. H., Baines, K. H., Buratti, B. J., et al., 2016. Meridional variation in tropospheric methane on Titan observed with AO spectroscopy at Keck and VLT. *Icarus* 270, 376–388.

- Ádámkovics, M., Wong, M. H., Laver, C., de Pater, I., 2007. Widespread morning drizzle on Titan. *Science* 318 (5852), 962–965.
- Aharonson, O., Hayes, A., Lopes, R., Lucas, A., Hayne, P., Perron, J., Soderblom, L., 2014. Titan's surface geology. In: *Titan: Surface, Atmosphere and Magnetosphere*. Cambridge Univ. Press Cambridge, U. K, pp. 43–75.
- Aharonson, O., Hayes, A., Lunine, J., Lorenz, R., Allison, M., Elachi, C., 2009. An asymmetric distribution of lakes on Titan as a possible consequence of orbital forcing. *Nature Geoscience* 2 (12), 851–854.
- Anderson, C., Nna-Mvondo, D., Samuelson, R., McLain, J., Dworkin, J., 2018. The SPECTRAL ice chamber: application to Titan's stratospheric ice clouds. *The Astrophysical Journal* 865 (1), 62.
- Anderson, C. M., Samuelson, R. E., Bjomaker, G. L., Achterberg, R. K., Jun. 2010. Particle size and abundance of HC_3N ice in Titan's lower stratosphere at high northern latitudes. *Icarus* 207, 914–922.
- Anderson, C. M., Samuelson, R. E., Yung, Y. L., McLain, J. L., Apr 2016. Solid-state photochemistry as a formation mechanism for Titan's stratospheric C_4N_2 ice clouds. *Geophysical Research Letters* 43 (7), 3088–3094. URL <http://dx.doi.org/10.1002/2016GL067795>
- Artemieva, N., Lunine, J., Aug. 2003. Cratering on Titan: impact melt, ejecta, and the fate of surface organics. *Icarus* 164, 471–480.
- Balucani, N., Leonori, F., Petrucci, R., Stazi, M., Skouteris, D., Rosi, M., Casavecchia, P., 2010. Formation of nitriles and imines in the atmosphere of Titan: combined crossed-beam and theoretical studies on the reaction dynamics of excited nitrogen atoms $\text{N}({}^2\text{D})$ with ethane. *Faraday Discussions* 147, 189–216.
- Bar-Nun, A., Dimitrov, V., Tomasko, M., 2008. Titan's aerosols: Comparison between our model and DISR findings. *Planetary and Space Science* 56 (5), 708–714.
- Barnes, J. W., Brown, R. H., Soderblom, L., Sotin, C., Le Mouëlic, S., Rodriguez, S., Jaumann, R., Beyer, R. A., Buratti, B. J., Pitman, K.,

et al., 2008. Spectroscopy, morphometry, and photoclinometry of Titan's dunefields from Cassini/VIMS. *Icarus* 195 (1), 400–414.

Barnes, J. W., Brown, R. H., Turtle, E. P., McEwen, A. S., Lorenz, R. D., Janssen, M., Schaller, E. L., Brown, M. E., Buratti, B. J., Sotin, C., Griffith, C., Clark, R., Perry, J., Fussner, S., Barbara, J., West, R., Elachi, C., Bouchez, A. H., Roe, H. G., Baines, K. H., Bellucci, G., Bibring, J., Capaccioni, F., Cerroni, P., Combes, M., Coradini, A., Cruikshank, D. P., Drossart, P., Formisano, V., Jaumann, R., Langevin, Y., Matson, D. L., McCord, T. B., Nicholson, P. D., Sicardy, B., Oct. 2005. A 5-Micron-Bright Spot on Titan: Evidence for Surface Diversity. *Science* 310, 92–95.

Barnes, J. W., Hayes, A. G., Soderblom, J. M., Mackenzie, S. M., Hofgartner, J. D., Lorenz, R. D., Turtle, E. P., Radebaugh, J., Burr, D., Lora, J., et al., 2022. New Frontiers Titan Orbiter. In: Planetary Science and Astrobiology Decadal Survey 2023-2032 white paper e-id. 317; Bulletin of the American Astronomical Society. Vol. 53.

Barnes, J. W., Lemke, L., Foch, R., McKay, C. P., Beyer, R. A., Radebaugh, J., Atkinson, D. H., Lorenz, R. D., Le Mouélic, S., Rodriguez, S., et al., 2012. AVIATR—Aerial Vehicle for In-situ and Airborne Titan Reconnaissance: A Titan airplane mission concept. *Experimental Astronomy* 33, 55–127.

Barnes, J. W., Lorenz, R. D., Radebaugh, J., Hayes, A. G., Arnold, K., Chandler, C., 2015. Production and global transport of Titan's sand particles. *Planetary Science* 4 (1), 1–19.

Barnes, J. W., Radebaugh, J., Brown, R. H., Wall, S., Soderblom, L., Lunine, J., Burr, D., Sotin, C., Le Mouélic, S., Rodriguez, S., et al., 2007. Near-infrared spectral mapping of Titan's mountains and channels. *Journal of Geophysical Research: Planets* 112 (E11).

Barnes, J. W., Turtle, E. P., Trainer, M. G., Lorenz, R. D., MacKenzie, S. M., Brinckerhoff, W. B., Cable, M. L., Ernst, C. M., Freissinet, C., Hand, K. P., Hayes, A. G., Hörist, S. M., Johnson, J. R., Karkoschka, E., Lawrence, D. J., Gall, A. L., Lora, J. M., McKay, C. P., Miller, R. S., Murchie, S. L., Neish, C. D., Newman, C. E., Núñez, J., Panning, M. P., Parsons, A. M., Peplowski, P. N., Quick, L. C., Radebaugh, J., Rafkin, S. C. R., Shiraishi, H., Soderblom, J. M., Sotzen, K. S., Stickle, A. M.,

- Stofan, E. R., Szopa, C., Tokano, T., Wagner, T., Wilson, C., Yingst, R. A., Zacny, K., Stähler, S. C., jul 2021. Science Goals and Objectives for the Dragonfly Titan Rotorcraft Relocatable Lander. *The Planetary Science Journal* 2 (4), 130.
- Barth, E. L., Rafkin, S. C., 2007. TRAMS: A new dynamic cloud model for Titan's methane clouds. *Geophysical Research Letters* 34 (3).
- Barth, E. L., Toon, O. B., 2004. Properties of methane clouds on Titan: Results from microphysical modeling. *Geophysical Research Letters* 31 (17).
- Barth, E. L., Toon, O. B., 2006. Methane, ethane, and mixed clouds in Titan's atmosphere: Properties derived from microphysical modeling. *Icarus* 182 (1), 230–250.
- Bernath, P., Fernando, A. M., 2021. Infrared absorption cross sections for hot isobutane in the CH stretching region. *Journal of Quantitative Spectroscopy and Radiative Transfer* 269, 107644.
- Bernath, P. F., Dodangodage, R., Zhao, J., Billinghamurst, B., 2023. Infrared absorption cross sections for propene broadened by N₂ (450–1250 cm⁻¹) and by H₂ (2680–3220 cm⁻¹). *Journal of Quantitative Spectroscopy and Radiative Transfer* 296, 108462.
- Birch, S., Hayes, A., Corlies, P., Stofan, E., Hofgartner, J., Lopes, R., Lorenz, R., Lunine, J., MacKenzie, S., Malaska, M., et al., 2018. Morphological evidence that Titan's southern hemisphere basins are paleoseas. *Icarus* 310, 140–148.
- Birch, S. P., Parker, G., Corlies, P., Soderblom, J. M., Miller, J. W., Palermo, R. V., Lora, J. M., Ashton, A. D., Hayes, A. G., Perron, J. T., 2023. Reconstructing river flows remotely on Earth, Titan, and Mars. *Proceedings of the National Academy of Sciences* 120 (29), e2206837120.
- Black, B. A., Perron, J. T., Burr, D. M., Drummond, S. A., Aug. 2012. Estimating erosional exhumation on Titan from drainage network morphology. *Journal of Geophysical Research (Planets)* 117, 8006.
- Bouchez, A. H., Brown, M. E., 2004. Statistics of Titan's south polar tropospheric clouds. *The Astrophysical Journal* 618 (1), L53.

- Brown, M., Schaller, E., Roe, H., Chen, C., Roberts, J., Brown, R., Baines, K., Clark, R., 2009. Discovery of lake-effect clouds on Titan. *Geophysical Research Letters* 36 (1).
- Brown, M. E., Bouchez, A. H., Griffith, C. A., Dec. 2002. Direct detection of variable tropospheric clouds near Titan's south pole. *Nature* 420, 795–797.
- Brown, M. E., Roberts, J. E., Schaller, E. L., Feb. 2010. Clouds on Titan during the Cassini prime mission: A complete analysis of the VIMS data. *Icarus* 205, 571–580.
- Brown, R. H., Soderblom, L. A., Soderblom, J. M., Clark, R. N., Jaumann, R., Barnes, J. W., Sotin, C., Buratti, B., Baines, K. H., Nicholson, P. D., jul 2008. The identification of liquid ethane in Titan's Ontario Lacus. *Nature* 454 (7204), 607–610.
- Burgalat, J., Rannou, P., Cours, T., Riviere, E., 2014. Modeling cloud microphysics using a two-moments hybrid bulk/bin scheme for use in Titan's climate models: Application to the annual and diurnal cycles. *Icarus* 231, 310–322.
- Burkhard, L. M., Smith-Konter, B. R., Fagents, S. A., Cameron, M. E., Collins, G. C., Pappalardo, R. T., 2022. Strike-slip faulting on Titan: Modeling tidal stresses and shear failure conditions due to pore fluid interactions. *Icarus* 371, 114700.
- Burr, D. M., Bridges, N. T., Marshall, J. R., Smith, J. K., White, B. R., Emery, J. P., 2015a. Higher-than-predicted saltation threshold wind speeds on Titan. *Nature* 517 (7532), 60–63.
- Burr, D. M., Bridges, N. T., Smith, J. K., Marshall, J. R., White, B. R., Williams, D. A., 2015b. The Titan wind tunnel: A new tool for investigating extraterrestrial aeolian environments. *Aeolian Research* 18, 205–214.
- Burr, D. M., Drummond, S. A., Cartwright, R., Black, B. A., Perron, J. T., Sep. 2013a. Morphology of fluvial networks on Titan: Evidence for structural control. *Icarus* 226, 742–759.
- Burr, D. M., Perron, J. T., Lamb, M. P., Irwin, R. P., Collins, G. C., Howard, A. D., Sklar, L. S., Moore, J. M., Ádámkovics, M., Baker, V. R., Drummond, S. A., Black, B. A., March/April 2013b. Fluvial features on Titan: Insights from morphology and modeling. *GSA Bulletin* 125 (3/4), 299–321.

- Burr, D. M., Sutton, S. L., Emery, J. P., Nield, E. V., Kok, J. F., Smith, J. K., Bridges, N. T., 2020. A wind tunnel study of the effect of intermediate density ratio on saltation threshold. *Aeolian Research* 45, 100601.
- Butler, B., Gurwell, M., 2004. Radio wavelength observations of Titan with the VLA. In: *AAS/Division for Planetary Sciences Meeting Abstracts #36*. Vol. 36. pp. 06–04.
- Cabane, M., Rannou, P., Chassefière, E., Israel, G., 1993. Fractal aggregates in Titan's atmosphere. *Planetary and Space Science* 41 (4), 257–267.
- Cable, M. L., Hörst, S. M., Hodyss, R., Beauchamp, P. M., Smith, M. A., Willis, P. A., Nov 2011. Titan Tholins: Simulating Titan Organic Chemistry in the Cassini-Huygens Era. *Chemical Reviews* 112 (3), 1882–1909.
URL <https://doi.org/10.1021/cr200221x>
- Cable, M. L., Porco, C., Glein, C. R., German, C. R., MacKenzie, S. M., Neveu, M., Hoehler, T. M., Hofmann, A. E., Hendrix, A. R., Eigenbrode, J., et al., 2021a. The science case for a return to Enceladus. *The Planetary Science Journal* 2 (4), 132.
- Cable, M. L., Runčevski, T., Maynard-Casely, H. E., Vu, T. H., Hodyss, R., 2021b. Titan in a test tube: organic co-crystals and implications for Titan mineralogy. *Accounts of Chemical Research* 54 (15), 3050–3059.
- Cable, M. L., Vu, T. H., Hodyss, R., Choukroun, M., Malaska, M. J., Beauchamp, P., aug 2014. Experimental determination of the kinetics of formation of the benzene-ethane co-crystal and implications for Titan. *Geophysical Research Letters* 41 (15), 5396–5401.
URL <https://doi.org/10.1002/2014gl060531>
- Cable, M. L., Vu, T. H., Malaska, M. J., Maynard-Casely, H. E., Choukroun, M., Hodyss, R., nov 2019. A Co-Crystal between Acetylene and Butane: A Potentially Ubiquitous Molecular Mineral on Titan. *ACS Earth and Space Chemistry* 3 (12), 2808–2815.
URL <https://doi.org/10.1021/acsearthspacechem.9b00275>
- Cable, M. L., Vu, T. H., Malaska, M. J., Maynard-Casely, H. E., Choukroun, M., Hodyss, R., jul 2020. Properties and Behavior of the Acetonitrile–Acetylene Co-Crystal under Titan Surface Conditions. *ACS Earth*

and Space Chemistry 4 (8), 1375–1385.

URL <https://doi.org/10.1021/acsearthspacechem.0c00129>

Cable, M. L., Vu, T. H., Maynard-Casely, H. E., Choukroun, M., Hoddyss, R., feb 2018. The Acetylene-Ammonia Co-crystal on Titan. ACS Earth and Space Chemistry 2 (4), 366–375.

URL <https://doi.org/10.1021/acsearthspacechem.7b00135>

Carrasco, N., Bourgalais, J., Vettier, L., Pernot, P., Giner, E., Spezia, R., 2022. A missing link in the nitrogen-rich organic chain on Titan. Astronomy & Astrophysics 663, A165.

Carrasco, N., Tigrine, S., Gavilan, L., Nahon, L., Gudipati, M. S., 2018. The evolution of Titan's high-altitude aerosols under ultraviolet irradiation. Nature Astronomy 2 (6), 489–494.

Cartwright, R., Clayton, J. A., Kirk, R. L., 2011. Channel morphometry, sediment transport, and implications for tectonic activity and surficial ages of Titan basins. Icarus 214 (2), 561–570.

Charnay, B., Forget, F., Tobie, G., Sotin, C., Wordsworth, R., 2014a. Titan's past and future: 3D modeling of a pure nitrogen atmosphere and geological implications. Icarus 241, 269–279.

Charnay, B., Forget, F., Tobie, G., Sotin, C., Wordsworth, R., 2014b. Titan's past and future: 3D modeling of a pure nitrogen atmosphere and geological implications. Icarus 241, 269–279.

Chatain, A., Carrasco, N., Vettier, L., Guaitella, O., 2023. Interaction dust– plasma in Titan's ionosphere: Feedbacks on the gas phase composition. Icarus 396, 115502.

Chatain, A., Rafkin, S. C., Soto, A., Hueso, R., Spiga, A., 2022. Air–Sea Interactions on Titan: Effect of Radiative Transfer on the Lake Evaporation and Atmospheric Circulation. The Planetary Science Journal 3 (10), 232.

Chatain, A., Wahlund, J.-E., Shebanits, O., Hadid, L. Z., Morooka, M., Edberg, N. J., Guaitella, O., Carrasco, N., 2021. Re-analysis of the Cassini RPWS/LP data in Titan's ionosphere: 1. Detection of several electron populations. Journal of Geophysical Research: Space Physics 126 (8), e2020JA028412.

- Choukroun, M., Grasset, O., Tobie, G., Sotin, C., Feb. 2010. Stability of methane clathrate hydrates under pressure: Influence on outgassing processes of methane on Titan. *Icarus* 205, 581–593.
- Choukroun, M., Sotin, C., 2012. Is Titan's shape caused by its meteorology and carbon cycle? *Geophys. Res. Lett.* 39, L04201.
- Clark, R. N., Curchin, J. M., Barnes, J. W., Jaumann, R., Soderblom, L., Cruikshank, D. P., Brown, R. H., Rodriguez, S., Lunine, J., Stephan, K., Hoefen, T. M., Le Mouélic, S., Sotin, C., Baines, K. H., Buratti, B. J., Nicholson, P. D., 2010. Detection and mapping of hydrocarbon deposits on Titan. *Journal of Geophysical Research: Planets* 115 (E10), n/a–n/a, e10005.
URL <http://dx.doi.org/10.1029/2009JE003369>
- Cleaves II, H. J., Neish, C., Callahan, M. P., Parker, E., Fernández, F. M., Dworkin, J. P., 2014. Amino acids generated from hydrated Titan tholins: Comparison with Miller–Urey electric discharge products. *Icarus* 237, 182–189.
- Cockell, C. S., Bush, T., Bryce, C., Direito, S., Fox-Powell, M., Harrison, J. P., Lammer, H., Landenmark, H., Martin-Torres, J., Nicholson, N., et al., 2016. Habitability: a review. *Astrobiology* 16 (1), 89–117.
- Coll, P., Navarro-González, R., Szopa, C., Poch, O., Ramírez, S. I., Coscia, D., Raulin, F., Cabane, M., Buch, A., Israël, G., 2013. Can laboratory tholins mimic the chemistry producing Titan's aerosols? A review in light of ACP experimental results. *Planetary and Space Science* 77, 91–103.
- Cook-Hallett, C., Barnes, J. W., Kattenhorn, S. A., Hurford, T., Radebaugh, J., Stiles, B., Beuthe, M., 2015. Global contraction/expansion and polar lithospheric thinning on Titan from patterns of tectonism. *Journal of Geophysical Research: Planets* 120 (6), 1220–1236.
- Cordier, D., Bonhommeau, D. A., Vu, T. H., Choukroun, M., García-Sánchez, F., 2021. Vertical compositional variations of liquid hydrocarbons in Titan's alkanofers. *Astronomy & Astrophysics* 653, A80.
- Cordier, D., Mousis, O., Lunine, J., Lebonnois, S., Rannou, P., Lavvas, P., Lobo, L., Ferreira, A., 2012. Titan's lakes chemical composition: Sources

of uncertainties and variability. Planetary and Space Science 61 (1), 99–107, Surfaces, atmospheres and magnetospheres of the outer planets and their satellites and ring systems: Part VII.

Cordier, D., Mousis, O., Lunine, J. I., Lavvas, P., Vuitton, V., dec 2009. An Estimate of the Chemical Composition of Titan’s Lakes. *The Astrophysical Journal* 707 (2), L128–L131.

Cordiner, M., Palmer, M., Nixon, C., Irwin, P., Teanby, N., Charnley, S., Mumma, M., Kisiel, Z., Serigano, J., Kuan, Y.-J., et al., 2015. Ethyl cyanide on Titan: Spectroscopic detection and mapping using ALMA. *The Astrophysical Journal Letters* 800 (1), L14.

Cordiner, M. A., Nixon, C. A., Teanby, N. A., Irwin, P. G. J., Serigano, J., Charnley, S. B., Milam, S. N., Mumma, M. J., Lis, D. C., Villanueva, G., Paganini, L., Kuan, Y.-J., Remijan, A. J., Nov. 2014. ALMA Measurements of the HNC and HC₃N Distributions in Titan’s Atmosphere. *Astrophys. J. Lett.* 795, L30.

Cordiner, M. A., Teanby, N. A., Nixon, C. A., Vuitton, V., Thelen, A. E., Charnley, S. B., Jul 2019. ALMA Spectral Imaging of Titan Contemporaneous with Cassini’s Grand Finale. *The Astronomical Journal* 158 (2), 76. URL <http://dx.doi.org/10.3847/1538-3881/ab2d20>

Coustenis, A., Atreya, S. K., Balint, T., Brown, R. H., Dougherty, M. K., Ferri, F., Fulchignoni, M., Gautier, D., Gowen, R. A., Griffith, C. A., Gurvits, L. I., Jaumann, R., Langevin, Y., Leese, M. R., Lunine, J. I., McKay, C. P., Moussas, X., Müller-Wodarg, I., Neubauer, F., Owen, T. C., Raulin, F., Sittler, E. C., Sohl, F., Sotin, C., Tobie, G., Tokano, T., Turtle, E. P., Wahlund, J.-E., Waite, J. H., Baines, K. H., Blamont, J., Coates, A. J., Dandouras, I., Krimigis, T., Lellouch, E., Lorenz, R. D., Morse, A., Porco, C. C., Hirtzig, M., Saur, J., Spilker, T., Zarnecki, J. C., Choi, E., Achilleos, N., Amils, R., Annan, P., Atkinson, D. H., Bénilan, Y., Bertucci, C., Bézard, B., Björaker, G. L., Blanc, M., Boireau, L., Bouman, J., Cabane, M., Capria, M. T., Chassefière, E., Coll, P., Combes, M., Cooper, J. F., Coradini, A., Crary, F., Cravens, T., Daglis, I. A., de Angelis, E., de Bergh, C., de Pater, I., Dunford, C., Durry, G., Dutuit, O., Fairbrother, D., Flasar, F. M., Fortes, A. D., Frampton, R., Fujimoto, M., Galand, M., Grasset, O., Grott, M., Haltigin, T., Herique, A., Hersant, F., Hussmann,

- H., Ip, W., Johnson, R., Kallio, E., Kempf, S., Knapmeyer, M., Kofman, W., Koop, R., Kostiuk, T., Krupp, N., Küppers, M., Lammer, H., Lara, L.-M., Lavvas, P., Le Mouélic, S., Lebonnois, S., Ledvina, S., Li, J., Livenessgood, T. A., Lopes, R. M., Lopez-Moreno, J.-J., Luz, D., Mahaffy, P. R., Mall, U., Martinez-Frias, J., Marty, B., McCord, T., Menor Salvan, C., Milillo, A., Mitchell, D. G., Modolo, R., Mousis, O., Nakamura, M., Neish, C. D., Nixon, C. A., Nna Mvondo, D., Orton, G., Paetzold, M., Pitman, J., Pogrebenko, S., Pollard, W., Prieto-Ballesteros, O., Rannou, P., Reh, K., Richter, L., Robb, F. T., Rodrigo, R., Rodriguez, S., Romani, P., Ruiz Bermejo, M., Sarris, E. T., Schenk, P., Schmitt, B., Schmitz, N., Schulze-Makuch, D., Schwingenschuh, K., Selig, A., Sicardy, B., Soderblom, L., Spilker, L. J., Stam, D., Steele, A., Stephan, K., Strobel, D. F., Szego, K., Szopa, C., Thissen, R., Tomasko, M. G., Toublanc, D., Vali, H., Vardavas, I., Vuitton, V., West, R. A., Yelle, R., Young, E. F., Mar. 2009. TandEM: Titan and Enceladus mission. *Experimental Astronomy* 23, 893–946.
- Coustenis, A., Salama, A., Lellouch, E., Encrenaz, T., Bjoraker, G. L., Samuelson, R. E., de Graauw, T., Feuchtgruber, H., Kessler, M. F., Aug. 1998. Evidence for water vapor in Titan's atmosphere from ISO/SWS data. *Astron. and Astrophys.* 336, L85–L89.
- Coustenis, A., Salama, A., Schulz, B., Ott, S., Lellouch, E., Encrenaz, T. H., Gautier, D., Feuchtgruber, H., Feb. 2003. Titan's atmosphere from ISO mid-infrared spectroscopy. *Icarus* 161, 383–403.
- Crida, A., Charnoz, S., Hsu, H.-W., Dones, L., 2019. Are Saturn's rings actually young? *Nature Astronomy* 3 (11), 967–970.
- Crósta, A., Silber, E., Lopes, R., Johnson, B., Bjonnes, E., Malaska, M., Vance, S., Sotin, C., Solomonidou, A., Soderblom, J., 2021. Modeling the formation of Menrva impact crater on Titan: Implications for habitability. *Icarus* 370, 114679.
- Ćuk, M., Dones, L., Nesvorný, D., 2016. Dynamical evidence for a late formation of Saturn's moons. *The Astrophysical Journal* 820 (2), 97.
- Czaplinski, E. C., Vu, T. H., Cable, M. L., Choukroun, M., Malaska, M. J., Hodyss, R., 2023. Experimental Characterization of the Pyridine: Acetylene Co-crystal and Implications for Titan's Surface. *ACS Earth and Space Chemistry* 7 (3), 597–608.

- Davies, A. G., Sotin, C., Choukroun, M., Matson, D. L., Johnson, T. V., 2016. Cryolava flow destabilization of crustal methane clathrate hydrate on Titan. *Icarus* 274, 23–32.
- de Kok, R. J., Teamby, N. A., Maltagliati, L., Irwin, P. G. J., Vinatier, S., Oct. 2014. HCN ice in Titan’s high-altitude southern polar cloud. *Nature* 514, 65–67.
- De Pater, I., Ádámkovics, M., Bouchez, A. H., Brown, M. E., Gibbard, S. G., Marchis, F., Roe, H. G., Schaller, E. L., Young, E., 2006. Titan imagery with Keck adaptive optics during and after probe entry. *Journal of Geophysical Research: Planets* 111 (E7).
- Del Genio, A. D., Zhou, W., Eichler, T. P., 1993. Equatorial superrotation in a slowly rotating GCM: Implications for Titan and Venus. *Icarus* 101 (1), 1–17.
- Dobrijevic, M., Hébrard, E., Loison, J., Hickson, K., Jan 2014. Coupling of oxygen, nitrogen, and hydrocarbon species in the photochemistry of Titan’s atmosphere. *Icarus* 228, 324–346.
URL <http://dx.doi.org/10.1016/j.icarus.2013.10.015>
- Dubinski, J., 2019. A recent origin for Saturn’s rings from the collisional disruption of an icy moon. *Icarus* 321, 291–306.
- Durante, D., Hemingway, D., Racioppa, P., Iess, L., Stevenson, D., 2019. Titan’s gravity field and interior structure after Cassini. *Icarus* 326, 123–132.
- Dutuit, O., Carrasco, N., Thissen, R., Vuitton, V., Alcaraz, C., Pernot, P., Balucani, N., Casavecchia, P., Canosa, A., Le Picard, S., et al., 2013. Critical review of N, N⁺, N₂⁺, N⁺⁺, and N₂⁺⁺ main production processes and reactions of relevance to Titan’s atmosphere. *The Astrophysical Journal Supplement Series* 204 (2), 20.
- Ennis, C., Cable, M. L., Hodyss, R., Maynard-Casely, H. E., jun 2020. Mixed hydrocarbon and cyanide ice compositions for Titan’s atmospheric aerosols: A ternary-phase co-crystal predicted by density functional theory. *ACS Earth and Space Chemistry* 4 (7), 1195–1200.
URL <https://doi.org/10.1021/acsearthspacechem.0c00130>

- Ewing, R. C., Hayes, A. G., Lucas, A., 2015. Sand dune patterns on Titan controlled by long-term climate cycles. *Nature Geoscience* 8 (1), 15–19.
- Fanson, J., Bernstein, R., Angeli, G., Ashby, D., Bigelow, B., Brossus, G., Bouchez, A., Burgett, W., Contos, A., Demers, R., et al., 2020. Overview and status of the Giant Magellan Telescope project. In: *Ground-based and airborne telescopes VIII*. Vol. 11445. SPIE, pp. 295–314.
- Fanson, J., Bernstein, R., Ashby, D., Bigelow, B., Brossus, G., Burgett, W., Demers, R., Fischer, B., Figueira, F., Groark, F., et al., 2022. Overview and status of the Giant Magellan Telescope project. In: *Ground-based and Airborne Telescopes IX*. Vol. 12182. SPIE, pp. 454–473.
- Fanson, J., McCarthy, P. J., Bernstein, R., Angeli, G., Ashby, D., Bigelow, B., Bouchez, A., Burgett, W., Chauvin, E., Contos, A., et al., 2018. Overview and status of the Giant Magellan Telescope project. In: *Ground-based and Airborne Telescopes VII*. Vol. 10700. SPIE, pp. 319–332.
- Faulk, S. P., Mitchell, J. L., Moon, S., Lora, J. M., oct 2017. Regional patterns of extreme precipitation on Titan consistent with observed alluvial fan distribution. *Nature Geoscience* 10 (11), 827–831.
URL <https://doi.org/10.1038/ngeo3043>
- Fortes, A., 2000. Exobiological implications of a possible ammonia–water ocean inside Titan. *Icarus* 146 (2), 444–452.
- Fortes, A. D., 2012. Titan’s internal structure and the evolutionary consequences. *Planetary and Space Science* 60 (1), 10–17.
- Fortes, A. D., Grindrod, P. M., Trickett, S. K., Vočadlo, L., May 2007. Ammonium sulfate on Titan: Possible origin and role in cryovolcanism. *Icarus* 188, 139–153.
- Francis, T. A., Maynard-Casely, H. E., Cable, M. L., Hodyss, R., Ennis, C., 2023. Simulation of Cocrystal Formation in Planetary Atmospheres: The C₆H₆: C₂H₂ Cocrystal Produced by Gas Deposition. *The Journal of Physical Chemistry A* 127 (10), 2322–2335.
- Friedson, A. J., West, R. A., Wilson, E. H., Oyafuso, F., Orton, G. S., 2009. A global climate model of Titan’s atmosphere and surface. *Planetary and Space Science* 57 (14-15), 1931–1949.

- Gans, B., Peng, Z., Carrasco, N., Gauyacq, D., Lebonnois, S., Pernot, P., 2013. Impact of a new wavelength-dependent representation of methane photolysis branching ratios on the modeling of Titan's atmospheric photochemistry. *Icarus* 223 (1), 330–343.
- Gautier, T., Carrasco, N., Schmitz-Afonso, I., Touboul, D., Szopa, C., Buch, A., Pernot, P., 2014. Nitrogen incorporation in Titan's tholins inferred by high resolution orbitrap mass spectrometry and gas chromatography–mass spectrometry. *Earth and Planetary Science Letters* 404, 33–42.
URL <https://www.sciencedirect.com/science/article/pii/S0012821X14004610>
- Gilmozzi, R., Spyromilio, J., 2007. The European Extremely Large Telescope (E-ELT). *The Messenger* 127 (11), 3.
- Glein, C. R., 2015. Noble gases, nitrogen, and methane from the deep interior to the atmosphere of Titan. *Icarus* 250, 570–586.
- Goossens, S., van Noort, B., Mateo, A., Mazarico, E., van der Wal, W., 2024. A low-density ocean inside Titan inferred from Cassini data. *Nature Astronomy*, 1–10.
- Grasset, O., Sotin, C., Deschamps, F., 2000. On the internal structure and dynamics of Titan. *Planetary and Space Science* 48 (7-8), 617–636.
- Graves, S., Mckay, C. P., Griffith, C. A., Ferri, F., Fulchignoni, M., 2008. Rain and hail can reach the surface of Titan. *Planetary and Space Science* 56 (3-4), 346–357.
- Griffith, C. A., Hall, J. L., Geballe, T. R., Oct. 2000. Detection of daily clouds on Titan. *Science* 290, 509–513.
- Griffith, C. A., Owen, T., Miller, G. A., Geballe, T., Oct. 1998. Transient clouds in Titan's lower atmosphere. *Nature* 395, 575–578.
- Griffith, C. A., Penteado, P., Rannou, P., Brown, R., Boudon, V., Baines, K. H., Clark, R., Drossart, P., Buratti, B., Nicholson, P., McKay, C. P., Coustenis, A., Negrao, A., Jaumann, R., Sep. 2006. Evidence for a Polar Ethane Cloud on Titan. *Science* 313, 1620–1622.

- Griffith, C. A., Zahnle, K., 1995. Influx of cometary volatiles to planetary moons: The atmospheres of 1000 possible Titans. *Journal of Geophysical Research* 100, 16907–16922.
- Grubisic, A., Trainer, M. G., Li, X., Brinckerhoff, W. B., van Amerom, F. H., Danell, R. M., Costa, J. T., Castillo, M., Kaplan, D., Zacny, K., 2021. Laser desorption mass spectrometry at Saturn's moon Titan. *International Journal of Mass Spectrometry* 470, 116707.
- Gu, H., Cui, J., Niu, D.-D., Wellbrock, A., Tseng, W.-L., Xu, X.-J., 2019. Monte Carlo calculations of the atmospheric sputtering yields on Titan. *Astronomy & Astrophysics* 623, A18.
- Hanel, R., Conrath, B., Flasar, F. M., Kunde, V., Maguire, W., Pearl, J. C., Pirraglia, J., Samuelson, R., Herath, L., Allison, M., Cruikshank, D. P., Gautier, D., Gierasch, P. J., Horn, L., Koppany, R., Ponnampерuma, C., Apr. 1981. Infrared observations of the Saturnian system from Voyager 1. *Science* 212, 192–200.
- Hayes, A., Aharonson, O., Callahan, P., Elachi, C., Gim, Y., Kirk, R., Lewis, K., Lopes, R., Lorenz, R., Lunine, J., Mitchell, K., Mitri, G., Stofan, E., Wall, S., May 2008. Hydrocarbon lakes on Titan: Distribution and interaction with a porous regolith. *Geophysical Research Letters* 35, 9204–+.
- Hayes, A. G., Jun 2016. The Lakes and Seas of Titan. *Annual Review of Earth and Planetary Sciences* 44 (1), 57–83.
URL <https://doi.org/10.1146/annurev-earth-060115-012247>
- Hayes, A. G., Birch, S., Dietrich, W. E., Howard, A. D., Kirk, R. L., Poggiali, V., Mastrogiovanni, M., Michaelides, R., Corlies, P., Moore, J., et al., 2017. Topographic constraints on the evolution and connectivity of Titan's lacustrine basins. *Geophysical Research Letters* 44 (23), 11–745.
- He, C., Smith, M. A., 2014. Identification of nitrogenous organic species in Titan aerosols analogs: Implication for prebiotic chemistry on Titan and early Earth. *Icarus* 238, 86–92.
- Hébrard, E., Bénilan, Y., Raulin, F., 2005. Sensitivity effects of photochemical parameters uncertainties on hydrocarbon production in the

- atmosphere of Titan. *Advances in Space Research* 36 (2), 268–273.
URL <https://www.sciencedirect.com/science/article/pii/S0273117705003856>
- Hébrard, E., Dobrijevic, M., Bénilan, Y., Raulin, F., 2006. Photochemical kinetics uncertainties in modeling Titan's atmosphere: A review. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews* 7 (4), 211–230.
URL <https://www.sciencedirect.com/science/article/pii/S138955670700007X>
- Hébrard, E., Dobrijevic, M., Bénilan, Y., Raulin, F., 2007. Photochemical kinetics uncertainties in modeling Titan's atmosphere: First consequences. *Planetary and Space Science* 55 (10), 1470–1489.
URL <https://www.sciencedirect.com/science/article/pii/S0032063307001213>
- Hébrard, E., Marty, B., 2014. Coupled noble gas–hydrocarbon evolution of the early Earth atmosphere upon solar UV irradiation. *Earth and Planetary Science Letters* 385, 40–48.
- Hilbert, D., 2000. Mathematical problems. *Bulletin of the American Mathematical Society* 37 (4), 407–436.
- Hirai, E., Sekine, Y., Zhang, N., Noda, N., Tan, S., Takahashi, Y., Kagi, H., 2023. Rapid aggregation and dissolution of organic aerosols in liquid methane on Titan. *Geophysical Research Letters* 50 (12), e2023GL103015.
- Hörst, S., Tolbert, M. A., 2013. In situ measurements of the size and density of Titan aerosol analogs. *The Astrophysical Journal Letters* 770 (1), L10.
- Hörst, S., Yelle, R., Buch, A., Carrasco, N., Cernogora, G., Dutuit, O., Quirico, E., Sciamma-O'Brien, E., Smith, M., Somogyi, Á., Szopa, C., Thissen, R., Vuitton, V., sep 2012. Formation of Amino Acids and Nucleotide Bases in a Titan Atmosphere Simulation Experiment. *Astrobiology* 12 (9), 809–817.
URL <https://doi.org/10.1089/ast.2011.0623>
- Hourdin, F., Talagrand, O., Sadourny, R., Courtin, R., Gautier, D., McKay, C. P., 1995. Numerical simulation of the general circulation of the atmosphere of Titan. *Icarus* 117 (2), 358–374.

- Howard, A. D., Breton, S., Moore, J. M., 2016. Formation of gravel pavements during fluvial erosion as an explanation for persistence of ancient cratered terrain on Titan and Mars. *Icarus* 270, 100–113.
- Hubbard, W. B., Hunten, D., Reitsema, H., Brosch, N., Nevo, Y., Carreira, E., Rossi, F., Wasserman, L., 1990. Results for Titan's atmosphere from its occultation of 28 Sagittarii. *Nature* 343 (6256), 353–355.
- Iess, L., Jacobson, R. A., Ducci, M., Stevenson, D. J., Lunine, J. I., Armstrong, J. W., Asmar, S. W., Racioppa, P., Rappaport, N. J., Tortora, P., 2012. The tides of Titan. *Science* 337 (6093), 457–459.
- Iess, L., Rappaport, N. J., Jacobson, R. A., Racioppa, P., Stevenson, D. J., Tortora, P., Armstrong, J. W., Asmar, S. W., Mar. 2010. Gravity Field, Shape, and Moment of Inertia of Titan. *Science* 327, 1367–.
- Israël, G., Szopa, C., Raulin, F., Cabane, M., Niemann, H. B., Atreya, S. K., Bauer, S. J., Brun, J.-F., Chassefière, E., Coll, P., et al., Nov 2005. Complex organic matter in Titan's atmospheric aerosols from in situ pyrolysis and analysis. *Nature* 438 (7069), 796–799.
 URL <http://dx.doi.org/10.1038/nature04349>
- Israël, G., Cabane, M., Brun, J.-F., Niemann, H., Way, S., Riedler, W., Steller, M., Raulin, F., Coscia, D., 2003. Huygens probe aerosol collector pyrolyser experiment. In: The Cassini-Huygens Mission: Overview, Objectives and Huygens Instrumentarium Volume 1. Springer, pp. 433–468.
- Jaumann, R., Brown, R. H., Stephan, K., Barnes, J. W., Soderblom, L. A., Sotin, C., Le Mouélic, S., Clark, R. N., Soderblom, J., Buratti, B. J., Wagner, R., McCord, T. B., Rodriguez, S., Baines, K. H., Cruikshank, D. P., Nicholson, P. D., Griffith, C. A., Langhans, M., Lorenz, R. D., Oct. 2008. Fluvial erosion and post-erosional processes on Titan. *Icarus* 197, 526–538.
- Jaumann, R., Kirk, R. L., Lorenz, R. D., Lopes, R. M., Stofan, E., Turtle, E. P., Keller, H. U., Wood, C. A., Sotin, C., Soderblom, L. A., et al., 2010. Geology and surface processes on Titan. In: Titan from Cassini-Huygens. Springer, pp. 75–140.
- Jennings, D. E., Nixon, C. A., Jolly, A., Bézard, B., Coustenis, A., Vinatier, S., Irwin, P. G. J., Teanby, N. A., Romani, P. N., Achterberg, R. K., Flasar,

- F. M., Jul. 2008. Isotopic Ratios in Titan's Atmosphere from Cassini CIRS Limb Sounding: HC₃N in the North. *Astrophys. J. Lett.* 681, L109–L111.
- Johnson, R. E., 1994. Plasma-induced sputtering of an atmosphere. *Space Science Reviews* 69 (3-4), 215–253.
- Johnson, R. E., 2009. Sputtering and heating of Titan's upper atmosphere. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 367 (1889), 753–771.
- Journaux, B., Daniel, I., Petitgirard, S., Cardon, H., Perrillat, J.-P., Caracas, R., Mezouar, M., 2017. Salt partitioning between water and high-pressure ices. Implication for the dynamics and habitability of icy moons and water-rich planetary bodies. *Earth and Planetary Science Letters* 463, 36–47.
- Journaux, B., Kalousová, K., Sotin, C., Tobie, G., Vance, S., Saur, J., Bol lengier, O., Noack, L., Rückriemen-Bez, T., Van Hoolst, T., et al., 2020. Large ocean worlds with high-pressure ices. *Space Science Reviews* 216, 1–36.
- Kaiser, R. I., Parker, D. S., Mebel, A. M., 2015. Reaction dynamics in astrochemistry: Low-temperature pathways to polycyclic aromatic hydrocarbons in the interstellar medium. *Annual Review of Physical Chemistry* 66, 43–67.
- Kalousová, K., Sotin, C., 2020a. Dynamics of Titan's high-pressure ice layer. *Earth and Planetary Science Letters* 545, 116416.
- Kalousová, K., Sotin, C., 2020b. The insulating effect of methane clathrate crust on Titan's thermal evolution. *Geophysical Research Letters* 47 (13), e2020GL087481.
- Kargel, J. S., 1991. Brine volcanism and the interior structures of asteroids and icy satellites. *Icarus* 94 (2), 368–390.
- Kargel, J. S., 1992. Ammonia-water volcanism on icy satellites: Phase relations at 1 atmosphere. *Icarus* 100 (2), 556–574.
- Kawai, J., Jagota, S., Kaneko, T., Obayashi, Y., Yoshimura, Y., Khare, B. N., Deamer, D. W., McKay, C. P., Kobayashi, K., 2013. Self-assembly of tholins in environments simulating Titan liquidospheres: implications

- for formation of primitive coacervates on Titan. International Journal of Astrobiology 12 (4), 282–291.
- Kawai, J., Kebukawa, Y., McKay, C. P., Kobayashi, K., 2019. Nucleic acid bases in Titan tholins and possible genetic systems in the Titan liquidosphere. Life Sciences in Space Research 20, 20–29.
- Kempf, S., Altobelli, N., Schmidt, J., Cuzzi, J. N., Estrada, P. R., Srama, R., 2023. Micrometeoroid infall onto Saturn's rings constrains their age to no more than a few hundred million years. Science Advances 9 (19), eadf8537.
- Khare, B. N., Sagan, C., Ogino, H., Nagy, B., Er, C., Schram, K. H., Arakawa, E. T., 1986. Amino acids derived from Titan Tholins. Icarus 68 (1), 176–184.
URL <https://www.sciencedirect.com/science/article/pii/0019103586900801>
- Kohlhase, C., Penzo, P. A., 1977. Voyager mission description. Space Science Reviews 21 (2), 77–101.
- Korycansky, D. G., Zahnle, K. J., Jan. 2011. Titan impacts and escape. Icarus 211, 707–721.
- Krasnopolsky, V. A., May 2009. A photochemical model of Titan's atmosphere and ionosphere. Icarus 201, 226–256.
- Krasnopolsky, V. A., Oct. 2010. The photochemical model of Titan's atmosphere and ionosphere: A version without hydrodynamic escape. Plan. & Space Sci. 58, 1507–1515.
- Krasnopolsky, V. A., 2016. Isotopic ratio of nitrogen on Titan: Photochemical interpretation. Planetary and Space Science 134, 61–63.
- Kress, M. E., McKay, C. P., Apr. 2004. Formation of methane in comet impacts: implications for Earth, Mars, and Titan. Icarus 168, 475–483.
- Kuiper, G. P., Nov. 1944. Titan: a Satellite with an Atmosphere. Astrophys. J. 100, 378–+.
- Kunde, V. G., Aikin, A. C., Hanel, R. A., Jennings, D. E., Maguire, W. C., Samuelson, R. E., Aug. 1981. C₄H₂, HC₃N and C₂N₂ in Titan's atmosphere. Nature 292, 686–688.

- Kutsop, N., Hayes, A., Corlies, P., Le Mouélic, S., Lunine, J., Nixon, C., Rannou, P., Rodriguez, S., Roman, M., Sotin, C., et al., 2022. Titan Stratospheric Haze Bands Observed in Cassini VIMS as Tracers of Meridional Circulation. *The Planetary Science Journal* 3 (5), 114.
- Lai, J. C.-Y., Cordner, M. A., Nixon, C. A., Achterberg, R. K., Molter, E. M., Teanby, N. A., Palmer, M. Y., Charnley, S. B., Lindberg, J. E., Kisiel, Z., et al., Nov 2017. Mapping Vinyl Cyanide and Other Nitriles in Titan's Atmosphere Using ALMA. *The Astronomical Journal* 154 (5), 206. URL <http://dx.doi.org/10.3847/1538-3881/aa8eef>
- Lainey, V., Jacobson, R. A., Tajeddine, R., Cooper, N. J., Murray, C., Robert, V., Tobie, G., Guillot, T., Mathis, S., Remus, F., et al., 2017. New constraints on Saturn's interior from Cassini astrometric data. *Icarus* 281, 286–296.
- Lalich, D., Poggiali, V., Hayes, A., Mastrogiovanni, M., Malaska, M., Schurmeier, L., 2022. Diverse evolution of mountains and hummocks on Titan as observed by the Cassini RADAR altimeter. *Icarus* 374, 114775.
- Lammer, H., Bauer, S., 1993. Atmospheric mass loss from Titan by sputtering. *Planetary and Space Science* 41 (9), 657–663.
- Lammer, H., Stumptner, W., Bauer, S., 1998. Dynamic escape of H from Titan as consequence of sputtering induced heating. *Planetary and Space Science* 46 (9-10), 1207–1213.
- Langhans, M., Jaumann, R., Stephan, K., Brown, R. H., Buratti, B., Clark, R. N., Baines, K. H., Nicholson, P. D., Lorenz, R. D., Soderblom, L. A., et al., 2012. Titan's fluvial valleys: Morphology, distribution, and spectral properties. *Planetary and Space Science* 60 (1), 34–51.
- Larson, E. J., Toon, O. B., West, R. A., Friedson, A. J., 2015. Microphysical modeling of Titan's detached haze layer in a 3D GCM. *Icarus* 254, 122–134.
- Larsson, R., McKay, C. P., 2013. Timescale for oceans in the past of Titan. *Planetary and Space Science* 78, 22–24.
- Lavvas, P., Coustenis, A., Vardavas, I., 2008a. Coupling photochemistry with haze formation in Titan's atmosphere, Part I: Model description. *Planetary and Space Science* 56 (1), 27 – 66.

URL <http://www.sciencedirect.com/science/article/B6V6T-4PN05S4-2/2f028ed3f317ff898f73c50d4b73fb21>

Lavvas, P., Sander, M., Kraft, M., Imanaka, H., Jan 2011. Surface Chemistry and Particle Shape: Processes for the Evolution of Aerosols in Titan's Atmosphere. *The Astrophysical Journal* 728 (2), 80.

URL <https://doi.org/10.1088/0004-637x/728/2/80>

Lavvas, P. P., Coustenis, A., Vardavas, I. M., 2008b. Coupling photochemistry with haze formation in Titan's atmosphere, Part II: Results and validation with Cassini/Huygens data. *Plan. & Space Sci.* 56 (1), 67 – 99.

URL <http://www.sciencedirect.com/science/article/B6V6T-4PN05S4-3/2/b3847e5b6c934fd53368891c25e62210>

Le Gall, A., Janssen, M. A., Paillou, P., Lorenz, R. D., Wall, S. D., Cassini Radar Team, Jun. 2010. Radar-bright channels on Titan. *Icarus* 207, 948–958.

Le Gall, A., Malaska, M. J., Lorenz, R. D., Janssen, M. A., Tokano, T., Hayes, A. G., Mastrogiovanni, M., Lunine, J. I., Veyssiére, G., Encrenaz, P., Karatekin, O., Feb 2016. Composition, seasonal change, and bathymetry of Ligeia Mare, Titan, derived from its microwave thermal emission. *Journal of Geophysical Research: Planets* 121 (2), 233–251.

URL <https://doi.org/10.1002/2015je004920>

Lebonnois, S., Burgalat, J., Rannou, P., Charnay, B., Mar 2012. Titan global climate model: A new 3-dimensional version of the IPSL Titan GCM. *Icarus* 218 (1), 707–722.

URL <http://dx.doi.org/10.1016/j.icarus.2011.11.032>

Lebreton, J.-P., 2002. The Huygens Probe: Science, Payload and Mission Overview. *Space Science Reviews* 104 (1/4), 59–100.

URL <https://doi.org/10.1023/a:1023657127549>

Lellouch, E., Bézard, B., Flasar, F. M., Vinatier, S., Achterberg, R., Nixon, C. A., Bjoraker, G. L., Gorius, N., Mar. 2014. The distribution of methane in Titan's stratosphere from Cassini/CIRS observations. *Icarus* 231, 323–337.

- Lellouch, E., Coustenis, A., Sebag, B., Cuby, J.-G., López-Valverde, M., Schmitt, B., Fouchet, T., Crovisier, J., Mar. 2003. Titan's 5- μ m window: observations with the Very Large Telescope. *Icarus* 162, 125–142.
- Lellouch, E., Gurwell, M. A., Moreno, R., Vinatier, S., Strobel, D. F., Moullet, A., Butler, B., Lara, L., Hidayat, T., Villard, E., apr 2019. An intense thermospheric jet on Titan. *Nature Astronomy* 3 (7), 614–619.
URL <https://doi.org/10.1038/s41550-019-0749-4>
- Lewis, J. S., 1971. Satellites of the outer planets: Their physical and chemical nature. *Icarus* 15 (2), 174–185.
- Lewis-Merrill, R. A., Moon, S., Mitchell, J. L., Lora, J. M., 2022. Assessing Environmental Factors of Alluvial Fan Formation on Titan. *The Planetary Science Journal* 3 (9), 223.
- Li, J., Yu, X., Sciamma-O'Brien, E., He, C., Sebree, J. A., Salama, F., Hörist, S. M., Zhang, X., 2022. A cross-laboratory comparison study of Titan haze analogs: Surface energy. *The Planetary Science Journal* 3 (1), 2.
- Litwin, K. L., Zygielbaum, B. R., Polito, P. J., Sklar, L. S., Collins, G. C., 2012. Influence of temperature, composition, and grain size on the tensile failure of water ice: Implications for erosion on Titan. *Journal of Geophysical Research: Planets* 117 (E8).
- Liu, Z. Y.-C., Radebaugh, J., Harris, R. A., Christiansen, E. H., Neish, C. D., Kirk, R. L., Lorenz, R. D., Team, C. R., et al., 2016a. The tectonics of Titan: Global structural mapping from Cassini RADAR. *Icarus* 270, 14–29.
- Liu, Z. Y.-C., Radebaugh, J., Harris, R. A., Christiansen, E. H., Rupper, S., 2016b. Role of fluids in the tectonic evolution of Titan. *Icarus* 270, 2–13.
- Loison, J., Hébrard, E., Dobrijevic, M., Hickson, K., Caralp, F., Hue, V., Gronoff, G., Venot, O., Bénilan, Y., Feb 2015. The neutral photochemistry of nitriles, amines and imines in the atmosphere of Titan. *Icarus* 247, 218–247.
URL <http://dx.doi.org/10.1016/j.icarus.2014.09.039>
- Lombardo, N. A., Nixon, C. A., Achterberg, R. K., Jolly, A., Sung, K., Irwin, P. G. J., Flasar, F. M., Jan 2019a. Spatial and seasonal variations

in C₃H_x hydrocarbon abundance in Titan's stratosphere from Cassini CIRS observations. *Icarus* 317, 454–469.

URL <http://dx.doi.org/10.1016/j.icarus.2018.08.027>

Lombardo, N. A., Nixon, C. A., Greathouse, T. K., Bézard, B., Jolly, A., Vinatier, S., Teanby, N. A., Richter, M. J., G Irwin, P. J., Coustenis, A., et al., Aug 2019b. Detection of Propadiene on Titan. *The Astrophysical Journal* 881 (2), L33.

URL <http://dx.doi.org/10.3847/2041-8213/ab3860>

Lopes, R., Stofan, E., Peckyno, R., Radebaugh, J., Mitchell, K., Mitri, G., Wood, C., Kirk, R., Wall, S., Lunine, J., et al., 2010. Distribution and interplay of geologic processes on Titan from Cassini radar data. *Icarus* 205 (2), 540–558.

Lopes, R., Wall, S., Elachi, C., Birch, S. P., Corlies, P., Coustenis, A., Hayes, A., Hofgartner, J., Janssen, M. A., Kirk, R., et al., 2019. Titan as revealed by the Cassini RADAR. *Space Science Reviews* 215, 1–50.

Lopes, R. M., 2017. Titan's 'Mount Doom'. *Nature Astronomy* 1 (9), 578–578.

Lopes, R. M., Malaska, M. J., Schoenfeld, A., Solomonidou, A., Birch, S., Florence, M., Hayes, A., Williams, D., Radebaugh, J., Verlander, T., et al., 2020. A global geomorphologic map of Saturn's moon Titan. *Nature astronomy* 4 (3), 228–233.

Lopes, R. M. C., Kirk, R. L., Mitchell, K. L., Legall, A., Barnes, J. W., Hayes, A., Kargel, J., Wye, L., Radebaugh, J., Stofan, E. R., Janssen, M. A., Neish, C. D., Wall, S. D., Wood, C. A., Lunine, J. I., Malaska, M. J., Mar. 2013. Cryovolcanism on Titan: New results from Cassini RADAR and VIMS. *Journal of Geophysical Research (Planets)* 118, 416–435.

Lopes, R. M. C., Mitchell, K. L., Stofan, E. R., Lunine, J. I., Lorenz, R., Paganelli, F., Kirk, R. L., Wood, C. A., Wall, S. D., Robshaw, L. E., Fortes, A. D., Neish, C. D., Radebaugh, J., Reffet, E., Ostro, S. J., Elachi, C., Allison, M. D., Anderson, Y., Boehmer, R., Boubin, G., Callahan, P., Encrenaz, P., Flamini, E., Francescetti, G., Gim, Y., Hamilton, G., Hensley, S., Janssen, M. A., Johnson, W. T. K., Kelleher, K., Muhleman, D. O., Ori, G., Orosei, R., Picardi, G., Posa, F., Roth, L. E., Seu, R.,

- Shaffer, S., Soderblom, L. A., Stiles, B., Vetrella, S., West, R. D., Wye, L., Zebker, H. A., Feb. 2007. Cryovolcanic features on Titan's surface as revealed by the Cassini Titan Radar Mapper. *Icarus* 186, 395–412.
- Lora, J. M., Battalio, J. M., Yap, M., Baciocco, C., 2022. Topographic and orbital forcing of Titan's hydroclimate. *Icarus* 384, 115095.
- Lora, J. M., Lunine, J. I., Russell, J. L., 2015. GCM simulations of Titan's middle and lower atmosphere and comparison to observations. *Icarus* 250, 516–528.
 URL <https://www.sciencedirect.com/science/article/pii/S0019103514007210>
- Lora, J. M., Tokano, T., Vatant d'Ollone, J., Lebonnois, S., Lorenz, R. D., 2019. A model intercomparison of Titan's climate and low-latitude environment. *Icarus* 333, 113–126.
 URL <https://www.sciencedirect.com/science/article/pii/S0019103518307838>
- Lorenz, R., 1995. Raindrops on Titan. *Advances in Space Research* 15 (3), 317–320.
 URL <https://www.sciencedirect.com/science/article/pii/S0273117799801033>
- Lorenz, R. D., 1998. Preliminary measurements of the cryogenic dielectric properties of water–ammonia ices: Implications for radar observations of icy satellites. *Icarus* 136 (2), 344–348.
- Lorenz, R. D., 2008. A review of balloon concepts for Titan. *Journal of the British Interplanetary Society* 61 (1), 2.
- Lorenz, R. D., 2014. The flushing of Ligeia: Composition variations across Titan's seas in a simple hydrological model. *Geophysical Research Letters* 41 (16), 5764–5770.
- Lorenz, R. D., Biolluz, G., Encrénaz, P., Janssen, M. A., West, R. D., Muöhleman, D. O., 2003. Cassini RADAR: prospects for Titan surface investigations using the microwave radiometer. *Planetary and Space Science* 51 (4-5), 353–364.

- Lorenz, R. D., Lopes, R. M., Paganelli, F., Lunine, J. I., Kirk, R. L., Mitchell, K. L., Soderblom, L. A., Stofan, E. R., Ori, G., Myers, M., et al., 2008. Fluvial channels on Titan: initial Cassini RADAR observations. *Planetary and Space Science* 56 (8), 1132–1144.
- Lorenz, R. D., Lunine, J. I., 1996. Erosion on Titan: Past and present. *Icarus* 122 (1), 79–91.
- Lorenz, R. D., Lunine, J. I., McKay, C. P., 1997. Titan under a red giant sun: A new kind of “habitable” moon. *Geophysical Research Letters* 24 (22), 2905–2908.
- Lorenz, R. D., MacKenzie, S. M., Neish, C. D., Le Gall, A., Turtle, E. P., Barnes, J. W., Trainer, M. G., Werynski, A., Hedgepeth, J., Karkoschka, E., 2021. Selection and characteristics of the Dragonfly landing site near Selk crater, Titan. *The Planetary Science Journal* 2 (1), 24.
- Lorenz, R. D., McKay, C. P., Lunine, J. I., 1997. Photochemically-induced collapse of Titan’s atmosphere. *Science* 275, 642–644.
- Lorenz, R. D., Mitchell, K. L., Kirk, R. L., Hayes, A. G., Aharonson, O., Zebker, H. A., Paillou, P., Radebaugh, J., Lunine, J. I., Janssen, M. A., Wall, S. D., Lopes, R. M., Stiles, B., Ostro, S., Mitri, G., Stofan, E. R., Jan. 2008. Titan’s inventory of organic surface materials. *Geophysical Research Letters* 35, 2206–+.
- Lorenz, R. D., Waite, Jr., J. H., 2008. Titan Explorer Flagship Mission Study. Tech. rep., Johns Hopkins University Applied Physics Laboratory.
- Lorenz, R. D., Wall, S., Radebaugh, J., Boubin, G., Reffet, E., Janssen, M., Stofan, E., Lopes, R., Kirk, R., Elachi, C., Lunine, J., Mitchell, K., Paganelli, F., Soderblom, L., Wood, C., Wye, L., Zebker, H., Anderson, Y., Ostro, S., Allison, M., Boehmer, R., Callahan, P., Encrenaz, P., Ori, G. G., Francescetti, G., Gim, Y., Hamilton, G., Hensley, S., Johnson, W., Kelleher, K., Muhleman, D., Picardi, G., Posa, F., Roth, L., Seu, R., Shaffer, S., Stiles, B., Vetrella, S., Flamini, E., West, R., May 2006. The Sand Seas of Titan: Cassini RADAR Observations of Longitudinal Dunes. *Science* 312, 724–727.
- Lucas, A., Rodriguez, S., Narteau, C., Charnay, B., Du Pont, S. C., Tokano, T., Garcia, A., Thiriet, M., Hayes, A. G., Lorenz, R. D., et al., 2014.

- Growth mechanisms and dune orientation on Titan. *Geophysical Research Letters* 41 (17), 6093–6100.
- Lunine, J. I., Stevenson, D. J., 1987. Clathrate and ammonia hydrates at high pressure: Application to the origin of methane on Titan. *Icarus* 70 (1), 61–77.
- Lutz, B. L., de Bergh, C., Owen, T., Jun. 1983. Titan - Discovery of carbon monoxide in its atmosphere. *Science* 220, 1374.
- MacKenzie, S. M., Runyon, K. D., Yu, X., Kok, J. F., Newman, C., Lorenz, R. D., Comola, F., 2023. Sediment-moving winds and abrasion on Titan: Implications for yardangs. *Icarus* 394, 115433.
- Maguire, W. C., Hanel, R. A., Jennings, D. E., Kunde, V. G., Samuelson, R. E., Aug. 1981. C₃H₈ and C₃H₄ in Titan's atmosphere. *Nature* 292, 683–686.
- Malaska, M. J., Hodyss, R., 2014. Dissolution of benzene, naphthalene, and biphenyl in a simulated Titan lake. *Icarus* 242, 74–81.
- Malaska, M. J., Lopes, R. M., Hayes, A. G., Radebaugh, J., Lorenz, R. D., Turtle, E. P., 2016. Material transport map of Titan: The fate of dunes. *Icarus* 270, 183–196.
- Malaska, M. J., Radebaugh, J., Lopes, R. M., Mitchell, K. L., Verlander, T., Schoenfeld, A. M., Florence, M. M., Le Gall, A., Solomonidou, A., Hayes, A. G., et al., 2020. Labyrinth terrain on Titan. *Icarus* 344, 113764.
- Mandt, K. E., Mousis, O., Lunine, J., Gautier, D., 2014. Protosolar ammonia as the unique source of Titan's nitrogen. *The Astrophysical Journal Letters* 788 (2), L24.
- Mandt, K. E., Waite, J. H., Lewis, W., Magee, B., Bell, J., Lunine, J., Mousis, O., Cordier, D., Dec. 2009. Isotopic evolution of the major constituents of Titan's atmosphere based on Cassini data. *Planetary and Space Science* 57, 1917–1930.
- Mandt, K. E., Waite, J. H., Teolis, B. D., Magee, B. A., Bell, J. M., Westlake, J. H., Nixon, C. A., Mousis, O., Lunine, J. I., 2012. The ¹²C/¹³C ratio on Titan from Cassini INMS measurements and implications for the evolution of Titan's methane. *Astrophys. J.* 749, 160.

- Marusiak, A. G., Vance, S., Panning, M. P., Bryant, A. S., Hesse, M. A., Carnahan, E., Journaux, B., 2022. The effects of methane clathrates on the thermal and seismic profile of Titan's icy lithosphere. *The Planetary Science Journal* 3 (7), 167.
- Mastrogiuseppe, M., Poggiali, V., Hayes, A., Lunine, J., Seu, R., Mitri, G., Lorenz, R., 2019. Deep and methane-rich lakes on Titan. *Nature Astronomy* 3 (6), 535–542.
- Matson, D. L., 2002. The Cassini/Huygens Mission to the Saturnian System. *Space Science Reviews* 104 (1/4), 1–58.
URL <http://dx.doi.org/10.1023/A:1023609211620>
- Matteoni, P., Mitri, G., Poggiali, V., Mastrogiuseppe, M., 2020. Geomorphological analysis of the southwestern margin of Xanadu, Titan: Insights on tectonics. *Journal of Geophysical Research: Planets* 125 (12), e2020JE006407.
- Maue, A. D., Levy, J. S., Burr, D. M., Matulka, P. R., Nathan, E., 2022. Rapid rounding of icy clasts during simulated fluvial transport in the Titan Tumbler. *Icarus* 375, 114831.
- Maynard-Casely, H. E., Hoddyss, R., Cable, M. L., Vu, T. H., Rahm, M., 2016. A co-crystal between benzene and ethane: a potential evaporite material for Saturn's moon Titan. *IUCrJ* 3 (3), 192–199.
- McKay, C., Coustenis, A., Samuelson, R., Lemmon, M., Lorenz, R., Cabane, M., Rannou, P., Drossart, P., 2001. Physical properties of the organic aerosols and clouds on Titan. *Planetary and Space Science* 49 (1), 79–99.
URL <https://www.sciencedirect.com/science/article/pii/S0032063300000519>
- McKay, C. P., Pollack, J. B., Courtin, R., Sep. 1991. The greenhouse and antigreenhouse effects on Titan. *Science* 253, 1118–1121.
- Miller, J., Birch, S., Hayes, A., Malaska, M., Lopes, R., Schoenfeld, A., Corlies, P., Burr, D., Farr, T., Perron, J., 2021. Fluvial features on Titan and Earth: Lessons from planform images in low-resolution SAR. *The Planetary Science Journal* 2 (4), 142.

- Miller, K. E., Glein, C. R., Waite, J. H., 2019. Contributions from accreted organics to Titan's atmosphere: new insights from cometary and chondritic data. *The Astrophysical Journal* 871 (1), 59.
- Mitchell, J. L., 2008. The drying of Titan's dunes: Titan's methane hydrology and its impact on atmospheric circulation. *Journal of Geophysical Research: Planets* 113 (E8).
- Mitchell, J. L., Ádámkovics, M., Caballero, R., Turtle, E. P., Sep. 2011. Locally enhanced precipitation organized by planetary-scale waves on Titan. *Nature Geoscience* 4, 589–592.
- Mitchell, J. L., Lora, J. M., 2016. The climate of Titan. *Annual Review of Earth and Planetary Sciences* 44, 353–380.
- Mitchell, J. L., Pierrehumbert, R. T., Frierson, D. M., Caballero, R., 2009. The impact of methane thermodynamics on seasonal convection and circulation in a model Titan atmosphere. *Icarus* 203 (1), 250–264.
- Mitri, G., Barnes, J., Coustenis, A., Flamini, E., Hayes, A., Lorenz, R. D., Mastrogiovanni, M., Orosei, R., Postberg, F., Reh, K., et al., 2021. Exploration of Enceladus and Titan: investigating ocean worlds' evolution and habitability in the Saturn system. *Experimental Astronomy*, 1–34.
- Mitri, G., Bland, M. T., Showman, A. P., Radebaugh, J., Stiles, B., Lopes, R. M. C., Lunine, J. I., Pappalardo, R. T., Oct. 2010. Mountains on Titan: Modeling and observations. *Journal of Geophysical Research (Planets)* 115, 10002–+.
- Mitri, G., Coustenis, A., Fanchini, G., Hayes, A. G., Iess, L., Khurana, K., Lebreton, J.-P., Lopes, R. M., Lorenz, R. D., Merigliola, R., et al., 2014a. The exploration of Titan with an orbiter and a lake probe. *Planetary and Space Science* 104, 78–92.
- Mitri, G., Merigliola, R., Hayes, A., Lefevre, A., Tobie, G., Genova, A., Lunine, J. I., Zebker, H., 2014b. Shape, topography, gravity anomalies and tidal deformation of Titan. *Icarus* 236, 169–177.
- Mitri, G., Showman, A. P., 2008. Thermal convection in ice-I shells of Titan and Enceladus. *Icarus* 193 (2), 387–396.

- Mitri, G., Showman, A. P., Lunine, J. I., Lopes, R. M., 2008. Resurfacing of Titan by ammonia-water cryomagma. *Icarus* 196 (1), 216–224.
- Mitri, G., Showman, A. P., Lunine, J. I., Lorenz, R. D., 2007. Hydrocarbon lakes on Titan. *Icarus* 186 (2), 385–394.
- Molter, E. M., Nixon, C. A., Cordiner, M. A., Serigano, J., Irwin, P. G. J., Teanby, N. A., Charnley, S. B., Lindberg, J. E., Aug 2016. ALMA Observations of HCN and its Isotopologues on Titan. *Astron. J.* 152 (2), 42. URL <http://dx.doi.org/10.3847/0004-6256/152/2/42>
- Moore, J. M., Howard, A. D., Morgan, A. M., 2014. The landscape of Titan as witness to its climate evolution. *Journal of Geophysical Research: Planets* 119 (9), 2060–2077.
- Moore, J. M., Pappalardo, R. T., Apr. 2011. Titan: An exogenic world? *Icarus* 212, 790–806.
- Mousis, O., Chassefiere, E., Holm, N. G., Bouquet, A., Waite, J. H., Geppert, W. D., Picaud, S., Aikawa, Y., Ali-Dib, M., Charlou, J.-L., et al., 2015. Methane clathrates in the solar system. *Astrobiology* 15 (4), 308–326.
- Mousis, O., Choukroun, M., Lunine, J. I., Sotin, C., 2014. Equilibrium composition between liquid and clathrate reservoirs on Titan. *Icarus* 239, 39–45.
- Mousis, O., Gautier, D., Bockelée-Morvan, D., Mar. 2002a. An Evolutionary Turbulent Model of Saturn’s Subnebula: Implications for the Origin of the Atmosphere of Titan. *Icarus* 156, 162–175.
- Mousis, O., Gautier, D., Coustenis, A., Sep. 2002b. The D/H Ratio in Methane in Titan: Origin and History. *Icarus* 159, 156–165.
- Mousis, O., Lunine, J. I., Picaud, S., Cordier, D., Waite, J. H., Mandt, K. E., 2011. Removal of Titan’s atmospheric Noble gases by their sequestration in surface clathrates. *The Astrophysical Journal Letters* 740 (1), L9.
- Muhleman, D. O., Grossman, A. W., Butler, B. J., Slade, M. A., may 1990. Radar Reflectivity of Titan. *Science* 248 (4958), 975–980. URL <https://doi.org/10.1126/science.248.4958.975>

- Neish, C., Malaska, M. J., Sotin, C., Lopes, R. M., Nixon, C. A., Affholder, A., Chatain, A., Cockell, C., Farnsworth, K. K., Higgins, P. M., et al., 2024. Organic Input to Titan's Subsurface Ocean Through Impact Cratering. *Astrobiology*.
- Neish, C., Somogyi, A., Imanaka, H., Lunine, J., Smith, M., 2008. Rate measurements of the hydrolysis of complex organic macromolecules in cold aqueous solutions: implications for prebiotic chemistry on the early Earth and Titan. *Astrobiology* 8 (2), 273–287.
- Neish, C. D., Lorenz, R. D., 2012. Titan's global crater population: a new assessment. *Plan. & Space Sci.* 60 (1), 26–33.
- Neish, C. D., Molaro, J., Lora, J., Howard, A., Kirk, R. L., Schenk, P., Bray, V., Lorenz, R., 2016. Fluvial erosion as a mechanism for crater modification on Titan. *Icarus* 270, 114–129.
- Neish, C. D., Somogyi, Á., Lunine, J. I., Smith, M. A., 2009. Low temperature hydrolysis of laboratory tholins in ammonia-water solutions: Implications for prebiotic chemistry on Titan. *Icarus* 201 (1), 412–421.
- Neish, C. D., Somogyi, A., Smith, M. A., 2010. Titan's primordial soup: formation of amino acids via low-temperature hydrolysis of tholins. *Astrobiology* 10 (3), 337–347.
- Nelson, J., Sanders, G. H., 2008. The status of the Thirty Meter Telescope project. In: *Ground-based and Airborne Telescopes II*. Vol. 7012. SPIE, pp. 504–521.
- Néri, A., Guyot, F., Reynard, B., Sotin, C., 2020. A carbonaceous chondrite and cometary origin for icy moons of Jupiter and Saturn. *Earth and Planetary Science Letters* 530, 115920.
- Nesvorný, D., Dones, L., De Prá, M., Womack, M., Zahnle, K. J., 2023. Impact rates in the outer solar system. *The Planetary Science Journal* 4 (8), 139.
- Neveu, M., Rhoden, A. R., 2019. Evolution of Saturn's mid-sized moons. *Nature Astronomy* 3 (6), 543–552.

Niemann, H. B., Atreya, S. K., Bauer, S. J., Biemann, K., Block, B., Carignan, G. R., Donahue, T. M., Frost, R. L., Gautier, D., Haberman, J. A., Harpold, D. N., Hunten, D. M., Israel, G., Lunine, J. I., Mauersberger, K., Owen, T. C., Raulin, F., Richards, J. E., Way, S. H., 2002. The Gas Chromatograph Mass Spectrometer for the Huygens Probe. *Space Science Reviews* 104 (1/4), 553–591.

URL <http://dx.doi.org/10.1023/A:1023680305259>

Niemann, H. B., Atreya, S. K., Bauer, S. J., Carignan, G. R., Demick, J. E., Frost, R. L., Gautier, D., Haberman, J. A., Harpold, D. N., Hunten, D. M., Israel, G., Lunine, J. I., Kasprzak, W. T., Owen, T. C., Paulkovich, M., Raulin, F., Raaen, E., Way, S. H., Dec. 2005. The abundances of constituents of Titan's atmosphere from the GCMS instrument on the Huygens probe. *Nature* 438, 779–784.

Niemann, H. B., Atreya, S. K., Demick, J. E., Gautier, D., Haberman, J. A., Harpold, D. N., Kasprzak, W. T., Lunine, J. I., Owen, T. C., Raulin, F., Dec. 2010. Composition of Titan's lower atmosphere and simple surface volatiles as measured by the Cassini-Huygens probe gas chromatograph mass spectrometer experiment. *Journal of Geophysical Research (Planets)* 115, 12006–+.

Nixon, C., Lorenz, R., Achterberg, R., Buch, A., Coll, P., Clark, R., Courtin, R., Hayes, A., Iess, L., Johnson, R., et al., Jun 2018. Titan's cold case files - Outstanding questions after Cassini-Huygens. *Planetary and Space Science* 155, 50–72.

URL <http://dx.doi.org/10.1016/j.pss.2018.02.009>

Nixon, C. A., Achterberg, R. K., Ádámkovics, M., Bézard, B., Bjoraker, G. L., Cornet, T., Hayes, A. G., Lellouch, E., Lemmon, M. T., López-Puertas, M., Rodriguez, S., Sotin, C., Teanby, N. A., Turtle, E. P., West, R. A., jan 2016a. Titan Science with the James Webb Space Telescope. *Publications of the Astronomical Society of the Pacific* 128 (959), 018007. URL <https://doi.org/10.1088/1538-3873/128/959/018007>

Nixon, C. A., Aslam, S., Guzewich, S., Esper, J., Thompson, S. N., Hewagama, T., Tran, D., Nehmetallah, G., 2019. TEAM-Titan Exploration Atmospheric Microprobes. In: 2019 IEEE Aerospace Conference. IEEE, pp. 1–16.

- Nixon, C. A., Kirchman, F., Esper, J., Folta, D., Mashiku, A., 2016b. Aero-capture design study for a Titan polar orbiter. In: 2016 IEEE Aerospace Conference. IEEE, pp. 1–16.
- Nixon, C. A., Teanby, N. A., Irwin, P., Hörst, S. M., May 2013. Upper limits for PH₃ and H₂S in Titan's atmosphere from Cassini CIRS. *Icarus* 224 (1), 253–256.
URL <http://dx.doi.org/10.1016/j.icarus.2013.02.024>
- Nixon, C. A., Temelso, B., Vinatier, S., Teanby, N. A., Bézard, B., Achterberg, R. K., Mandt, K. E., Sherrill, C. D., Irwin, P. G. J., Jennings, D. E., Romani, P. N., Coustenis, A., Flasar, F. M., 2012. Isotopic ratios in Titan's methane: measurements and modeling. *Astrophys. J.* 749, 159.
- Nixon, C. A., Thelen, A. E., Cordiner, M. A., Kisiel, Z., Charnley, S. B., Molter, E. M., Serigano, J., Irwin, P. G. J., Teanby, N. A., Kuan, Y.-J., oct 2020. Detection of Cyclopropenylidene on Titan with ALMA. *The Astronomical Journal* 160 (5), 205.
URL <https://doi.org/10.3847/1538-3881/abb679>
- Oleson, S. R., Lorenz, R., Paul, M., 2015. Titan submarine: exploring the depths of Kraken Mare. In: AIAA Space 2015 Conference and Exposition. p. 4445.
- Owen, T., Niemann, H., 2009. The origin of Titan's atmosphere: some recent advances. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 367 (1889), 607–615.
- Pailou, P., Lunine, J., Ruffié, G., Encrénaz, P., Wall, S., Lorenz, R., Janssen, M., 2008. Microwave dielectric constant of Titan-relevant materials. *Geophysical Research Letters* 35 (18).
- Palmer, M. Y., Cordiner, M. A., Nixon, C. A., Charnley, S. B., Teanby, N. A., Kisiel, Z., Irwin, P. G. J., Mumma, M. J., Jul 2017. ALMA Detection and Astrobiological Potential of Vinyl Cyanide on Titan. *Science Advances* 3 (7), e1700022.
URL <http://dx.doi.org/10.1126/sciadv.1700022>
- Pasek, M. A., Mousis, O., Lunine, J. I., Apr. 2011. Phosphorus chemistry on Titan. *Icarus* 212, 751–761.

- Penteado, P. F., Griffith, C. A., 2010. Ground-based measurements of the methane distribution on Titan. *Icarus* 206 (1), 345–351.
- Penteado, P. F., Griffith, C. A., Greathouse, T. K., de Bergh, C., Aug. 2005. Measurements of CH₃D and CH₄ in Titan from Infrared Spectroscopy. *The Astrophysical Journal Letters* 629, L53–L56.
- Perrin, Z., Carrasco, N., Chatain, A., Jovanovic, L., Vettier, L., Ruscassier, N., Cernogora, G., 2021. An atmospheric origin for HCN-derived polymers on Titan. *Processes* 9 (6), 965.
- Perron, J. T., Lamb, M. P., Koven, C. D., Fung, I. Y., Yager, E., Ádámkovics, M., 2006. Valley formation and methane precipitation rates on Titan. *Journal of Geophysical Research: Planets* 111 (E11).
- Pinto, J. P., Lunine, J. I., Kim, S., Yung, Y. L., Jan. 1986. D to H ratio and the origin and evolution of Titan's atmosphere. *Nature* 319, 388–390.
- Poch, O., Coll, P., Buch, A., Ramírez, S., Raulin, F., 2012. Production yields of organics of astrobiological interest from H₂O–NH₃ hydrolysis of Titan's tholins. *Planetary and Space Science* 61 (1), 114–123.
- Postberg, F., Sekine, Y., Klenner, F., Glein, C. R., Zou, Z., Abel, B., Furuya, K., Hillier, J. K., Khawaja, N., Kempf, S., et al., 2023. Detection of phosphates originating from Enceladus's ocean. *Nature* 618 (7965), 489–493.
- Radebaugh, J., Lorenz, R., Farr, T., Paillou, P., Savage, C., Spencer, C., 2010. Linear dunes on Titan and Earth: Initial remote sensing comparisons. *Geomorphology* 121 (1-2), 122–132.
- Radebaugh, J., Lorenz, R. D., Kirk, R. L., Lunine, J. I., Stofan, E. R., Lopes, R. M., Wall, S. D., Team, C. R., et al., 2007. Mountains on Titan observed by Cassini RADAR. *Icarus* 192 (1), 77–91.
- Radebaugh, J., Ventra, D., Lorenz, R. D., Farr, T., Kirk, R., Hayes, A., Malaska, M. J., Birch, S., Liu, Z. Y.-C., Lunine, J., et al., 2018. Alluvial and fluvial fans on Saturn's moon Titan reveal processes, materials and regional geology. *Geological Society, London, Special Publications* 440 (1), 281–305.

- Rafkin, S. C., Soto, A., 2020. Air-sea interactions on Titan: Lake evaporation, atmospheric circulation, and cloud formation. *Icarus* 351, 113903.
- Rahm, M., Lunine, J., Usher, D., Shalloway, D., 2016. Polymorphism and electronic structure of polyimine and its potential significance for prebiotic chemistry on Titan. *PNAS; Proceedings of the National Academy of Sciences* 113 (29), 8121–8126.
- Rannou, P., Cabane, M., Chassefiere, E., Botet, R., McKay, C., Courtin, R., 1995. Titan's Geometric Albedo: Role of the Fractal Structure of the Aerosols. *Icarus* 118 (2), 355–372.
URL <https://www.sciencedirect.com/science/article/pii/S0019103585711967>
- Rannou, P., McKay, C., Botet, R., Cabane, M., 1999. Semi-empirical model of absorption and scattering by isotropic fractal aggregates of spheres. *Planetary and Space Science* 47 (3-4), 385–396.
- Rannou, P., McKay, C., Lorenz, R., 2003. A model of Titan's haze of fractal aerosols constrained by multiple observations. *Planetary and Space Science* 51 (14-15), 963–976.
- Raulin, F., Brassé, C., Poch, O., Coll, P., 2012. Prebiotic-like chemistry on Titan. *Chemical Society Reviews* 41 (16), 5380–5393.
- Rodriguez, S., Garcia, A., Lucas, A., Appéré, T., Le Gall, A., Reffet, E., Le Corre, L., Le Mouélic, S., Cornet, T., Courrech du Pont, S., Narteau, C., Bourgeois, O., Radebaugh, J., Arnold, K., Barnes, J. W., Stephan, K., Jaumann, R., Sotin, C., Brown, R. H., Lorenz, R. D., Turtle, E. P., Feb. 2014. Global mapping and characterization of Titan's dune fields with Cassini: Correlation between RADAR and VIMS observations. *Icarus* 230, 168–179.
- Rodriguez, S., Le Mouélic, S., Rannou, P., Sotin, C., Brown, R. H., Barnes, J. W., Griffith, C. A., Burgalat, J., Baines, K. H., Buratti, B. J., Clark, R. N., Nicholson, P. D., Nov. 2011. Titan's cloud seasonal activity from winter to spring with Cassini/VIMS. *Icarus* 216, 89–110.
- Rodriguez, S., Le Mouélic, S., Rannou, P., Tobie, G., Baines, K. H., Barnes, J. W., Griffith, C. A., Hirtzig, M., Pitman, K. M., Sotin, C., Brown, R. H.,

- Buratti, B. J., Clark, R. N., Nicholson, P. D., Jun. 2009. Global circulation as the main source of cloud activity on Titan. *Nature* 459, 678–682.
- Rodriguez, S., Vinatier, S., Cordier, D., Tobie, G., Achterberg, R. K., Anderson, C. M., Badman, S. V., Barnes, J. W., Barth, E. L., Bézard, B., Carrasco, N., Charnay, B., Clark, R. N., Coll, P., Cornet, T., Coustenis, A., Couturier-Tamburelli, I., Dobrijevic, M., Flasar, F. M., de Kok, R., Freissinet, C., Galand, M., Gautier, T., Geppert, W. D., Griffith, C. A., Gudipati, M. S., Hadid, L. Z., Hayes, A. G., Hendrix, A. R., Jaumann, R., Jennings, D. E., Jolly, A., Kalousova, K., Koskinen, T. T., Lavvas, P., Lebonnois, S., Lebreton, J.-P., Gall, A. L., Lellouch, E., Mouélic, S. L., Lopes, R. M. C., Lora, J. M., Lorenz, R. D., Lucas, A., MacKenzie, S., Malaska, M. J., Mandt, K., Mastrogiovanni, M., Newman, C. E., Nixon, C. A., Radebaugh, J., Rafkin, S. C., Rannou, P., Sciamma-O'Brien, E. M., Soderblom, J. M., Solomonidou, A., Sotin, C., Stephan, K., Strobel, D., Szopa, C., Teanby, N. A., Turtle, E. P., Vuitton, V., West, R. A., jan 2022. Science goals and new mission concepts for future exploration of Titan's atmosphere, geology and habitability: Titan POlar scout/orbitEr and in situ lake lander and DrONe explorer (POSEIDON). *Experimental Astronomy*.
- URL <https://doi.org/10.1007/s10686-021-09815-8>
- Roe, H., Bouchez, A., Trujillo, C., Schaller, E., Brown, M., 2004. Discovery of temperate latitude clouds on Titan. *The Astrophysical Journal* 618 (1), L49.
- Roe, H., De Pater, I., Macintosh, B., McKay, C., 2002. Titan's Clouds from Gemini and Keck Adaptive Optics Imaging. *The Astrophysical Journal* 581 (2), 1399.
- Roe, H. G., Greathouse, T., Richter, M., Lacy, J., 2003. Propane on Titan. *The Astrophysical Journal* 597 (1), L65.
- Roman, M. T., West, R. A., Banfield, D. J., Gierasch, P. J., Achterberg, R. K., Nixon, C. A., Thomas, P. C., 2009. Determining a tilt in Titan's north-south albedo asymmetry from Cassini images. *Icarus* 203 (1), 242–249.
- Rubin, D. M., Hesp, P. A., 2009. Multiple origins of linear dunes on Earth and Titan. *Nature Geoscience* 2 (9), 653–658.

- Sanders, G. H., 2013. The Thirty Meter Telescope (TMT): An international observatory. *Journal of Astrophysics and Astronomy* 34, 81–86.
- Savage, C. J., Radabaugh, J., Christiansen, E. H., Lorenz, R. D., 2014. Implications of dune pattern analysis for Titan's surface history. *Icarus* 230, 180–190.
- Schaller, E., Roe, H., Schneider, T., Brown, M., 2009. Storms in the tropics of Titan. *Nature* 460 (7257), 873–875.
- Schaller, E. L., Brown, M. E., Roe, H. G., Bouchez, A. H., May 2006a. A large cloud outburst at Titan's south pole. *Icarus* 182, 224–229.
- Schaller, E. L., Brown, M. E., Roe, H. G., Bouchez, A. H., Trujillo, C. A., Oct. 2006b. Dissipation of Titan's south polar clouds. *Icarus* 184, 517–523.
- Schneider, T., Graves, S. D. B., Schaller, E. L., Brown, M. E., Jan. 2012. Polar methane accumulation and rainstorms on Titan from simulations of the methane cycle. *Nature* 481, 58–61.
- Schulz, F., Maillard, J., Kaiser, K., Schmitz-Afonso, I., Gautier, T., Afonso, C., Carrasco, N., Gross, L., Feb 2021. Imaging Titan's Organic Haze at Atomic Scale. *The Astrophysical Journal Letters* 908 (1), L13.
URL <https://doi.org/10.3847/2041-8213/abd93e>
- Schurmeier, L. R., Dombard, A. J., Malaska, M. J., Fagents, S. A., Radabaugh, J., Lalich, D. E., 2023. An intrusive cryomagmatic origin for northern radial labyrinth terrains on Titan and implications for the presence of crustal clathrates. *Icarus*, 115664.
- Serigano, J., Nixon, C. A., Cordiner, M. A., Irwin, P. G. J., Teanby, N. A., Charnley, S. B., Lindberg, J. E., Apr 2016. Isotopic Ratios Of Carbon And Oxygen In Titan's CO Using ALMA. *The Astrophysical Journal* 821 (1), L8.
URL <http://dx.doi.org/10.3847/2041-8205/821/1/L8>
- Sharkey, J., Teanby, N. A., Sylvestre, M., Mitchell, D. M., Seviour, W. J., Nixon, C. A., Irwin, P. G., 2020. Mapping the zonal structure of Titan's northern polar vortex. *Icarus* 337, 113441.

- Shematovich, V., Johnson, R., Michael, M., Luhmann, J., 2003. Nitrogen loss from Titan. *Journal of Geophysical Research: Planets* 108 (E8).
- Sicardy, B., Brahic, A., Ferrari, C., Gautier, D., Lecacheux, J., Lellouch, E., Reques, F., Arlot, J., Colas, F., Thuillot, W., et al., 1990. Probing Titan's atmosphere by stellar occultation. *Nature* 343 (6256), 350–353.
- Sicardy, B., Colas, F., Widemann, T., Bellucci, A., Beisker, W., Kretlow, M., Ferri, F., Lacour, S., Lecacheux, J., Lellouch, E., et al., 2006. The two Titan stellar occultations of 14 November 2003. *Journal of Geophysical Research: Planets* 111 (E11).
- Sicardy, B., Ferri, F., Roques, F., Lecacheux, J., Pau, S., Brosch, N., Nevo, Y., Hubbard, W., Reitsema, H., Blanco, C., et al., 1999. The structure of Titan's stratosphere from the 28 Sgr occultation. *Icarus* 142 (2), 357–390.
- Skidmore, W., et al., 2015. Thirty Meter Telescope detailed science case: 2015. *Research in Astronomy and Astrophysics* 15 (12), 1945.
- Smith, P. H., Lemmon, M. T., Lorenz, R. D., Sromovsky, L. A., Caldwell, J. J., Allison, M. D., Feb. 1996. Titan's Surface, Revealed by HST Imaging. *Icarus* 119, 336–349.
- Snowden, D., Higgins, A., 2021. A Monte Carlo model of energy deposition, ionization, and sputtering due to thermal ion precipitation into Titan's upper atmosphere. *Icarus* 354, 113929.
- Soderblom, L. A., Tomasko, M. G., Archinal, B. A., Becker, T. L., Bushroe, M. W., Cook, D. A., Doose, L. R., Galuszka, D. M., Hare, T. M., Howington-Kraus, E., et al., 2007. Topography and geomorphology of the Huygens landing site on Titan. *Planetary and Space Science* 55 (13), 2015–2024.
- Soderlund, K. M., 2019. Ocean dynamics of outer solar system satellites. *Geophysical Research Letters* 46 (15), 8700–8710.
- Soderlund, K. M., Rovira-Navarro, M., Le Bars, M., Schmidt, B. E., Gerkema, T., 2023. The physical oceanography of ice-covered moons. *Annual Review of Marine Science* 16.

- Sohl, F., Solomonidou, A., Wagner, F., Coustenis, A., Hussmann, H., Schulze-Makuch, D., 2014. Structural and tidal models of Titan and inferences on cryovolcanism. *Journal of Geophysical Research: Planets* 119 (5), 1013–1036.
- Solomonidou, A., Bampasidis, G., Hirtzig, M., Coustenis, A., Kyriakopoulos, K., Seymour, K. S., Bratsolis, E., Moussas, X., 2013. Morphotectonic features on Titan and their possible origin. *Planetary and Space Science* 77, 104–117.
- Solomonidou, A., Coustenis, A., Hirtzig, M., Rodriguez, S., Stephan, K., Lopes, R., Drossart, P., Sotin, C., Le Mouélic, S., Lawrence, K., et al., 2016. Temporal variations of Titan's surface with Cassini/VIMS. *Icarus* 270, 85–99.
- Solomonidou, A., Coustenis, A., Lopes, R. M., Malaska, M. J., Rodriguez, S., Drossart, P., Elachi, C., Schmitt, B., Philippe, S., Janssen, M., et al., 2018. The spectral nature of Titan's major geomorphological units: constraints on surface composition. *Journal of Geophysical Research: Planets* 123 (2), 489–507.
- Solomonidou, A., Le Gall, A., Malaska, M., Birch, S., Lopes, R., Coustenis, A., Rodriguez, S., Wall, S., Michaelides, R., Nasr, M., et al., 2020. Spectral and emissivity analysis of the raised ramparts around Titan's northern lakes. *Icarus* 344, 113338.
- Sotin, C., Jaumann, R., Buratti, B. J., Brown, R. H., Clark, R. N., Soderblom, L. A., Baines, K. H., Bellucci, G., Bibring, J., Capaccioni, F., Cerroni, P., Combes, M., Coradini, A., Cruikshank, D. P., Drossart, P., Formisano, V., Langevin, Y., Matson, D. L., McCord, T. B., Nelson, R. M., Nicholson, P. D., Sicardy, B., Le Mouelic, S., Rodriguez, S., Stephan, K., Scholz, C. K., Jun. 2005. Release of volatiles from a possible cryovolcano from near-infrared imaging of Titan. *Nature* 435, 786–789.
- Sotin, C., Lawrence, K. J., Reinhardt, B., Barnes, J. W., Brown, R. H., Hayes, A. G., Le Mouélic, S., Rodriguez, S., Soderblom, J. M., Soderblom, L. A., Baines, K. H., Buratti, B. J., Clark, R. N., Jaumann, R., Nicholson, P. D., Stephan, K., Nov. 2012. Observations of Titan's Northern lakes at 5 μm: Implications for the organic cycle and geology. *Icarus* 221, 768–786.

- Spiga, A., Lefèvre, M., d'Ollone, J. V., Lebonnois, S., Rodriguez, S., Chatain, A., Bonnefoy, L., Lucas, A., 2020. Developping a new model for mesoscale and large-eddy simulations of Titan's atmosphere. Tech. rep., Copernicus Meetings.
- Stevenson, J. M., Fouad, W. A., Shalloway, D., Usher, D., Lunine, J., Chapman, W. G., Clancy, P., 2015. Solvation of nitrogen compounds in Titan's seas, precipitates, and atmosphere. *Icarus* 256, 1–12.
 URL <https://www.sciencedirect.com/science/article/pii/S0019103515001621>
- Stofan, E., Lorenz, R., Lunine, J., Bierhaus, E. B., Clark, B., Mahaffy, P. R., Ravine, M., 2013. TiME—the Titan Mare Explorer. In: 2013 IEEE Aerospace Conference. IEEE, pp. 1–10.
- Strobel, D., 1982. Chemistry and evolution of Titan's atmosphere. *Planetary and Space Science* 30 (8), 839–848.
 URL <https://www.sciencedirect.com/science/article/pii/0032063382901167>
- Strobel, D. F., 1974. The photochemistry of hydrocarbons in the atmosphere of Titan. *Icarus* 21 (4), 466–470.
 URL <https://www.sciencedirect.com/science/article/pii/0019103574901493>
- Sulaiman, A. H., Achilleos, N., Bertucci, C., Coates, A., Dougherty, M., Hadid, L., Holmberg, M., Hsu, H.-W., Kimura, T., Kurth, W., et al., 2021. Enceladus and Titan: emerging worlds of the Solar System. *Experimental Astronomy*, 1–28.
- Sung, K., Steffens, B., Toon, G. C., Nemchick, D. J., Smith, M. A. H., 2020. Pseudoline parameters to represent n-butane ($n\text{-C}_4\text{H}_{10}$) cross-sections measured in the 7–15 μm region for the Titan atmosphere. *Journal of Quantitative Spectroscopy and Radiative Transfer* 251, 107011.
- Sung, K., Toon, G. C., Drouin, B. J., Mantz, A. W., Smith, M. A. H., 2018. FT-IR measurements of cold propene (C_3H_6) cross-sections at temperatures between 150 and 299 K. *Journal of Quantitative Spectroscopy and Radiative Transfer* 213, 119–132.

- Sung, K., Toon, G. C., Mantz, A. W., Smith, M. A. H., 2013. FT-IR measurements of cold C_3H_8 cross sections at 7—15 μm for Titan atmosphere. *Icarus*.
- Tamai, R., Spyromilio, J., 2014. European Extremely Large Telescope: progress report. In: *Ground-based and Airborne Telescopes V*. Vol. 9145. SPIE, pp. 510–518.
- Tan, S. P., Kargel, J. S., Marion, G. M., 2013. Titan's atmosphere and surface liquid: New calculation using Statistical Associating Fluid Theory. *Icarus* 222 (1), 53–72.
- Teanby, N. A., Cordiner, M. A., Nixon, C. A., Irwin, P. G. J., Hörst, S. M., Sylvestre, M., Serigano, J., Thelen, A. E., Richards, A. M. S., Charnley, S. B., May 2018. The Origin of Titan's External Oxygen: Further Constraints from ALMA Upper Limits on CS and CH_2NH . *The Astronomical Journal* 155 (6), 251.
URL <http://dx.doi.org/10.3847/1538-3881/aac172>
- Teanby, N. A., Irwin, P. G. J., de Kok, R. J., Apr 2010. Compositional evidence for Titan's stratospheric tilt. *Planetary and Space Science* 58 (5), 792–800.
URL <http://dx.doi.org/10.1016/j.pss.2009.12.005>
- Teodoro, L. F., Kegerreis, J. A., Estrada, P. R., Ćuk, M., Eke, V. R., Cuzzi, J. N., Massey, R. J., Sandnes, T. D., 2023. A recent impact origin of Saturn's rings and mid-sized moons. *The Astrophysical Journal* 955 (2), 137.
- Tewelde, Y., Perron, J. T., Ford, P., Miller, S., Black, B., 2013. Estimates of fluvial erosion on Titan from sinuosity of lake shorelines. *Journal of Geophysical Research: Planets* 118 (10), 2198–2212.
- Thakur, A. C., Remsing, R. C., 2023. Molecular Structure, Dynamics, and Vibrational Spectroscopy of the Acetylene: Ammonia (1: 1) Plastic Co-Crystal at Titan Conditions. *ACS Earth and Space Chemistry* 7 (2), 479–489.
- Thelen, A. E., Cordiner, M. A., Nixon, C. A., Vuitton, V., Kisiel, Z., Charnley, S. B., Palmer, M. Y., Teanby, N. A., Irwin, P. G. J., Nov 2020. Detection of CH_3C_3N in Titan's Atmosphere. *The Astrophysical Journal* 903 (1),

L22.

URL <https://doi.org/10.3847/2041-8213/abc1e1>

Thelen, A. E., Nixon, C., Chanover, N., Cordiner, M., Molter, E., Teanby, N., Irwin, P., Serigano, J., Charnley, S., Feb 2019. Abundance measurements of Titan's stratospheric HCN, HC₃N, C₃H₄, and CH₃CN from ALMA observations. *Icarus* 319, 417–432.

URL <http://dx.doi.org/10.1016/j.icarus.2018.09.023>

Thelen, A. E., Nixon, C., Chanover, N., Molter, E., Cordiner, M., Achterberg, R., Serigano, J., Irwin, P., Teanby, N., Charnley, S., Jun 2018. Spatial variations in Titan's atmospheric temperature: ALMA and Cassini comparisons from 2012 to 2015. *Icarus* 307, 380–390.

URL <http://dx.doi.org/10.1016/j.icarus.2017.10.042>

Thomas, C., Mousis, O., Ballenegger, V., Picaud, S., 2007. Clathrate hydrates as a sink of noble gases in Titan's atmosphere. *Astronomy & Astrophysics* 474 (2), L17–L20.

Thomas, C., Picaud, S., Mousis, O., Ballenegger, V., 2008. A theoretical investigation into the trapping of noble gases by clathrates on Titan. *Planetary and Space Science* 56 (12), 1607–1617.

Thompson, W. R., Squyres, S. W., 1990. Titan and other icy satellites: Dielectric properties of constituent materials and implications for radar sounding. *Icarus* 86 (2), 336–354.

Tobie, G., Choukroun, M., Grasset, O., Le Mouélic, S., Lunine, J. I., Sotin, C., Bourgeois, O., Gautier, D., Hirtzig, M., Lebonnois, S., Le Corre, L., Feb. 2009. Evolution of Titan and implications for its hydrocarbon cycle. *Royal Society of London Philosophical Transactions Series A* 367, 617–631.

Tobie, G., Lunine, J. I., Sotin, C., Mar. 2006. Episodic outgassing as the origin of atmospheric methane on Titan. *Nature* 440, 61–64.

Tobie, G., Teanby, N., Coustenis, A., Jaumann, R., Raulin, F., Schmidt, J., Carrasco, N., Coates, A., Cordier, D., De Kok, R., et al., Dec 2014. Science goals and mission concept for the future exploration of Titan and Enceladus. *Planetary and Space Science* 104, 59–77.

URL <http://dx.doi.org/10.1016/j.pss.2014.10.002>

- Tokano, T., 2008. Dune-forming winds on Titan and the influence of topography. *Icarus* 194 (1), 243–262.
- Tokano, T., 2009. Impact of seas/lakes on polar meteorology of Titan: Simulation by a coupled GCM-Sea model. *Icarus* 204 (2), 619–636.
- Tokano, T., 2013. Wind-induced equatorial bulge in Venus and Titan general circulation models: Implication for the simulation of superrotation. *Geophysical Research Letters* 40 (17), 4538–4543.
- Tokano, T., 2019. Orbitally and geographically caused seasonal asymmetry in Titan's tropospheric climate and its implication for the lake distribution. *Icarus* 317, 337–353.
- Tomasko, M. G., Archinal, B., Becker, T., Bézard, B., Bushroe, M., Combes, M., Cook, D., Coustenis, A., de Bergh, C., Dafoe, L. E., Doose, L., Douté, S., Eibl, A., Engel, S., Gliem, F., Grieger, B., Holso, K., Howington-Kraus, E., Karkoschka, E., Keller, H. U., Kirk, R., Kramm, R., Küppers, M., Lanagan, P., Lellouch, E., Lemmon, M., Lunine, J., McFarlane, E., Moores, J., Prout, G. M., Rizk, B., Rosiek, M., Rueffer, P., Schröder, S. E., Schmitt, B., See, C., Smith, P., Soderblom, L., Thomas, N., West, R., Dec. 2005. Rain, winds and haze during the Huygens probe's descent to Titan's surface. *Nature* 438, 765–778.
- Trainer, M., Brinckerhoff, W., Grubisic, A., Danell, R., Kaplan, D., van Amerom, F., Li, X., Freissinet, C., Szopa, C., Buch, A., et al., 2021. Development of the Dragonfly Mass Spectrometer (DraMS) for Titan. In: Lunar and Planetary Science Conference. No. 2548. p. 1532.
- Tseng, W.-L., Ip, W.-H., Kopp, A., 2008. Exospheric heating by pickup ions at Titan. *Advances in Space Research* 42 (1), 54–60.
- Turtle, E., Perry, J., Barbara, J., Del Genio, A., Rodriguez, S., Le Mouélic, S., Sotin, C., Lora, J., Faulk, S., Corlies, P., et al., 2018. Titan's meteorology over the Cassini mission: Evidence for extensive subsurface methane reservoirs. *Geophysical Research Letters* 45 (11), 5320–5328.
- Turtle, E. P., Del Genio, A. D., Barbara, J. M., Perry, J. E., Schaller, E. L., McEwen, A. S., West, R. A., Ray, T. L., Feb. 2011a. Seasonal changes in Titan's meteorology. *Geophysical Research Letters* 38, 3203.

- Turtle, E. P., Perry, J. E., Hayes, A. G., Lorenz, R. D., Barnes, J. W., McEwen, A. S., West, R. A., Del Genio, A. D., Barbara, J. M., Lunine, J. I., Schaller, E. L., Ray, T. L., Lopes, R. M. C., Stofan, E. R., Mar. 2011b. Rapid and Extensive Surface Changes Near Titan's Equator: Evidence of April Showers. *Science* 331, 1414–.
- Turtle, E. P., Perry, J. E., Hayes, A. G., McEwen, A. S., Apr. 2011c. Shoreline retreat at Titan's Ontario Lacus and Arrakis Planitia from Cassini Imaging Science Subsystem observations. *Icarus* 212, 957–959.
- Turtle, E. P., Perry, J. E., McEwen, A. S., Del Genio, A. D., Barbara, J., West, R. A., Dawson, D. D., Porco, C. C., Jan. 2009. Cassini imaging of Titan's high-latitude lakes, clouds, and south-polar surface changes. *Geophysical Research Letters* 36, 2204.
- Tyler, R., 2014. Comparative estimates of the heat generated by ocean tides on icy satellites in the outer Solar System. *Icarus* 243, 358–385.
- Tyler, R. H., 2008. Strong ocean tidal flow and heating on moons of the outer planets. *Nature* 456 (7223), 770–772.
- Vance, S. D., Panning, M. P., Stähler, S., Cammarano, F., Bills, B. G., Tobie, G., Kamata, S., Kedar, S., Sotin, C., Pike, W. T., et al., 2018. Geophysical investigations of habitability in ice-covered ocean worlds. *Journal of Geophysical Research: Planets* 123 (1), 180–205.
- Vinatier, S., Schmitt, B., Bézard, B., Rannou, P., Dauphin, C., de Kok, R., Jennings, D., Flasar, F., Aug 2018. Study of Titan's fall southern stratospheric polar cloud composition with Cassini/CIRS: Detection of benzene ice. *Icarus* 310, 89–104.
 URL <http://dx.doi.org/10.1016/j.icarus.2017.12.040>
- Vincent, D., Lambrechts, J., Tyler, R. H., Karatekin, Ö., Dehant, V., Deleersnijder, É., 2022. A numerical study of the liquid motion in Titan's subsurface ocean. *Icarus* 388, 115219.
- Vu, T., Choukroun, M., Sotin, C., Muñoz-Iglesias, V., Maynard-Casely, H., 2020a. Rapid formation of clathrate hydrate from liquid ethane and water ice on Titan. *Geophysical Research Letters* 47 (4), e2019GL086265.

- Vu, T. H., Cable, M. L., Choukroun, M., Hodyss, R., Beauchamp, P., 2014. Formation of a new benzene–ethane co-crystalline structure under cryogenic conditions. *The Journal of Physical Chemistry A* 118 (23), 4087–4094.
- Vu, T. H., Maynard-Casely, H. E., Cable, M. L., Hodyss, R., Choukroun, M., Malaska, M. J., 2020b. Anisotropic thermal expansion of the acetylene–ammonia co-crystal under Titan’s conditions. *Journal of Applied Crystallography* 53 (6), 1524–1530.
- Vuitton, V., Yelle, R. V., Klippenstein, S. J., Hörst, S. M., Lavvas, P., 2019. Simulating the density of organic species in the atmosphere of Titan with a coupled ion-neutral photochemical model. *Icarus*, 120–197.
- Vuitton, V., Yelle, R. V., McEwan, M. J., Nov. 2007. Ion chemistry and N-containing molecules in Titan’s upper atmosphere. *Icarus* 191, 722–742.
- Waite, J. H., Niemann, H., Yelle, R. V., Kasprzak, W. T., Cravens, T. E., Luhmann, J. G., McNutt, R. L., Ip, W.-H., Gell, D., De La Haye, V., Müller-Wordag, I., Magee, B., Borggren, N., Ledvina, S., Fletcher, G., Walter, E., Miller, R., Scherer, S., Thorpe, R., Xu, J., Block, B., Arnett, K., May 2005. Ion Neutral Mass Spectrometer Results from the First Flyby of Titan. *Science* 308, 982–986.
- Waite, J. H., Young, D. T., Cravens, T. E., Coates, A. J., Crary, F. J., Magee, B., Westlake, J., May 2007. The Process of Tholin Formation in Titan’s Upper Atmosphere. *Science* 316, 870–.
- Waite, Jr., J. H., Lewis, W. S., Magee, B. A., Lunine, J. I., McKinnon, W. B., Glein, C. R., Mousis, O., Young, D. T., Brockwell, T., Westlake, J., Nguyen, M.-J., Teolis, B. D., Niemann, H. B., McNutt, R. L., Perry, M., Ip, W.-H., Jul. 2009. Liquid water on Enceladus from observations of ammonia and ^{40}Ar in the plume. *Nature* 460, 487–490.
- Wilson, E., Atreya, S., 2003. Chemical sources of haze formation in Titan’s atmosphere. *Planetary and Space Science* 51 (14-15), 1017–1033.
- Wilson, E. H., Atreya, S. K., Sep. 2009. Titan’s Carbon Budget and the Case of the Missing Ethane. *Journal of Physical Chemistry A* 113, 11221–11226.

- Wisdom, J., Dbouk, R., Militzer, B., Hubbard, W. B., Nimmo, F., Downey, B. G., French, R. G., 2022. Loss of a satellite could explain Saturn's obliquity and young rings. *Science* 377 (6612), 1285–1289.
- Witasse, O., Lebreton, J.-P., Bird, M. K., Dutta-Roy, R., Folkner, W. M., Preston, R. A., Asmar, S. W., Gurvits, L. I., Pogrebenko, S. V., Avruch, I. M., et al., 2006. Overview of the coordinated ground-based observations of Titan during the Huygens mission. *Journal of Geophysical Research: Planets* 111 (E7).
- Wong, A.-S., Morgan, C. G., Yung, Y. L., Owen, T., Feb. 2002. Evolution of CO on Titan. *Icarus* 155, 382–392.
- Wong, E., Brasser, R., Werner, S., Kirchoff, M., 2023. Saturn's ancient regular satellites. *Icarus* 406, 115763.
- Wong, M. L., Yung, Y. L., Randall Gladstone, G., 2015. Pluto's implications for a Snowball Titan. *Icarus* 246, 192–196, special Issue: The Pluto System.
URL <https://www.sciencedirect.com/science/article/pii/S0019103514002723>
- Wood, C. A., Radebaugh, J., 2020. Morphologic evidence for volcanic craters near Titan's north polar region. *Journal of Geophysical Research: Planets* 125 (8), e2019JE006036.
- Wye, L. C., Zebker, H. A., Ostro, S. J., West, R. D., Gim, Y., Lorenz, R. D., Team, C. R., et al., 2007. Electrical properties of Titan's surface from Cassini RADAR scatterometer measurements. *Icarus* 188 (2), 367–385.
- Yu, X., Hörst, S. M., He, C., Bridges, N. T., Burr, D. M., Sebree, J. A., Smith, J. K., 2017a. The effect of adsorbed liquid and material density on saltation threshold: Insight from laboratory and wind tunnel experiments. *Icarus* 297, 97–109.
- Yu, X., Hörst, S. M., He, C., McGuigan, P., Bridges, N. T., 2017b. Direct measurement of interparticle forces of Titan aerosol analogs (“tholin”) using atomic force microscopy. *Journal of Geophysical Research: Planets* 122 (12), 2610–2622.

- Yu, X., Hörst, S. M., He, C., McGuigan, P., Crawford, B., 2018. Where does Titan sand come from: insight from mechanical properties of Titan sand candidates. *Journal of Geophysical Research: Planets* 123 (9), 2310–2321.
- Yu, X., Hörst, S. M., He, C., McGuigan, P., Kristiansen, K., Zhang, X., 2020. Surface energy of the Titan aerosol analog “tholin”. *The Astrophysical Journal* 905 (2), 88.
- Yung, Y. L., Allen, M., Pinto, J. P., Jul. 1984. Photochemistry of the atmosphere of Titan - Comparison between model and observations. *Astrophys. J. Suppl.* 55, 465–506.
- Zahnle, K. J., Korycansky, D. G., Nixon, C. A., 2014. Transient climate effects of large impacts on Titan. *Icarus* 229, 378–391.
- Zalucha, A., Fitzsimmons, A., Elliot, J., Thomas-Osip, J., Hammel, H., Dhillon, V., Marsh, T., Taylor, F., Irwin, P., 2007. The 2003 November 14 occultation by Titan of TYC 1343-1865-1: II. Analysis of light curves. *Icarus* 192 (2), 503–518.