



Use of crops by the partridge (*Perdix perdix*) in the context of methanization

Lucille Capitaine, Rémi Helder, E. Guillon

► To cite this version:

Lucille Capitaine, Rémi Helder, E. Guillon. Use of crops by the partridge (*Perdix perdix*) in the context of methanization. Ecology and Behaviour, Aug 2024, Villiers-en-Bois, France. hal-04902108

HAL Id: hal-04902108

<https://hal.science/hal-04902108v1>

Submitted on 20 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Use of crops by the partridge (*Perdix perdix*) in the context of methanization


L.CAPITAIN^{1,2}, R.HELDER¹, E.GUILLO²

¹ Centre de Recherche et de Formation en Eco-éthologie – Université de Reims Champagne-Ardenne - France

² Institut de chimie Moléculaire de Reims – UMR 7312 – Université de Reims Champagne-Ardenne - France

Background


For methanization, farmers use new crops called intermediate energy crops to put into the biogas plant. These new crops lead to **changes in the landscape**.



Objectives

- Study habitat selection by partridges over the course of the year in cereal plains.
- Study the selection of energy crops by partridges.


Capture



Analyses



Winter : 1 December to 29 February



Spring : 1 March to 31 May

1-For each period : calculation of Home Range (HR) and generating random locations



2- For each period : Calculation of Ivlev Electivity index^[2]


$$E = \frac{pi - p}{pi + p}$$

pi : Proportion of partridge GPS point in a habitat for individual i
p : Proportion of random point for the same habitat

→ Index from -1 to 1


Value of -1 : strong avoidance
Value of 1 : strong preference

Importance of height ?




Results

a) Winter : 01/12 au 29/02 n = 6 ind



b) Spring : 01/03 au 31/05 n = 7 ind



First conclusions

- Partridge don't seem to use winter energy crops in any season.
- They use crops differently depending on the season, probably due to the change in crop height.
- The greater use of paths in spring could be explained by the greater height of crops or by the greater food availability in grass strip.
- Energy crops are higher than classic crops, which may explain why winter energy crops are not selected.

References

- [1] Kranstauber, B., Kays, R., LaPoint, S. D., Wikelski, M., & Safi, K. (2012). A dynamic Brownian bridge movement model to estimate utilization distributions for heterogeneous animal movement. *Journal of Animal Ecology*, 81(4), 738–746. <https://doi.org/10.1111/j.1365-2656.2012.01955.x>
- [2] Gautrelet, M., Gerard, J.-F., Helder, R., Fournier, P., Fournier-Chambrillon, C., Hubert, P., Isère-Lauqué, E., Capitaine, L., Dupuy, M., Dispain De Floran, L., Alleman, C., & Devos, A. (2024). First look on the home range, movement, and habitat selection of the invasive Northern raccoon (*Procyon lotor*) in France through two contrasted populations. *European Journal of Wildlife Research*, 70(1), 7. <https://doi.org/10.1007/s10344-023-01756-1>