

Table 2: Number of periods of digestive drug uses according to atropinic exposure status

	<b>Unexposed (AB*=0) N=5,720</b>	<b>Exposed (AB*&gt;0) N=2,652</b>	<i>P<sup>1</sup></i>	<b>Exposed (AB* [1-3]) N=1,386</b>	<b>Exposed (AB* [3-9]) N=1,152</b>	<b>Exposed (AB*≥9) N=114</b>	<i>P<sup>2</sup></i>
<b>Number of periods of digestive drug uses</b>	<b>4.0±3.8</b>	<b>5.0±4.2</b>	<b>&lt;.0001</b>	<b>4.9±4.0</b>	<b>5.1±4.3</b>	<b>6.1±4.8</b>	<b>&lt;.0001</b>
Drugs for constipation	0.3±1.0	0.4±1.2	<.0001	0.4±1.0	0.4±1.3	0.5±1.3	<.0001
Drugs for diarrhea	2.0±1.8	2.5±2.1	<.0001	2.5±2.1	2.5±2.1	2.8±2.2	<.0001
Drugs for vomiting or gastroesophageal reflux disease	2.2±2.7	2.8±3.1	<.0001	2.7±3.0	2.9±3.2	3.2±3.4	<.0001
Drugs for gastro-intestinal pain	0.5±1.1	0.7±1.2	<.0001	0.7±1.1	0.7±1.2	1.1±1.9	<.0001

\*AB= Atropinic Burden

P<sup>1</sup>: P value for the comparisons of the exposed and unexposed groups

P<sup>2</sup>: P value for the comparisons of the different atropinic burden groups