



HAL
open science

Addressing adverse synergies between chemical and biological pollutants at schools-The ‘ SynAir-G ’ hypothesis

Nikolaos Papadopoulos, Cezmi Akdis, Mubeccel Akdis, Athanasios Damialis, Giuseppina Esposito, Ioana Fergadiotou, Christian Goroncy, Pierre Guitton, Maia Gotua, Kleanthis Erotokritou, et al.

► **To cite this version:**

Nikolaos Papadopoulos, Cezmi Akdis, Mubeccel Akdis, Athanasios Damialis, Giuseppina Esposito, et al.. Addressing adverse synergies between chemical and biological pollutants at schools-The ‘ SynAir-G ’ hypothesis. *Allergy*, 2023, 79 (2), pp.294-301. 10.1111/all.15857 . hal-04904605

HAL Id: hal-04904605

<https://hal.science/hal-04904605v1>

Submitted on 21 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

REVIEW ARTICLE

Addressing adverse synergies between chemical and biological pollutants at schools—The ‘SynAir-G’ hypothesis

Nikolaos G. Papadopoulos^{1,2}  | Cezmi A. Akdis³  | Mubeccel Akdis³  | Athanasios Damialis⁴  | Giuseppina Esposito⁵  | Ioana Fergadiotou⁶  | Christian Goroncy⁷ | Pierre Guitton⁸ | Maia Gotua⁹  | Kleantlis Erotokritou¹⁰  | Tuomas Jartti^{11,12,13}  | Clare Murray¹  | Athanasios Nenes¹⁴  | Sotirios Nikolettseas¹⁵  | Susetta Finotto¹⁶  | Spyros N. Pandis¹⁷  | Valeria Ramiconi¹⁸ | Angela Simpson¹  | Aki Soudunsaari¹⁹ | Anna Stårbröst²⁰ | Maria Staiano²¹  | Antonio Varriale^{21,22}  | Paraskevi Xepapadaki²  | Torsten Zuberbier^{23,24,25}  | Isabella Annesi-Maesano²⁶  | The SynAir-G Consortium

Correspondence

Nikolaos G. Papadopoulos, Division of Immunology, Immunity to Infection and Respiratory Medicine, School of Biological Sciences, The University of Manchester, Manchester, UK.
Email: ngp@allergy.gr

Funding information

HORIZON EUROPE Health, Grant/Award Number: 101057271

Abstract

While the number and types of indoor air pollutants is rising, much is suspected but little is known about the impact of their potentially synergistic interactions, upon human health. Gases, particulate matter, organic compounds but also allergens and viruses, fall within the ‘pollutant’ definition. Distinct populations, such as children and allergy and asthma sufferers are highly susceptible, while a low socioeconomic background is a further susceptibility factor; however, no specific guidance is available. We spend most of our time indoors; for children, the school environment is of paramount importance and potentially amenable to intervention. The interactions between some pollutant classes have been studied. However, a lot is missing with respect to understanding interactions between specific pollutants of different classes in terms of concentrations, timing and sequence, to improve targeting and upgrade standards. SynAir-G is a European Commission-funded project aiming to reveal and quantify synergistic interactions between different pollutants affecting health, from mechanisms to real life, focusing on the school setting. It will develop a comprehensive and responsive multipollutant monitoring system, advance environmentally friendly interventions, and disseminate the generated knowledge to relevant stakeholders in accessible and actionable formats. The aim of this article is to put forward the SynAir-G hypothesis, and describe its background and objectives.

KEYWORDS

aerobiology, allergy, artificial intelligence, asthma, biosensors, children, school, virus, volatile organic compounds

See Appendix for The SynAir-G Consortium.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd.

1 | THERE IS A MAJOR UNMET NEED TO CHARACTERISE, MONITOR AND IMPROVE AIR QUALITY TO SUPPORT HEALTH

The profound impact of air pollution on human health has been thoroughly studied and reported.^{1,2} The risk involves most systems of our body, with skin, respiratory, cardiovascular and central nervous systems predominating.² Overall, acute and chronic illnesses, as well as exacerbations of chronic diseases, can result in humans and animals following damage of epithelial barriers, activation of inflammatory pathways, DNA damage and cell death responses.³ The list of substances with negative potential impacts on human, as well as animal and environmental, health, is large and growing.¹ Some, such as fine particulate matter (PM), O₃, CO, NO_x and SO₂ are well characterized and monitored, while others, such as ultrafine particles (UPF), or radon, are uncertain or unknown. In many environments, the threat is constant and affects almost everybody. More than 10 million deaths can be attributed to air pollution every year.⁴ It is both fortunate and tragic that these events are largely preventable.² Sadly, much of the impact predominantly involves the developing world.^{2,5} Although this is not always the case, many of the adverse health effects can rapidly improve and reverse if the source of pollution is removed.² For example, the smoking ban in indoor places has resulted in improved health not only for hospitality workers but also for the general population.² Improving air quality probably has benefits even beyond the currently accepted standards. Furthermore, targeting air quality has often added considerable value to wider pollution, climate change and planetary health. There is, therefore, a major unmet need to characterise, monitor and improve air quality to support health.

2 | WE SPEND MOST OF OUR TIMES INDOORS - INDOOR AIR QUALITY (IAQ) HAS NOT RECEIVED AS MUCH ATTENTION

In the last decades, outdoor, or 'ambient', air pollution has attracted the most attention in research and media alike. However, nowadays there is increased concern and focus on household and more generally indoor air pollution.^{1,2,6,7} We spend more than 90% of our time indoors; furthermore, buildings are increasingly airtight to conserve energy among others.^{8,9} Some pollutants have higher concentrations in the ambient air; however, others accumulate indoors, while different outdoor-indoor gradients are generated depending on pollutant source, ventilation and local conditions.¹⁰ Key pollutants for human health are generated by cooking, heating and cleaning. These can vary widely between households; in contrast to outdoor air pollution that can be monitored through a relatively small number of stations and geographical or atmospheric and meteorological models or remote sensors,¹¹ indoor air monitoring does not apply in most environments and indoor air quality can literally be unpredictable because of a huge variety of factors that influence it.^{9,12} In recent years, comparative risk studies performed by the Environmental

Protection Agency of the USA have consistently ranked indoor air pollution among the top five environmental risks to public health.¹³

3 | THE EFFECTS OF INDOOR AIR POLLUTION CAN BE PARTICULARLY PRONOUNCED IN DISTINCT POPULATIONS. CHILDREN WITH ALLERGY AND ASTHMA ARE A MAJOR SUCH POPULATION

The variability of indoor air quality, but also the differential susceptibility of discrete people groups, result in considerable inequalities and pronounced health effects in different populations.^{14,15} Economy and access to clean technologies is undoubtedly a major determinant. Across populations, children are more susceptible than adults, as air pollutants may directly interfere with their development.^{8,16,17} Also, in children, respiratory rate is faster, bringing in proportionally more pollutants, while airways are smaller and therefore easier to become obstructed.⁸ Individuals with hypersensitivities, such as allergies and asthma, are also more susceptible: they tend to have exaggerated and persistent inflammation, triggered by specific and non-specific stimuli, including tobacco smoke, particular matter and other pollutants. Children with low socioeconomic status are also more vulnerable to air pollution effects due to higher exposure, nutrition problems, lack of capacity to anticipate, cope and adapt to the hazards, and reduced health access.¹⁸⁻²⁰ While the political connotations of socioeconomic disequilibrium are beyond the scope of this paper, an extended discussion on environmental justice and radical measures to achieve it are absolutely necessary. Furthermore, the developing diseases are not only affecting the organ of exposure, such as the respiratory system or skin; activated inflammatory cells from these organs can migrate to distant tissues and exacerbate chronic diseases there.²¹ A strong link to autoimmune diseases has been reported and there is emerging data demonstrating the influence of respiratory exposure to air pollutants on neuropsychiatric conditions. Headaches, fatigue, shortness of breath, sinus congestion, coughing, sneezing, dizziness, nausea and irritation of the eyes, nose, throat and skin are some of the symptoms.²² Risks include both increased symptoms or disease exacerbations, but also inability to prevent or treat, due to socioeconomic drawbacks.

4 | THE IMPACT OF ALLERGY AND ASTHMA

Evidently, these are not niche populations! There are more than 10-30 million people in Europe suffering from asthma.²³ Asthma is most often associated with allergy: the allergy epidemic is by itself the most common chronic condition across the life cycle, starting from early childhood, persisting through the productive years, and having an impact on healthy ageing. Allergic rhinitis, the most frequent respiratory allergy co-exists with asthma in most cases, while other forms, either organ-specific (e.g., conjunctivitis, dermatitis),

or systemic (e.g., anaphylaxis) co-exist in the same patient, increasing the personal and societal burden, and costing over €150 billion annually.²⁴

5 | POLLUTANTS ARE MANY, INCLUDING CHEMICAL AND BIOLOGICAL SUBSTANCES

The definition of 'air pollution' according to the WHO is 'contamination of the indoor or outdoor environment by any *chemical, physical or biological agent* that modifies the natural characteristics of the atmosphere'.²⁵ Different groups of pollutants of public health relevance have been identified and characterized: particulate matter, gases such as carbon monoxide, nitrogen dioxide, ozone, polycyclic aromatic hydrocarbons (PAHs) and a multitude of volatile organic compounds (VOCs).^{8,9} It has to be noted that more than 350,000 new chemical substances alone or in combination have been introduced to human lives with almost no regulatory control on their health effects after the 1960s. Many of them ended up as pollutants. The identities of many of these remain publicly unknown because they are claimed as confidential (over 50,000) or ambiguously described (up to 70,000).²⁶

While most often the concept of pollution and pollutants has been associated with chemical substances, there is an increased realization that biological pollutants are at least equally important.⁵ This gained considerable attention with the COVID-19 pandemic: while many microorganisms present in bioaerosols are part of the natural atmosphere and necessary for our microbiome, others can be highly detrimental, resulting in considerable morbidity and mortality.²⁷ Often, this depends upon the concentration of the microbial agent, as well as the immune status of an individual, which can be influenced by societal factors, such as congregation.^{1,28,29} For the large allergic population, allergens to which they are sensitized act as pollutants. In this case, as well, it is not only the individual susceptibility that defines the pollutant, but also the result of human interventions that radically change the concentration of for example, pollen or spores in the air. The increasing abundance of pollinating plants, such as the invasive *Ambrosia* spp., and their rising atmospheric pollen concentrations as a result of global warming, is a well-established, typical example.^{24,30,31} Outdoor air pollutants are a major source for indoor pollutants; in some cases, outdoor pollutants may be present in increased concentrations indoors, however, the composition of pollutants in indoor spaces—and therefore their potential interactions—can be highly variable.

6 | THERE ARE STRONG INDICATIONS THAT CHEMICAL AND BIOLOGICAL POLLUTANTS INTERACT AND POSSIBLY SYNERGISE

There is evidence suggesting that the impact of air pollutants on health goes beyond the cumulative effect of the different

pollutants. Although still not completely characterised, the interactions between chemical and biological pollutants, including allergens, may be in some cases synergistic.³²


6.1 | Interactions of chemical pollutants with allergens

Diesel exhaust (DE) emissions have been shown to be able to underpin allergen sensitisation³³ but also augment allergen-induced airway disease. In vitro exposure of peripheral blood mononuclear cells of allergic patient to DE, together with the major mite allergen Der p1, synergistically increased the production of the neutrophil chemoattractant IL-8.³⁴ In a French epidemiological study, levels of aldehydes in the house had a significant synergistic effect with the presence of the major dog allergen Can f1, although this was not the case for other interactions.³⁵ Using an in vivo exposure model in humans, allergic individuals were exposed for 2 h to particulate matter, followed by a bronchial allergen challenge. Levels of eosinophils, IL-5 and ECP were increased.³⁶ Additive, but not synergistic, effects have also been shown in a similar experimental setting, on the expression of microRNAs.³⁷ In contrast, in another epigenetic study, when exposure was separated by a 4-h time interval, differential methylation was observed to a much larger extent; furthermore, the timing of exposure altered the results.³⁸ In yet another human study, co-exposure to DE and allergen (house dust mite, birch or grasses) not only altered the allergen-mediated secretome by enhancing or suppressing specific proteins but also induced protein changes unique to co-exposure.³⁹ Lastly, there is evidence of interaction between chemical air pollutants and pollen grains. Chemical pollutants might facilitate pollen allergen release, act as adjuvants to stimulate IgE-mediated responses, modify allergenic potential and enhance the expression of some allergens in pollen grains.³⁹

6.2 | Interactions of viral infections with allergens

The allergen–virus interaction has been particularly studied in the context of asthma. In studies assessing exposures of either adults or children hospitalized for an acute exacerbation of asthma, it has been shown that the risk for such an event increased synergistically when they were infected and also exposed to an allergen to which they are sensitized.^{20,40} A typical upper respiratory infection, most often associated with human rhinoviruses, may induce airway hyper-responsiveness (AHR) even in healthy individuals.⁴¹ AHR is a cardinal characteristic of asthma and can be prolonged up to five times, in atopic children through repeated infections.⁴² An allergen can act either through IgE recognition or directly on the epithelium through protease activity.¹⁹ Timing of exposure appears to be crucial, as both in vitro¹⁹ and in vivo¹⁸ studies showed that synergy may not occur when allergen exposure happens first. Interestingly, recent observations suggested that allergen exposure, in sensitized individuals, may in fact protect them from infection with SARS-CoV2!⁴³ However,

FIGURE 1 Chemical pollutants, allergens and viruses affect the epithelial barrier and the immune system compromising health. While direct effects (blue lines) have been studied extensively, the interactions between chemical and biological pollutants (red lines) have not been adequately evaluated. Although all may be affected, young children and particularly those with allergy and asthma, are more susceptible to high indoor concentrations of pollutants at school.



pollen exposure is associated with an increased risk of viral infections, including both rhinoviruses (RVs) and SARS-CoV2.^{41,44–46} Therefore, when evaluating the impact of combined exposures, it is important to consider the immune background of individuals or groups, as well as the type and sequence of exposures.

6.3 | Interactions of chemical pollutants with viruses

Epidemiological data from children demonstrate more severe viral-induced asthma symptoms and worse lung function outcomes in the presence of increased exposure to high concentrations of chemical air pollutants.^{46,47} Ambient PM and DE particles compromise the integrity of the epithelial barrier, while NO₂ exposure increases epithelial expression of ICAM-1, the major receptor for RVs.⁴⁸ Indeed, PM increases susceptibility to RV epithelial infection *in vitro*.⁴⁹ Furthermore, exposure to air pollutants increases substantially oxidative stress, which can diminish Th1 and favour Th2 responses, but also disrupt the balance between Th1 and cytotoxic T lymphocyte responses, potentially affecting the antiviral defence mechanisms.⁵⁰ A large body of data has recently confirmed the effect of ambient pollution on susceptibility to SARS-CoV2 and the development of severe disease.²⁸

It should not be overlooked that additional interactions within the wide groups of 'chemical' and 'biological' pollutants have also been identified or suggested to different extents, such as interactions between viruses and bacteria⁵¹ or between ozone and other chemicals,⁵² as well as in the presence of allergens.

7 | SCHOOLS ARE PARTICULARLY IMPORTANT FOR CHILDREN

Children spend up to 10h a day at school, most of the time indoors^{6,8} and the density of persons per square meter during this time is much higher than in the usual working place or at home. Concentrations of particulate matter and PAHs higher than those

required by international guidelines have been found in around 30% of schools.^{8,53–55} Levels of exposure are amenable to change, as it has been shown for example, following school building renovation.⁵⁶ Nevertheless, it is not surprising that the highest levels of chemical pollutant exposure happen in schools located in highly polluted, low socioeconomic background areas. Simple field interventions, such as ventilation schedules and/or using air purifiers, have the capacity of reducing some of the effects, however many rely on changing habits, which is challenging.^{9,12,57} Furthermore, schools are the breeding grounds of respiratory viruses. The risk of transmission of SARS-CoV2 in schools has attracted a lot of attention; while the extent and relative importance of this mode of transmission is still scrutinized, there is no doubt that viral transmission either through aerosols or by hand contact is important.⁵⁸ Typically, high peaks of respiratory infections, followed by asthma attacks in susceptible individuals and their families, take place during the month of September in the Northern Hemisphere, coinciding with the return of schoolchildren after their summer vacation—the so-called 'September epidemic',⁵⁹ attributed mostly to rhinoviruses. At the same time, cat allergens, brought from home on the clothes of children are present in increased concentrations within classrooms. French data suggested that the combined action of sensitization to house dust mites and viruses on asthma attacks during the epidemics among the pupils might be quantitatively increased by the presence of stress as assessed objectively through the Strength and Difficulties Questionnaire.^{60,61} Many more allergens, such as cockroaches, mites and moulds, with the potential to sensitize and produce symptoms in sensitized individuals, are present at schools.⁶⁰ It is therefore plausible that schools are ideal grounds for interacting pollutants to affect the health of children—particularly those with hypersensitivity.

8 | THE SynAir-G HYPOTHESIS

To further explore the nature and impact of interactions between common chemical and biological pollutants, and in the context of the Horizon Europe project 'SynAir-G' (Disrupting Noxious Synergies

of Indoor Air Pollutants and their Impact in Childhood Health and Wellbeing, using Advanced Intelligent Multisensing and Green Interventions) we have generated the following working hypothesis: 'We hypothesise that major indoor air effects on respiratory, immune and mental health are modulated by synergistic interactions between biological and chemical pollutants' (Figure 1). This hypothesis needs to be evaluated both epidemiologically, in different populations and subgroups, and mechanistically, using state-of-the-art in vitro and in vivo models.

9 | NEEDS AND PROJECTED ACTIONS AND INNOVATION

The above hypothesis will be evaluated thoroughly in real-life conditions, as well as in experimental systems in the context of SynAir-G, with the participation of 18 partners from 11 countries. Details about the protocols and technical approaches used can be found in the project's website.⁶² SynAir-G will develop and deploy novel sensor technologies in schools, as well as intelligent wearable devices,⁶³ and evaluate the impact of the exposome on health, combining mathematical, statistical and computational means, including advanced methods from machine learning and artificial intelligence (AI). Citizen science and participatory design principles will be used in the establishment and follow-up of the cohorts. Recent studies indicate that when air pollution sensors are coupled with physiological sensors and location, a stronger connection can be made between a person's exposure environment and health indicators.⁶⁴ SynAir-G builds on the cross-fertilisation and analysis of data collected from various sources (questionnaires, sensors, wearables), accounting for privacy and embedding ethics in the development of innovative AI methods.⁶⁵ Several interventions, including air filtering and a green wall, will also be evaluated. Socioeconomic determinants will also be put into context. While it is expected that these investigations will improve our targeting and capacity for addressing the problem, the list of needs and potential mitigation measures is large and should not be delayed. Monitoring systems, air filtration measures as well as lifestyle interventions to develop healthy habits, are imminently required, together with measures to address inequalities, to support the next generation to develop and retain optimal health. Coordinated efforts by all stakeholders including scientists from different disciplines and regulatory authorities are urgently needed, to implement novel discoveries and opportunities highlighted here.⁶⁶

AUTHOR CONTRIBUTIONS

Study conception: NGP. Study design: All authors. Critical revision of the manuscript: All authors.

AFFILIATIONS

¹Division of Immunology, Immunity to Infection and Respiratory Medicine, School of Biological Sciences, The University of Manchester, Manchester, UK

²Allergy Department, 2nd Paediatric Clinic, National and Kapodistrian University of Athens, Athens, Greece

³Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich,

Davos, Switzerland, Christine Kühne-Center for Allergy Research and Education (CK-CARE), Davos, Switzerland

⁴Department of Ecology, School of Biology, Aristotle University of Thessaloniki, Thessaloniki, Greece

⁵Bio Check Up (BCU), Naples, Italy

⁶INLECOM INNOVATION, Kifissia, Greece

⁷DIN Deutsches Institut fuer Normung EV, Berlin, Germany

⁸TEQOYA Company, Paris, France

⁹Center for Allergy and Immunology Research (CAIR), Tbilisi, Georgia

¹⁰CY.R.I.C Cyprus Research and Innovation Center Ltd, Nicosia, Cyprus

¹¹PEDEGO Research Unit, University of Oulu, Oulu, Finland

¹²Department of Pediatrics, Oulu University Hospital, Oulu, Finland

¹³Department of Pediatrics and Adolescent Medicine, Turku University Hospital and University of Turku, Turku, Finland

¹⁴Laboratory of Atmospheric Processes and their Impacts, School of Architecture, Civil and Environmental Engineering, Swiss Institute of Technology, Lausanne, Switzerland

¹⁵Computer Engineering and Informatics Department, University of Patras, Patras, Greece

¹⁶Molecular Pneumology Department, University Hospital of Erlangen, Erlangen, Germany

¹⁷Institute of Chemical Engineering Sciences (ICEHT), Foundation for Research and Technology Hellas (FORTH), Patras, Greece

¹⁸The European Federation of Allergy and Airways Diseases Patients' Association (EFA), Brussels, Belgium

¹⁹NAAVA, Helsinki, Finland

²⁰Lulea University of Technology, Lulea, Sweden

²¹Institute of Food Science, CNR Italy, Avellino, Italy

²²URT-ISA, CNR at Department of Biology, University of Naples Federico II, Naples, Italy

²³Institute of Allergology, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt Universität zu Berlin, Berlin, Germany

²⁴Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, Allergology and Immunology, Berlin, Germany

²⁵Global Allergy & Asthma European Network of Excellence-GA²LEN, Berlin, Germany

²⁶Department of Allergic and Respiratory Disease, Institut Desbrest of Epidemiology and Public Health, University of Montpellier and INSERM, Montpellier University Hospital, Montpellier, France

ACKNOWLEDGMENTS

The project is funded by the Horizon Europe project SynAir-G, grant agreement ID: 101057271.

CONFLICT OF INTEREST STATEMENT


NGP declares contracts from Capricare, Nestle, Numil, Vianex, REG, consulting fees from Abbott, Abbvie, Astra Zeneca, GSK, HAL, Medscape, Menarini/Faes Farma, Mylan, Novartis, Nutricia, OM Pharma, Regeneron/Sanofi outside the submitted work. C. A. Akdis declares research grants from the Swiss National Science Foundation, European Union (EU CURE, EU Syn-Air-G), Novartis Research Institutes, (Basel, Switzerland), Stanford University (Redwood City, Calif), Seed Health (Boston, USA) and SciBase (Stockholm, Sweden); is the Co-Chair for EAACI Guidelines on Environmental Science in Allergic diseases and Asthma; Chair of the EAACI Epithelial Cell Biology Working Group is on the Advisory Boards of Sanofi/Regeneron (Bern, Switzerland, New York, USA), Stanford University Sean Parker Asthma Allergy Center (CA, USA), Novartis (Basel, Switzerland), Glaxo Smith Kline (Zurich, Switzerland), Bristol-Myers Squibb (New York, USA), Seed Health (Boston, USA) and SciBase (Stockholm, Sweden); and is the Editor-in-Chief of Allergy. PX declares PX reports personal

fees from Nutricia, Nestle, Friesland, Uriach, Novartis Pharma AG and GlaxoSmithkline outside the submitted work. VR declares unrestricted Grants from Astra Zeneca, Chiesi, Sanofi, Regeneron, Pfizer, OM Pharma, DBV Technologies, Viatrix, Roche, Boehringer Ingelheim, AbbVie, Leo Pharma outside the submitted work. MA, AD, GE, IF, CG, PG, MG, KE, TJ, CM, AN, SN, SF, SP, AS, AK, AN, MS, AV, TS, IA-M, declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Nikolaos G. Papadopoulos  <https://orcid.org/0000-0002-4448-3468>

Cezmi A. Akdis  <https://orcid.org/0000-0001-8020-019X>

Mubeccel Akdis  <https://orcid.org/0000-0003-0554-9943>

Athanasios Damialis  <https://orcid.org/0000-0003-2917-5667>

Giuseppina Esposito  <https://orcid.org/0000-0003-2811-4678>

Ioana Fergadiotou  <https://orcid.org/0000-0002-8940-2094>

Maia Gotua  <https://orcid.org/0000-0003-2497-4128>

Kleanthis Erotokritou  <https://orcid.org/0000-0002-7284-104X>

Tuomas Jartti  <https://orcid.org/0000-0003-2748-5362>

Clare Murray  <https://orcid.org/0000-0002-8961-8055>

Athanasios Nenes  <https://orcid.org/0000-0003-3873-9970>

Sotirios Nikolettseas  <https://orcid.org/0000-0003-3765-5636>

Susetta Finotto  <https://orcid.org/0000-0001-7623-7977>

Spyros N. Pandis  <https://orcid.org/0000-0001-8085-9795>

Angela Simpson  <https://orcid.org/0000-0003-2733-6666>

Maria Staiano  <https://orcid.org/0000-0003-0223-1825>

Antonio Varriale  <https://orcid.org/0000-0003-0985-9877>

Paraskevi Xepapadaki  <https://orcid.org/0000-0001-9204-1923>

Torsten Zuberbier  <https://orcid.org/0000-0002-1466-8875>

Isabella Annesi-Maesano  <https://orcid.org/0000-0002-6340-9300>

REFERENCES

- Moelling K, Broecker F. Air microbiome and pollution: composition and potential effects on human health, including SARS coronavirus infection. *J Environ Public Health*. 2020;2020:1646943.
- Schraufnagel DE, Balmes JR, de Matteis S, et al. Health benefits of air pollution reduction. *Ann Am Thorac Soc*. 2019;16(12):1478-1487.
- Akdis CA, Nadeau KC. Human and planetary health on fire. *Nat Rev Immunol*. 2022;22(11):651-652.
- Vohra K, Vodonos A, Schwartz J, Marais EA, Sulprizio MP, Mickley LJ. Global mortality from outdoor fine particle pollution generated by fossil fuel combustion: results from GEOS-Chem. *Environ Res*. 2021;195:110754.
- Wellbeing, and the changing structure of communities. In: *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*.
- Annesi-Maesano I, Baiz N, Banerjee S, Rudnai P, Rive S, Group S. Indoor air quality and sources in schools and related health effects. *J Toxicol Environ Health B Crit Rev*. 2013;16(8):491-550.
- Akar-Ghbiril N, Phipatanakul W. The indoor environment and childhood asthma. *Curr Allergy Asthma Rep*. 2020;20(9):43.
- Oliveira M, Slezakova K, Delerue-Matos C, Pereira MC, Morais S. Children environmental exposure to particulate matter and polycyclic aromatic hydrocarbons and biomonitoring in school environments: a review on indoor and outdoor exposure levels, major sources and health impacts. *Environ Int*. 2019;124:180-204.
- Zauli-Sajani S, Marchesi S, Boselli G, et al. Effectiveness of a protocol to reduce children's exposure to particulate matter and NO₂ in schools during alert days. *Int J Environ Res Public Health*. 2022;19(17):11019.
- Takahashi Y, Takano K, Suzuki M, et al. Two routes for pollen entering indoors: ventilation and clothes. *J Investig Allergol Clin Immunol*. 2008;18(5):382-388.
- Air quality and health impacts. https://www.researchgate.net/publication/363412159_Air_Quality_and_Health_Impacts
- Sauliene I, Valiulis A, Keriene I, et al. Airborne pollen and fungi indoors: evidence from primary schools in Lithuania. *Heliyon*. 2023;9(1):e12668.
- Why indoor air quality is important to schools. <https://www.epa.gov/iaq-schools/why-indoor-air-quality-important-schools>
- Hooper LG, Kaufman JD. Ambient air pollution and clinical implications for susceptible populations. *Ann Am Thorac Soc*. 2018;15(Suppl 2):S64-S68.
- Hulin M, Simoni M, Viegi G, Annesi-Maesano I. Respiratory health and indoor air pollutants based on quantitative exposure assessments. *Eur Respir J*. 2012;40(4):1033-1045.
- Ahdoot S, Pacheco SE, Council On Environmental H. Global climate change and children's health. *Pediatrics*. 2015;136(5):e1468-e1484.
- Bush A. Impact of early life exposures on respiratory disease. *Paediatr Respir Rev*. 2021;40:24-32.
- de Kluijver J, Evertse CE, Sont JK, et al. Are rhinovirus-induced airway responses in asthma aggravated by chronic allergen exposure? *Am J Respir Crit Care Med*. 2003;168(10):1174-1180.
- Bossios A, Gourgios D, Skevaki CL, et al. Rhinovirus infection and house dust mite exposure synergize in inducing bronchial epithelial cell interleukin-8 release. *Clin Exp Allergy*. 2008;38(10):1615-1626.
- Murray CS, Poletti G, Kebabdz T, et al. Study of modifiable risk factors for asthma exacerbations: virus infection and allergen exposure increase the risk of asthma hospital admissions in children. *Thorax*. 2006;61(5):376-382.
- Akdis CA. Does the epithelial barrier hypothesis explain the increase in allergy, autoimmunity and other chronic conditions? *Nat Rev Immunol*. 2021;21(11):739-751.
- WHO guidelines for indoor air quality. https://www.euro.who.int/_data/assets/pdf_file/0009/128169/e94535.pdf
- Selroos O, Kupczyk M, Kuna P, et al. National and regional asthma programmes in Europe. *Eur Respir Rev*. 2015;24(137):474-483.
- Global atlas of allergy. <https://eaaci-cdn-vod02-prod.azureedge.net/KnowledgeHub/education/books/Global%20Atlas%20of%20Allergy%20-%20English%20Version.pdf>
- Air Pollution. https://www.who.int/health-topics/air-pollution#tab=tab_1
- Wang Z, Walker GW, Muir DCG, Nagatani-Yoshida K. Toward a global understanding of chemical pollution: a first comprehensive analysis of national and regional chemical inventories. *Environ Sci Technol*. 2020;54(5):2575-2584.
- Damialis A, Gilles S, Sofiev M, et al. Higher airborne pollen concentrations correlated with increased SARS-CoV-2 infection rates, as evidenced from 31 countries across the globe. *Proc Natl Acad Sci*. 2021;118(12):e2019034118.
- Copat C, Cristaldi A, Fiore M, et al. The role of air pollution (PM and NO₂) in COVID-19 spread and lethality: a systematic review. *Environ Res*. 2020;191:110129.
- Loreggia A, Passarelli A, Pini MS. The influence of environmental factors on the spread of COVID-19 in Italy. *Procedia Comput Sci*. 2022;207:573-582.
- Damialis A, Traidl-Hoffmann C, Treudler R. Climate change and pollen allergies. 2019. doi:10.1007/978-3-030-02318-8_3

31. Ziska LH, Makra L, Harry SK, et al. Temperature-related changes in airborne allergenic pollen abundance and seasonality across the northern hemisphere: a retrospective data analysis. *Lancet Planet Health*. 2019;3(3):e124-e131. doi:[10.1016/S2542-5196\(19\)30015-4](https://doi.org/10.1016/S2542-5196(19)30015-4)
32. Biagioni B, Annesi-Maesano I, D'Amato G, Cecchi L. The rising of allergic respiratory diseases in a changing world: from climate change to migration. *Expert Rev Respir Med*. 2020;14(10):973-986.
33. Diaz-Sanchez D, Garcia MP, Wang M, Jyrala M, Saxon A. Nasal challenge with diesel exhaust particles can induce sensitization to a neoallergen in the human mucosa. *J Allergy Clin Immunol*. 1999;104(6):1183-1188.
34. Fahy O, Hammad H, Senechal S, et al. Synergistic effect of diesel organic extracts and allergen Der p 1 on the release of chemokines by peripheral blood mononuclear cells from allergic subjects: involvement of the map kinase pathway. *Am J Respir Cell Mol Biol*. 2000;23(2):247-254.
35. Baiz N, Billonnet C, Kirchner S, de Blay F, Annesi-Maesano I. Indoor pet allergen exposures modify the effects of chemical air pollutants on respiratory symptoms. *Int J Tuberc Lung Dis*. 2021;25(5):350-357.
36. Carlsten C, Blomberg A, Pui M, et al. Diesel exhaust augments allergen-induced lower airway inflammation in allergic individuals: a controlled human exposure study. *Thorax*. 2016;71(1):35-44.
37. Rider CF, Yamamoto M, Gunther OP, et al. Controlled diesel exhaust and allergen coexposure modulates microRNA and gene expression in humans: effects on inflammatory lung markers. *J Allergy Clin Immunol*. 2016;138(6):1690-1700.
38. Clifford RL, Jones MJ, MacIsaac JL, et al. Inhalation of diesel exhaust and allergen alters human bronchial epithelium DNA methylation. *J Allergy Clin Immunol*. 2017;139(1):112-121.
39. Mookherjee N, Piyadasa H, Ryu MH, et al. Inhaled diesel exhaust alters the allergen-induced bronchial secretome in humans. *Eur Respir J*. 2018;51(1):1701385.
40. Wang T, Dong H, Jiang W, et al. Viral etiology and atopic characteristics in high-risk asthmatic children hospitalized for lower respiratory tract infection. *Transl Pediatr*. 2020;9(4):541-550.
41. Gilles S, Blume C, Wimmer M, et al. Pollen exposure weakens innate defense against respiratory viruses. *Allergy*. 2020;75(3):576-587.
42. Xepapadaki P, Papadopoulos NG, Bossios A, Manoussakis E, Manousakas T, Saxoni-Papageorgiou P. Duration of postviral airway hyperresponsiveness in children with asthma: effect of atopy. *J Allergy Clin Immunol*. 2005;116(2):299-304.
43. Jackson DJ, Busse WW, Bacharier LB, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol*. 2020;146(1):203-206.e3.
44. Damialis A, Gilles S, Sofiev M, et al. Higher airborne pollen concentrations correlated with increased SARS-CoV-2 infection rates, as evidenced from 31 countries across the globe. *Proc Natl Acad Sci U S A*. 2021;118(12):e2019034118.
45. Choi YJ, Lee KS, Lee YS, Kim KR, Oh JW. Analysis of the association among air pollutants, allergenic pollen, and respiratory virus infection of children in Guri, Korea during recent 5 years. *Allergy Asthma Immunol Res*. 2022;14(3):289-299.
46. Lee SW, Yon DK, James CC, et al. Short-term effects of multiple outdoor environmental factors on risk of asthma exacerbations: age-stratified time-series analysis. *J Allergy Clin Immunol*. 2019;144(6):1542-1550.e1.
47. Chauhan AJ, Inskip HM, Linaker CH, et al. Personal exposure to nitrogen dioxide (NO₂) and the severity of virus-induced asthma in children. *Lancet*. 2003;361(9373):1939-1944.
48. Pfeffer PE, Mudway IS, Grigg J. Air pollution and asthma: mechanisms of harm and considerations for clinical interventions. *Chest*. 2021;159(4):1346-1355.
49. Muller L, Usemann J, Alves MP, Latzin P. Diesel exposure increases susceptibility of primary human nasal epithelial cells to rhinovirus infection. *Physiol Rep*. 2021;9(18):e14994.
50. Pfeffer PE, Ho TR, Mann EH, et al. Urban particulate matter stimulation of human dendritic cells enhances priming of I CD8 T lymphocytes. *Immunology*. 2018;153(4):502-512.
51. Skevaki CL, Tzialta P, Trochoutsou AI, et al. Associations between viral and bacterial potential pathogens in the nasopharynx of children with and without respiratory symptoms. *Pediatr Infect Dis J*. 2015;34(12):1296-1301.
52. Mauderly JL, Samet JM. Is there evidence for synergy among air pollutants in causing health effects? *Environ Health Perspect*. 2009;117(1):1-6.
53. Szabados M, Csako Z, Kotlik B, et al. Indoor air quality and the associated health risk in primary school buildings in Central Europe—the InAirQ study. *Indoor Air*. 2021;31(4):989-1003.
54. Kabirikopaei A, Lau J, Nord J, Bovaird J. Identifying the K-12 classrooms' indoor air quality factors that affect student academic performance. *Sci Total Environ*. 2021;786:147498.
55. de Giuli V, da Pos O, de Carli M. Indoor environmental quality and pupil perception in Italian primary schools. *Build Environ*. 2012;56:335-345.
56. Zaeh SE, Koehler K, Eakin MN, et al. Indoor air quality prior to and following school building renovation in a mid-Atlantic school district. *Int J Environ Res Public Health*. 2021;18(22):12149.
57. Duill FF, Schulz F, Jain A, Krieger L, van Wachem B, Beyrau F. The impact of large mobile air purifiers on aerosol concentration in classrooms and the reduction of airborne transmission of SARS-CoV-2. *Int J Environ Res Public Health*. 2021;18(21):11523.
58. Science M, Caldeira-Kulbakas M, Parekh RS, et al. Effect of wearing a face mask on hand-to-face contact by children in a simulated school environment: the Back-to-school COVID-19 simulation randomized clinical trial. *JAMA Pediatr*. 2022;176(12):1169-1175.
59. Johnston NW, Johnston SL, Duncan JM, et al. The September epidemic of asthma exacerbations in children: a search for etiology. *J Allergy Clin Immunol*. 2005;115(1):132-138.
60. Mortelliti CL, Banzon TM, Vieira CZ, Phipatanakul W. Environmental exposures impact pediatric asthma within the school environment. *Immunol Allergy Clin North Am*. 2022;42(4):743-760.
61. Annesi-Maesano I, Banerjee S, Pham-Thi N. Stress, asthma and allergic sensitisation to house dust mite in children. *Int J Tuberc Lung Dis*. 2021;25(8):675-676.
62. SynAir-G. <https://synairg.eu/>
63. Dai B, Gao C, Xie Y. Flexible wearable devices for intelligent health monitoring. *View*. 2022;3(5):20220027. doi:[10.1002/VIW.20220027](https://doi.org/10.1002/VIW.20220027)
64. Snyder EG, Watkins TH, Solomon PA, et al. The changing paradigm of air pollution monitoring. *Environ Sci Technol*. 2013;47(20):11369-11377. doi:[10.1021/es4022602](https://doi.org/10.1021/es4022602)
65. McLennan S, Fiske A, Tigard D, Müller R, Haddadin S, Buyx A. Embedded ethics: a proposal for integrating ethics into the development of medical AI. *BMC Med Ethics*. 2022;23:6. doi:[10.1186/s12910-022-00746-3](https://doi.org/10.1186/s12910-022-00746-3)
66. Rosario Filho NA, Urrutia-Pereira M, D'Amato G, et al. Air pollution and indoor settings. *World Allergy Organ J*. 2021;14(1):100499.

How to cite this article: Papadopoulos NG, Akdis CA, Akdis M, et al. Addressing adverse synergies between chemical and biological pollutants at schools—The 'SynAir-G' hypothesis. *Allergy*. 2024;79:294-301. doi:[10.1111/all.15857](https://doi.org/10.1111/all.15857)

APPENDIX

The SynAir-G Consortium: Nikolaos G. Papadopoulos, Paraskevi Xepapadaki, Irene Stergiou, Maria Kritikou, Stella Taka, Leonardos Mageiros, Konstantina Pachi, Eleni Stamou, Isabella Annesi-Maesano, Davide Caimmi, Luciana Kase Tanno, Pascal Demoly, Tuomas Jartti, Varpu Elenius, Susetta Finotto, Laura Kropf, Anna Stahlbrost, Abdolrasoul Habibipour, Sotiris Nikolettseas, Gavrilis Filios, PantelisTzamalīs, Christoforos Raptopoulos, Dimitrios Markantonatos, Fotios Kalioras, Panagiotis Kapetanidis, Athanasios Damialis, Antonio Varriale, Sabato D' Auria, Maria Staiano, Angela Pennacchio, Alessandro Capo, Concetta Montagnese, Maia Gotua,

Tamar Abramidze, Maia Rukhadze, Spyros N. Pandis, Eleni Dovrou, Christian Goroncy, Stefan Kelnberger, Valeria Ramiconi, Agata Papotto, Panagiotis Chaslaridis, Angelica Rossi, Torsten Zuberbier, Ingrid van Hofman, Yash Chadha, Ioanna Fergadiotou, Maria Kampa, Makis Kouloubis, Konstantinos Loupos, Tasos Kakouris, Stathis Antoniou, Georgia Priniotaki, Aki Soudunsaari, Niko Järvinen, Pierre Guitton, Pascal Larigaldie, Antonis Hadjiantonis, Alessandro Giusti, Kleanthis Erotokritou, Antonis Hadjantonis, Andreas Papadopoulos, Panayiota Demosthenous, Giuseppina Esposito, Filomena Schena, Marco Randon, Athanasios Nenes, Cezmi Akdis, Mübeccel Akdis, Yagiz Pat, Clare Murray, Angela Simpson.