



**HAL**  
open science

# A non-equilibrium species distribution model reveals unprecedented depth of time lag responses to past environmental change trajectories

Etienne Lalechère, Ronan Marrec, Jonathan Lenoir

## ► To cite this version:

Etienne Lalechère, Ronan Marrec, Jonathan Lenoir. A non-equilibrium species distribution model reveals unprecedented depth of time lag responses to past environmental change trajectories. *Ecology Letters*, 2025, 28 (1), 10.1111/ele.70040 . hal-04909787

**HAL Id: hal-04909787**

**<https://hal.science/hal-04909787v1>**

Submitted on 24 Jan 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title**

2 A non-equilibrium species distribution model reveals unprecedented depth of time lag responses to  
3 past environmental change trajectories.

4

5 **Short running title**

6 Non-equilibrium species distribution modelling

7

8 **Type of article:** Letter

9

10 **Authors**

11 Etienne Lalechère<sup>a1</sup>, Ronan Marrec<sup>b</sup>, Jonathan Lenoir<sup>b</sup>

12 <sup>a</sup> Center for Ecological and Evolutionary Synthesis, Department of Biosciences, University of Oslo,  
13 Norway

14 <sup>b</sup> UMR CNRS 7058 « Ecologie et Dynamique des Systèmes Anthropisés » (EDYSAN), Université  
15 de Picardie Jules Verne, 1 rue des Louvels, 80037, Amiens Cedex, France

16 <sup>1</sup> Corresponding author

17 e-mail addresses of the authors:

18 Etienne Lalechère: [lalechereetienne@gmail.com](mailto:lalechereetienne@gmail.com); +33 3 22 82 77 57

19 Ronan Marrec: [ronan.marrec@u-picardie.fr](mailto:ronan.marrec@u-picardie.fr)

20 Jonathan Lenoir: [jonathan.lenoir@u-picardie.fr](mailto:jonathan.lenoir@u-picardie.fr)

21

22 **Statement of authorship**

23 EL, RM and JL contributed to the conceptualization of the study, EL performed modeling work and

24 output analyzes, EL wrote the first draft of the manuscript and all authors contributed very  
25 significantly to the manuscript edition and revision.

26

### 27 **Data accessibility statement**

28 The data and R code supporting the results are freely available and have been archived in Dryad  
29 (<https://doi.org/10.5061/dryad.cfxpvnvxff>).

30

### 31 **Other information**

32 Number of words in the abstract: 151

33 Number of words in the main text: 5372

34 Number of references: 59

35 Number of figures: 3

36 Submission in the format of another journal: Nature style format

37

### 38 **Abstract**

39 Previous studies have demonstrated legacy effects of current species distributions to past  
40 environmental conditions, but the temporal extent of such time lag dynamics remains unknown. Here,  
41 we have developed a non-equilibrium SDM approach quantifying the temporal extent that must be  
42 taken into account to capture 95 % of the effect that a given time series of past environmental  
43 conditions has on the current distribution of a species. We applied this approach on the distribution  
44 of 92 European forest birds in response to past trajectories of change in forest cover and climate. We  
45 found that non-equilibrium SDMs outperformed traditional SDMs for 95 % of the species. Non-  
46 equilibrium SDMs suggest unprecedented long-lasting effects of past global changes (average time  
47 lag extent ranged from 9 years to 231 years). This framework can help to relax the equilibrium

48 hypothesis of traditional SDMs and to improve future predictions of biodiversity redistribution in  
49 response to global changes.

50

51 **Keywords:** Lagged dynamics – Delayed dynamics – Relaxation time – Colonization credit –  
52 Extinction debt – Climatic debt – Disequilibrium hypothesis – Biodiversity redistribution – Climate  
53 change – Land use change

54

## 55 **Introduction**

56 Lagging dynamics in ecology occur when biodiversity is not in equilibrium with current  
57 environmental conditions, meaning that current species distribution or community composition still  
58 depend on past environmental conditions. The concept of extinction debt was thus defined as a  
59 delayed loss in species number due to a decline in habitat suitability and connectivity (Diamond, 1972;  
60 Tilman et al., 1994). Immigration credit was then used to describe a delayed increase in species  
61 number due to a lack of colonization events towards newly suitable areas (Nagelkerke et al., 2002).  
62 For instance, by relying on empirical observations of past species distributions, recent syntheses on  
63 biodiversity redistribution have demonstrated evidence for extinction debts and immigration credits  
64 at the trailing and leading edge of species distributions, respectively (*e.g.*, Lenoir et al., 2020; Zurell  
65 et al., 2024). The question of whether or not to abandon the concept of equilibrium in ecology has  
66 been raised, due to the overwhelming body of evidence suggesting time-lagged dynamics in  
67 ecological responses to past environmental changes (Coulson, 2021). However, the concept of  
68 equilibrium is still widely used in ecology, especially so by the scientific community working with  
69 species distribution models (SDMs), for which the equilibrium between a species distribution and the  
70 current state of its environmental drivers is the main underlying assumption (Fig. 1C; Essl et al.,  
71 2024). Hence, the assessment of the consequences of environmental changes on biodiversity  
72 redistribution often rely on a questionable assumption.

73 The relevance of legacy effects on current and future biodiversity dynamics has been argued from a  
74 variety of approaches ranging from SDMs (*e.g.*, Dullinger et al., 2012; Rumpf et al., 2019), but also  
75 analyses of time series of community weighted means for specific functional traits (*e.g.*, Bertrand et  
76 al., 2011; Lajeunesse et al., 2022), space-for-time substitutions (Kuussaari et al., 2009), species-area  
77 relationships (*e.g.*, Chen et al., 2017; Ridding et al., 2021), and mechanistic species or community  
78 models (*e.g.*, Aguadé- Gorgorió et al., 2024). Few studies have applied these methods to map the  
79 spatial patterns of the extinction debts and the immigration credits. Some of them provided evidence  
80 of long-lasting legacy effects of anthropogenic pressures aggregated at the country or the biome levels  
81 (Devictor et al., 2012; Dullinger et al., 2013; Liao et al., 2022). Other studies have reported delayed  
82 biodiversity dynamics by paying more attention to their spatial patterns (Bertrand et al., 2016; Chen  
83 et al., 2017; Duchenne et al., 2021; Haddou et al., 2022), which suggest that extinction debts could  
84 mostly be located in areas that do not match with the current network of protected areas (Chen et al.,  
85 2017; Haddou et al., 2022). For instance, Haddou et al. (2022) mapped the distribution of birds across  
86 the conterminous United States in 2016 and compared the dependence of bird distribution on  
87 landscape composition in 2016 and 2001, demonstrating lagging dynamics to past landscape  
88 composition.

89 However, to estimate lagging effects, these studies chiefly relied on just two distant events in time at  
90 a given location (*i.e.*, diachronic analyses with one or two standardized or opportunistic biodiversity  
91 sampling associated with environmental conditions recorded at two dates; *e.g.*, Fig. 1C), thus missing  
92 the complex dynamic that may happen in between and that can only be captured by refined time series  
93 (Daskalova et al., 2020; Liao et al., 2022). First, the detected duration of a time lag depends on the  
94 total duration of the time series available for the environmental variable under consideration. The  
95 starting date of this time series is often defined according to data availability, and is not necessarily  
96 old enough to capture the real temporal extent of a time lag effect (Fig. 1B vs. Fig. 1A). Second, the  
97 direction and magnitude of the effect for a given environmental factor is not sufficient to understand

98 its importance. Indeed, the rate at which the effect decreases over time (*i.e.*, the shape of the time  
99 lagged effect) also indicates if it has a more recent or historic influence (Fig. 1A).

100 To solve the aforementioned knowledge gaps, we developed a non-equilibrium SDM based on  
101 multivariate and temporally-weighted generalized linear regressions to infer a time lag parameter that  
102 captures two important features (Fig. 1A): (i) the time lag extent that must be considered to capture  
103 95 % of the effect that a given time series of past environmental conditions has on the current species  
104 distribution and (ii) the shape of the time-lagged dynamic along the time lag extent. First, to illustrate  
105 the capabilities of our non-equilibrium SDM approach, we performed a simulation exercise that  
106 captures the temporal extent of a lagging response we imposed to a given predictor variable. We then  
107 applied this framework to the current (2013-2017) distribution of 92 European forest birds using long-  
108 term time series of past forest cover (850-2017) and climate (1850-2017). Finally, we illustrated the  
109 ability of our non-equilibrium approach to map the extinction debt and immigration credit by  
110 comparing outputs from our non-equilibrium SDMs against outputs from traditional equilibrium  
111 SDMs. We expect this study to pave the way for the use of non-equilibrium SDMs to better  
112 understand the ability of species to track global changes.

113

## 114 **Materials and methods**

### 115 Biodiversity data

116 For the purpose of our work and to illustrate the relevance of our framework, we focused on forest  
117 bird species. Birds are suitable biological models because they have long been monitored by  
118 ornithologists and now by citizen science and they are highly sensitive to anthropogenic disturbances  
119 (Rosenberg et al., 2019; Keller et al., 2020). Our focus on forest-dwelling species stems from the  
120 large body of evidence that has accumulated in the scientific literature on the delayed effects that both  
121 changes in forest cover and climate conditions may have on forest biodiversity (Bertrand et al., 2011;  
122 Perring et al., 2018; Daskalova et al., 2020; Richard et al., 2021). Birds are fast-dynamic species,

123 which means that the identification of time lags for this taxonomic group is more likely to be  
124 generalized to other taxonomic groups of slower dynamics in terms of life history traits (Daskalova  
125 et al., 2020).

126 We used the European Breeding Bird Atlas (EBBA2) which provides a systematic survey of species  
127 presence-absence data at a spatial resolution of 50 km × 50 km between 2013 and 2017 (median year  
128 = 2015; Keller et al., 2020). Data from European atlases have the advantage of providing exhaustive  
129 and homogenized data on species distribution, which limits potential biases compared to opportunistic  
130 records from presence-only datasets (Keller et al., 2020). We considered the 92 species  
131 (Supplementary Information, Appendix S1) of the EBBA2 whose habitat is classified as either forest  
132 or woodland according to Tobias et al. (2022). The coarse resolution of the EBBA2's mesh size and  
133 the continental extent of the analysis have the advantage of limiting spatial autocorrelation signal due  
134 to dispersal processes on a landscape extent, which were not considered in our analysis.

135

#### 136 Environmental predictors

137 We related species presence-absence data during 2013-2017 to long-term time series of changes in  
138 forest cover and climate conditions, which have been identified as the main determinants of bird  
139 species distributions at continental extent (Cooper et al., 2023). Long-term time series of changes in  
140 forest cover and climate conditions at a yearly resolution were extracted from the Land-Use  
141 Harmonization (LUH2; Hurtt et al., 2020) and the CHELSA Climate (Karger et al., 2017) databases,  
142 respectively. The LUH2 database provides time series of land use since 850 at a spatial resolution of  
143  $0.25^\circ \times 0.25^\circ$ , *i.e.*, approximately 28 km × 28 km at the equator, and at a yearly resolution. We focused  
144 on the land use categories 'primary forest' and 'secondary forest' since they reflect different trends:  
145 an average decrease in the cover of primary forests, mainly to the benefit of secondary forests, and  
146 an average increase in the cover of secondary forests elsewhere in Europe, mostly due to a decline of  
147 agricultural land uses (Hurtt et al., 2020; Supplementary Information, Appendix S2). Forests are

148 defined as ‘primary’ if a land use has remained forest continuously over time since 850, independently  
 149 of management practices. Forests are defined as ‘secondary’ if they have replaced another land use  
 150 type at any time during the same period. Cases of current forests with multiple swinging states  
 151 between forests and agricultural lands are considered ‘secondary’. We used standardized yearly  
 152 averaged maximum temperature (hereafter called temperature) and annual sum of precipitation  
 153 (hereafter called precipitation) since 1850 at a spatial resolution of 30 arcseconds, approximately 1  
 154 km × 1 km at the equator. Given the coarse spatial resolution and the continental extent of the atlas  
 155 data, the predictors were projected in the same coordinate reference system (ETRS89-LAEA) and  
 156 aggregated, using the mean, at a spatial resolution of 50 km × 50 km to match the EBBA2’s mesh  
 157 size. Because the period covered by the EBBA2 is a 5-year period (2013-2017), we used a 5-year  
 158 sliding window to average forest cover and climate conditions backward from 2015 (average year of  
 159 the EBBA2 dataset) and at a yearly resolution until 852 AD for forest cover and until 1852 for climate  
 160 conditions. There were no multicollinearity issues between the averaged predictor variables  
 161 considering that the absolute Spearman correlation coefficient was systematically < 0.7 (Dormann et  
 162 al., 2013).

163

#### 164 Non-equilibrium species distribution models

165 Inspired by previous works in spatial ecology (Chandler et al., 2016; Miguet et al., 2017), we  
 166 developed a multivariate and temporally-weighted generalized linear regression to model the  
 167 response variable, that is here a binary variable (presence-absence of a given species), as a function  
 168 of current and past conditions (Eq. 1):

$$169 \quad \mu_i = \alpha + \sum_{j=1}^N \beta_j \sum_{\Delta t=0}^{\max(\Delta t)} X_{ij\Delta t} W_j(\Delta t) + \varepsilon_i \text{ (Eq. 1)}$$

170 where  $\mu_i$  refers to each observation  $i$  of the response variable in the model formula,  $\alpha$  is the intercept  
 171 parameter and  $\beta_j$  is the slope estimate of the predictor variable  $j$  out of the  $N$  covariates included as



172 predictor variables in the model formula. The residual error term  $\varepsilon$  is assumed to be randomly  
173 distributed. The slope estimate  $\beta_j$  depends on a temporal weighting function  $W_j = e^{-0.5 \left(\frac{\Delta t}{\gamma_j \cdot \max(\Delta t)}\right)^2}$ ,  
174 which is used to weight the covariate value  $X_{ij\Delta t}$  according to its age  $\Delta t$ . For the year 2015,  $\Delta t = 0$   
175 and it increases as we go back in time. The older the date  $t$  for the covariate  $X_{ij}$ , the larger the  $\Delta t$  and  
176 the lower is the weight assigned to  $X_{ij\Delta t}$ . For instance, if we consider the covariate of primary forest  
177 cover, the older date at which the weight is the lowest is 852 which corresponds to an age of 1,163  
178 years. Thus, we assumed that the effect of a covariate decreases as we go back in time. To test the  
179 sensitivity of the results to the definition of the weighting function, we also considered an alternative  
180 hypothesis of a negative exponential decrease of the effects of the covariates with  $W_j = e^{-\frac{\Delta t}{\gamma_j \cdot \max(\Delta t)}}$ .  
181 We defined ‘the shape of the time lag’ as the shape of the weighting function that depends on the time  
182 lag parameter  $\gamma_j$ , which describes how the effect of a covariate  $j$  decreases, linearly or non-linearly  
183 as we go back in time. The time lag parameter corresponds to the year at which approximately one  
184 third (63.2 %) or one quarter (75.8 %) of the effect of the covariate has occurred over time, for a  
185 negative exponential and a half-Gaussian weighting function, respectively. We also used these  
186 functions to define the time-window that should be considered to account for 95 % of the effect that  
187 a given time series of past environmental condition has on the current distribution of a species, when  
188 summing all weighted values of that focal predictor before inferring its effect size (called ‘the time  
189 lag extent’, Fig. 1A). We applied this framework, using a binomial family with a logit link function,  
190 to relate the presence-absence data of a species to primary forest cover, secondary forest cover,  
191 temperature and precipitation altogether in one single multivariate model. All the analyses were  
192 performed using the R software and the *scalescape* R package for parameter inference (Lowe et al.,  
193 2022). The *scalescape* package was initially designed to fit spatially-weighted generalized linear  
194 regressions from ‘landscape matrices’ which include predictor values for different distance classes.  
195 Thus, the analysis of the bidimensional (*i.e.*, latitude and longitude) spatial context is reduced to a

196 one-dimensional problem, since cumulated landscape features are taken into account according to  
197 their distance only, but not their direction, from the focal plot where the response variable is sampled.  
198 Here, we replaced the ‘landscape matrices’ by ‘temporal matrices’ including the predictor values for  
199 each  $\Delta t$ , instead of distance classes, in order to fit the temporally-weighted generalized linear  
200 regressions. We used the maximum possible temporal extent of the predictors in order to maximize  
201 the possibility of correctly estimating the temporal extent of each time lag. We did not interpret time  
202 lag parameters leading to temporal extents that were larger than or equal to the maximum possible  
203 temporal extent available for the predictors (*i.e.*, 1,163 years for primary and secondary forest  
204 cover). We performed a simulation exercise to illustrate the capabilities of our non-equilibrium SDM  
205 approach to capture the temporal extent of a lagging response to a given predictor variable for which  
206 we imposed and thus know the magnitude of the lagging dynamic (Supplementary Information,  
207 Appendix S3).

208

#### 209 Model selection, validation and predictions

210 We compared the Akaike Information Criterion (AIC) of our non-equilibrium SDMs with an  
211 alternative equilibrium model relying on traditional SDM approaches and for which the predictors  
212 are basically averaged over the 5-year time-window of the EBBA2, namely 2013-2017. Models were  
213 considered significantly different if the AIC difference was  $> 2$  (Burnham & Anderson, 2002). The  
214 comparison between the equilibrium SDMs and the non-equilibrium ones were not very sensitive to  
215 the definition of the weighting functions. Both the Gaussian and the negative exponential weighting  
216 functions of the non-equilibrium SDMs outperformed the equilibrium SDMs for the majority of the  
217 studied bird species (87 and 79 out of 92 species, respectively). For further analyses, we retained only  
218 the Gaussian non-equilibrium SDMs given that it leads to a better fit compared to the negative  
219 exponential non-equilibrium SDMs for the vast majority of the species (for 82 out of 92 species).

220 The validity of the tested models depends on the typical assumptions underlying SDMs (Zurell et al,

221 2020). Obviously, the hypothesis of species-environment equilibrium is expected under traditional  
222 equilibrium SDMs, while the non-equilibrium SDM relaxes this assumption. For both equilibrium  
223 and non-equilibrium SDMs, observations are assumed to be unbiased and independent. The predictors  
224 are assumed to be free of errors and sufficient to capture the main determinants of European bird  
225 distributions given previous results (Cooper et al., 2023).

226 The residual errors were calculated, applying the *resid()* function from the *base* R package. For each  
227 species, model validity was assessed from the distribution of the residual errors, which was assumed  
228 to be Gaussian and centered on zero (tested from Student's *t* test). We tested for positive or negative  
229 spatial autocorrelation signals in the pooled residuals using the Moran's *I* index, calculated for  
230 uniformly distributed distance classes with 500 km increments, using the *correlog* function from the  
231 *ncf* R package (Bjornstad et al., 2001). We expected that the distribution of this index would be  
232 centered on 0 (no spatial autocorrelation) and tested this using a Student's *t* test. The distribution of  
233 the pooled residual errors, and the pooled Moran's *I*, among all the species are shown in the  
234 Supplementary Information (Appendix S4).

235 The predictive ability of the best model was evaluated using the Area Under the Curve (AUC), the  
236 specificity and the sensitivity indices, averaged over 20 replicates of a bootstrap cross-validation  
237 procedure (70 % and 30 % partitioning for calibration and validation, respectively; Fielding & Bell,  
238 1997). We have not considered a clustered selection of cross-validated samples given that there is no  
239 extrapolation thanks to the use a European atlas with an exhaustive sampling grid and given the  
240 absence of future projections (Wenger & Olden, 2012). Species distribution models with an AUC <  
241 0.7 were excluded from further analyses to prevent inaccurate predictions. For each predictor variable,  
242 we compared the estimates of the equilibrium SDMs against the estimates of the non-equilibrium  
243 SDMs to identify potential discrepancies in direction, magnitude (using Student's *t* test) and  
244 variability (using Fischer's *F* test).

245 We converted presence probabilities into binary presence-absence predictions using the maximum

246 true skill statistic (TSS) as a threshold adapted for presence-absence data (Liu et al., 2013). To map  
247 the extinction debt and the immigration credit of forest birds across Europe, we followed Haddou et  
248 al. (2022) who subtracted the effective number of species as predicted by stacking all species binary  
249 maps based on the non-equilibrium SDMs from that obtained by the equilibrium SDMs.

250

## 251 **Results**

252 We found that non-equilibrium SDMs had the lowest Akaike Information Criterion (AIC) for 95 %  
253 of the species (87 out of 92 species) confirming that equilibrium SDMs are less relevant than non-  
254 equilibrium ones, in most cases. Analyzing the effects of long-term yearly time series of land use and  
255 climate (Supplementary Information, Appendix S2), we found that time lag extents were extremely  
256 variable (Fig. 2) and were larger for changes in primary forests (mean = 231 years, SD = 229 years)  
257 compared to changes in secondary forests (mean = 137 years, SD = 190 years), and for changes in  
258 precipitation (mean = 20 years, SD = 29 years) compared to changes in temperature regimes (mean  
259 = 9 years, SD = 28 years). The shape of the temporal weighting function used in the model formula  
260 (*i.e.*, half-Gaussian compared to exponential negative as we go back in time) underlined a slower  
261 decrease of the weight attributed to environmental conditions towards the onset of the temporal extent  
262 covered by the lagging dynamic.

263 The non-equilibrium SDMs were valid, considering that species' residual errors followed a normal  
264 distribution centered on zero (Student's  $t$  p-value > 0.05; Supplementary Information, Appendix S4)  
265 and that the spatial autocorrelation signal of the species' residual errors in the correlogram of the  
266 Moran's  $I$  index was also centered on zero (Student's  $t$  p-value > 0.05; Supplementary Information,  
267 Appendix S4). The cross-validated Area Under the Curve (AUC) index indicated a good overall  
268 predictive accuracy (mean = 0.82, SD = 0.06) after excluding three species with low AUC values  
269 (AUC < 0.7). Overall, the non-equilibrium SDMs tended to better predict presences than absences  
270 (mean sensitivity = 0.83, SD = 0.06; mean specificity = 0.71, SD = 0.07).

271 We found some discrepancies between equilibrium and non-equilibrium SDMs when comparing the  
272 direction, magnitude and variance of the slope estimates for each predictor variable (Supplementary  
273 Information, Appendix S4). For 26 % of the studied species (27 out of 87 species), non-equilibrium  
274 SDMs indicated a positive effect of secondary forests while it was the opposite (*i.e.*, a negative effect  
275 of secondary forests) when relying on equilibrium SDMs. For the other predictor variables, the  
276 discrepancies on the direction of the effect between equilibrium and non-equilibrium SDMs  
277 concerned fewer species ( $\leq 6\%$ ). We found a larger effect size and a larger variability in the slope  
278 estimates of primary forests when considering equilibrium SDMs, while the opposite was true for  
279 secondary forests (Student's  $t$  test  $\leq 0.05$ ; Fischer's  $F$  test  $\leq 0.05$ ; Supplementary Information,  
280 Appendix S4).

281 We found evidence for extinction debts to be particularly pronounced in primary boreal forests, near  
282 the Carpathians, and north of the Black Sea while evidence for immigration credits were mostly  
283 pronounced in secondary forests in the Mediterranean part of the Alps and Pyrenees, as well as in the  
284 Caucasus and in the mountainous areas north of the Caspian Sea (Fig. 3; Supplementary Information,  
285 Appendix S2).

286

## 287 **Discussion**

288 Overall, we found less support for traditional species distribution models (SDMs) that rely on the  
289 equilibrium hypothesis and which showed lower fit than non-equilibrium SDMs and this was true for  
290 95 % of the 92 European forest bird species we studied. In line with former studies focusing on the  
291 fate of avian biodiversity in the context of global changes (*e.g.*, Devictor et al., 2012; Haddou et al.,  
292 2022), our result advocates for the abandonment of the equilibrium assumption in SDMs. Here, we  
293 have reported unprecedented long-lasting effects of global changes on current avian biodiversity  
294 distribution (up to 246 years for forest cover change) compared to previous studies (Figueiredo et al.,  
295 2019; Daskalova et al., 2020; Liao et al., 2022), even when considering potential sources of

296 inaccuracies (Essl et al., 2024). The shape of the time-lag response curve (*i.e.*, the shape of the  
297 temporal weighting function being half-Gaussian or exponential negative) along a given predictor  
298 variable also underlined the long-lasting effect of ancient environmental conditions. Because of that,  
299 the use of equilibrium SDMs can lead to biased direction, magnitude and variability in the coefficient  
300 estimates of the underlying predictor variables. The persistent use of models relying on the  
301 equilibrium hypothesis in the SDM literature is unlikely to be explained solely by a lack of available  
302 datasets, such as long-term time series of environmental conditions, because strong evidence of non-  
303 equilibrium biodiversity dynamics has already been proven and generalized from the currently  
304 available datasets, at different spatiotemporal scales (Figueiredo et al., 2019). We rather suggest that  
305 the poor use of models relying on the non-equilibrium hypothesis is more likely due to a lack of  
306 generic and ready-to-use modelling frameworks, something we now provide here, but also to an  
307 underestimation of the importance and implications of the temporal extent behind time lags.

308

309 The low consideration for time lag dynamics in SDMs, despite recent literature (Essl et al. 2024),  
310 might also be explained by the lack of a common definition of a time lag and how to measure it.  
311 Indeed, time lags can either refer to the time elapsed since the occurrence of a causative perturbation  
312 until the time it reaches a peak in population responses (Daskalova et al., 2020) or to the time elapsed  
313 for a debt to be completely paid off or for a credit to be completely spent (Figueiredo et al., 2019), or  
314 even to the time span elapsed between two dates in diachronic analyses (Haddou et al., 2020).  
315 However, time-lagged responses of biodiversity to environmental change trajectories were initially  
316 associated with the so-called ‘relaxation time’ which has a clear definition: the complete time window  
317 during which a disturbance has consequences on biodiversity until it returns to a new equilibrium  
318 state (Diamond, 1972; Tilman, 1995). References to the concept of relaxation time remain scarce in  
319 the scientific literature (*e.g.*, Lalechère et al., 2018), compared to the use of the term time lag, perhaps  
320 because a return to an equilibrium state is more theoretical in the context of continuous global changes,

321 and is rarely verified in the natural world (Essl et al., 2024). Finally, the concept of time lag remains  
322 ambiguous, although it is more commonly used than the concept of relaxation time, which has a clear  
323 definition but lacks empirical evidence. As a consequence, we propose to use the temporal extent of  
324 a time lag to accurately define its duration, abandoning the concepts of equilibrium and relaxation  
325 time, given the temporal non-interruption of global changes during the Anthropocene.

326

327 Our approach offers the possibility to assess the extent of a time lag associated with each individual  
328 predictor variable involved in explaining the contemporary distribution of a given species. For  
329 instance, we found that the distribution of European birds is strongly dependent on the historical  
330 extent of primary forests, whose decline takes longer to be paid off compared to the ongoing  
331 immigration credit to be spent following the historical increase in secondary forests. This result  
332 highlights that interacting land use changes threaten the resilience of communities, considering that  
333 some species preferentially present in primary forests are not adapted to colonize secondary forests  
334 that can have different compositional and structural characteristics (*e.g.*, Sabatini et al., 2018;  
335 Lalechère et al., 2024). The long-lasting effects of global changes that we have found here advocate  
336 for reconsidering the debate on their relative importance taking into account the dependance of  
337 species to past environmental conditions (Lalechère et al., 2022; Thompson et al., 2023). We found  
338 larger temporal extents of time lags for precipitation compared to temperature. Time lags related to  
339 precipitation is rarely considered in the scientific literature. As a consequence, previously estimated  
340 time-lagged consequences of climate change may have been underestimated because most studies use  
341 the community temperature index to estimate time lags, while some results also highlight the  
342 importance of precipitation in habitat suitability (Devictor et al., 2012; Tehrani et al., 2021;  
343 Lajeunesse et al., 2023).

344

345 Time lags can be artificially created by imperfect data, such as those caused by an inaccurate matching

346 between species observations, records in a database, environmental conditions and their averaging,  
347 especially when derived from model projections rather than from *in situ* measurements (Essl et al.,  
348 2024). Here, we used land use projections from an Earth System Model that was harmonized for long-  
349 term studies (LUH2; Hurtt et al., 2020). The inherent limitations (*e.g.*, Goldewijk & Verburg, 2013)  
350 due to the coarse spatial resolution and the coarse definition of land-use classes, defined in global  
351 earth system and climatic models, may potentially affect the inferred parameters and the predictions  
352 we made in our study. Then, some inaccuracies and uncertainties can be associated to the inferred  
353 time lag parameters (Supplementary Information, Appendix S3). However, these limitations are  
354 unlikely to bias the relative differences we found between equilibrium and non-equilibrium SDMs.  
355 The extent of the time lags we report here is disproportionated (several decades or centuries)  
356 compared to the potential biases that may be introduced by these different sources of inaccuracy,  
357 excepting for temperature with time lags that are mostly of the order of several years. In this case,  
358 time lags that are very close to the lower or the upper limit of  $\Delta t$  most likely indicates that the  
359 temporal extent of the temperature time series is too small for a proper inference of the lagging effect  
360 attributed to temperature changes (Arroyo-Rodríguez et al., 2022).

361

362 When compared with outputs from equilibrium SDMs, predictions from non-equilibrium SDMs  
363 allow us to assess the extinction debt and immigration credit (Haddou et al., 2022). We have shown  
364 that European bird communities are mostly affected by an extinction debt and not so much by an  
365 immigration credit. The extinction debt is particularly pronounced in boreal forests, which is  
366 consistent with the strong decline of primary forests therein. Overall, the immigration credit is mostly  
367 located in Mediterranean forests, the Caucasus and the mountainous areas north of the Caspian Sea,  
368 where secondary forests are increasing in cover. The presence of an extinction debt in some areas  
369 where secondary forests increase indicates an interaction between the progressive colonization of  
370 secondary forests and an ongoing decline of species with a low thermal tolerance due to increasing



371 temperature (Jiguet et al., 2010). These complex antagonistic interactions may have complicated the  
372 inference of the time lags related to temperature in areas with both delayed colonization and extinction  
373 dynamics.

374

375 Few other studies have ever attempted to map the extinction debt and the immigration credit of  
376 biodiversity distribution (but see, Bertrand et al., 2016; Chen et al., 2017; Duchenne et al., 2021;  
377 Haddou et al., 2022). For example, Chen et al. (2017) mapped the delayed extinction patterns of entire  
378 communities at the global extent, analyzing their dependence to forest areas in a diachronic analysis  
379 that compared the 2000s against the 16th century. They showed that delayed extinctions did not  
380 overlap with hotspots of species richness or high extinction-risk areas based on IUCN threatened  
381 status. Focusing on avian biodiversity, Haddou et al. (2022) have mapped changes that have yet to be  
382 realized over the conterminous United States, pointing out areas of conservation concern. However,  
383 they recognize that non-equilibrium spatial patterns could be obtained by considering multiple time  
384 points over a longer time period in order to improve projections of past, current and future biodiversity.  
385 Our conceptual and modelling framework paves the way to fill this gap and to anticipate future  
386 projections of non-equilibrium biodiversity patterns using shared socio-economic pathways and land  
387 use projections. Indeed, the inference procedure in our non-equilibrium SDM is not based on a  
388 specific year or a specific temporal baseline period, but it accounts for time-series of past  
389 environmental conditions. Note that it can be also applied to time-series of biodiversity monitoring  
390 data to account for temporal variation in species distribution parameters in order to anticipate changes  
391 in time lags and species affinity to future environmental conditions.

392

393 After establishing a proof of concept, one of the main perspectives of this study is to refine how  
394 species do or do not depend on past environmental conditions according to the model selection  
395 procedure. For example, this procedure could be fine-tuned considering alternative hypotheses about

396 the decrease of the effect size, in order to better model the dependence of current species to past  
397 environmental conditions. This can be especially true for temperature for which the temporal extent  
398 of time-lag effects is mostly shallow and vary little among species, while the literature indicates a  
399 strong variability according to species thermal tolerances (Jiguet et al., 2010). Model selection  
400 procedure can also be used to test if additional and potentially missing predictors could be identified  
401 to improve the balance of the predictive accuracy between presences and absences. However, the  
402 accessibility of long-term time series on environmental conditions is still very limited beyond land  
403 use and climate. Therefore, it would be necessary to adopt a mixed approach, by considering an  
404 equilibrium as well as a non-equilibrium hypothesis, depending on data accessibility of each predictor  
405 variable and not only based on the ecological significance. This can be easily done within the  
406 modelling framework we provide here by simply adding predictors that are invariant over time  
407 directly in Eq. 1.

408 A second perspective is to test whether it is possible or not to identify the functional characteristics  
409 of the species which have contrasted time lag extents and which do not respond in the same way to  
410 the model selection procedure (Tobias et al., 2022). For example, we can expect that species with a  
411 short life expectancy or able to disperse at long distances are less sensible to past environmental  
412 conditions (Kitzes & Harte, 2015; Lalechère et al., 2017; Lalechère et al., 2019; Daskalova et al.,  
413 2020; Couet et al., 2022). We could also expect time lags to be related to the trophic position of the  
414 species and impacted by cascading effects of cumulated time lags throughout the trophic food web,  
415 from primary producers to consumers, for example (Essl et al., 2015).

416 A third perspective is to improve our non-equilibrium modelling framework with the integration of  
417 species dispersal kernels, or the spatial extent of effect of each individual drivers of biodiversity, that  
418 are two ways to account for spatial processes (Barbet-Massin et al., 2012; Miguet et al., 2017). In this  
419 study, the coarse spatial resolution of the bird's atlas dataset had the advantage to limit spatial  
420 autocorrelation issues due to small-scale variation in environmental conditions. However, this coarse

421 spatial resolution can partly affect inference and predictions, by overlooking local microclimatic  
422 variation due to the buffering effect of vegetation, for example, which might also explain the small  
423 extent and variability of the time lags we found for temperature (Kim et al., 2022). Given that our  
424 approach is inspired by the concept of the spatial scale of effect, a logical follow-up would be to  
425 model spatiotemporal response curves to account for a decrease in the effect of a driver the further  
426 we go away, in space or time, from a given biodiversity survey. This perspective could pave the way  
427 to the inference of spatial and temporal extent of the effects, providing a common framework for  
428 multi-taxa analyses including species which do not respond to global changes at the same extents  
429 (Sandel, 2015).

430

431 To conclude, we contribute to fill the gap of a conceptual and modelling framework adapted to study  
432 non-equilibrium species distribution in the context of gradual global changes. We propose to model  
433 the temporal extent of time lags that account for the full effects (95 %) of past global changes on  
434 current biodiversity. We found that it can reach several centuries, that is much more than previously  
435 reported in the scientific literature. Our non-equilibrium SDM approach revealed the spatial patterns  
436 of the mismatch of the species-environment relationships at large biogeographical extent. This  
437 approach, which challenges the concept of equilibrium, also has the important implication of making  
438 it unnecessary to define an arbitrary baseline time period to project the future distribution of species.

439

#### 440 **Acknowledgments**

441 We would like to thank Ingolf Kühn for his friendly proofreading, which greatly improved the content  
442 of this study. We also thank Torbjørn Ergon for his important feedbacks. We also thank the two  
443 reviewers for their ideas and comments, which greatly helped to improve this work. This has received  
444 funding from the European Union's Horizon Europe research and innovation programme under the  
445 Marie Skłodowska-Curie grant agreement No. 101126636.

446

447 **References**

448 Aguadé-Gorgorió, G., Arnoldi, J. F., Barbier, M., & Kéfi, S. (2024). A taxonomy of multiple stable  
449 states in complex ecological communities. *Ecology Letters*, 27(4), e14413.

450 Arroyo-Rodríguez, V., Martínez-Ruiz, M., Bezerra, J. S., Galán-Acedo, C., San-José, M., & Fahrig,  
451 L. (2023). Does a species' mobility determine the scale at which it is influenced by the surrounding  
452 landscape pattern?. *Current Landscape Ecology Reports*, 8(1), 23-33.

453 Barbet-Massin, M., Thuiller, W., & Jiguet, F. (2012). The fate of European breeding birds under  
454 climate, land-use and dispersal scenarios. *Global Change Biology*, 18(3), 881-890.

455 Bertrand, R., Lenoir, J., Piedallu, C., Riofrío-Dillon, G., De Ruffray, P., Vidal, C., ... & Gégout, J. C.  
456 (2011). Changes in plant community composition lag behind climate warming in lowland forests.  
457 *Nature*, 479(7374), 517-520.

458 Bertrand, R., Riofrío-Dillon, G., Lenoir, J., Drapier, J., De Ruffray, P., Gégout, J. C., & Loreau, M.  
459 (2016). Ecological constraints increase the climatic debt in forests. *Nature Communications*, 7(1),  
460 12643.

461 Bjornstad, O. N., & Falck, W. (2001). Nonparametric spatial covariance functions: estimation and  
462 testing. *Environmental and Ecological Statistics*, 8, 53-70.

463 Burnham, K. P., & Anderson, D. R. (2002). Model selection and multimodel inference: a practical  
464 information-theoretic approach 2nd edition. Springer, New York.

465 Chandler, R., & Hepinstall-Cymerman, J. (2016). Estimating the spatial scales of landscape effects  
466 on abundance. *Landscape Ecology*, 31, 1383-1394.

467 Chen, Y., & Peng, S. (2017). Evidence and mapping of extinction debts for global forest-dwelling  
468 reptiles, amphibians and mammals. *Scientific Reports*, 7(1), 44305.

469 Couet, J., Marjakangas, E. L., Santangeli, A., Kålås, J. A., Lindström, Å., & Lehikoinen, A. (2022).

470 Short-lived species move uphill faster under climate change. *Oecologia*, 198(4), 877-888.

471 Cooper, W. J., McShea, W. J., Songer, M., Huang, Q., & Luther, D. A. (2023). Harmonizing spatial  
472 scales and ecological theories to predict avian richness and functional diversity within forest  
473 ecosystems. *Proceedings of the Royal Society B*, 290(2001), 20230742.

474 Coulson, T. (2021). We live in a changing world, but that shouldn't mean we abandon the concept of  
475 equilibrium. *Ecology Letters*, 24(1), 3-5.

476 Daskalova, G. N., Myers-Smith, I. H., Bjorkman, A. D., Blowes, S. A., Supp, S. R., Magurran, A. E.,  
477 & Dornelas, M. (2020). Landscape-scale forest loss as a catalyst of population and biodiversity  
478 change. *Science*, 368(6497), 1341-1347.

479 Devictor, V., Van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J., ... & Jiguet, F.  
480 (2012). Differences in the climatic debts of birds and butterflies at a continental scale. *Nature climate  
481 change*, 2(2), 121-124.

482 Diamond, J. M. (1972). Biogeographic kinetics: estimation of relaxation times for avifaunas of  
483 southwest Pacific islands. *Proceedings of the National Academy of Sciences*, 69(11), 3199-3203.

484 Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., ... & Lautenbach, S. (2013).  
485 Collinearity: a review of methods to deal with it and a simulation study evaluating their performance.  
486 *Ecography*, 36(1), 27-46.

487 Duchenne, F., Martin, G., & Porcher, E. (2021). European plants lagging behind climate change pay  
488 a climatic debt in the North, but are favoured in the South. *Ecology Letters*, 24(6), 1178-1186.

489 Dullinger, S., Gatttringer, A., Thuiller, W., Moser, D., Zimmermann, N. E., Guisan, A., ... & Hülber,  
490 K. (2012). Extinction debt of high-mountain plants under twenty-first-century climate change. *Nature  
491 climate change*, 2(8), 619-622.

492 Dullinger, S., Essl, F., Rabitsch, W., Erb, K. H., Gingrich, S., Haberl, H., ... & Hulme, P. E. (2013).  
493 Europe's other debt crisis caused by the long legacy of future extinctions. *Proceedings of the National*

494 *Academy of Sciences*, 110(18), 7342-7347.

495 Dullinger, S., Essl, F., Rabitsch, W., Erb, K. H., Gingrich, S., Haberl, H., ... & Hulme, P. E. (2013).  
496 Europe's other debt crisis caused by the long legacy of future extinctions. *Proceedings of the National*  
497 *Academy of Sciences*, 110(18), 7342-7347.

498 Essl, F., Dullinger, S., Rabitsch, W., Hulme, P. E., Pyšek, P., Wilson, J. R., & Richardson, D. M.  
499 (2015). Historical legacies accumulate to shape future biodiversity in an era of rapid global change.  
500 *Diversity and Distributions*, 21(5), 534-547.

501 Essl, F., García-Rodríguez, A., Lenzner, B., Alexander, J. M., Capinha, C., Gaüzère, P., ... &  
502 Dullinger, S. (2024). Potential sources of time lags in calibrating species distribution models. *Journal*  
503 *of Biogeography*, 51(1), 89-102.

504 Fielding, A. H., & Bell, J. F. (1997). A review of methods for the assessment of prediction errors in  
505 conservation presence/absence models. *Environmental Conservation*, 24(1), 38-49.

506 Figueiredo, L., Krauss, J., Steffan-Dewenter, I., & Sarmiento Cabral, J. (2019). Understanding  
507 extinction debts: spatio-temporal scales, mechanisms and a roadmap for future research. *Ecography*,  
508 42(12), 1973-1990.

509 Goldewijk, K., & Verburg, P. H. (2013). Uncertainties in global-scale reconstructions of historical  
510 land use: an illustration using the HYDE data set. *Landscape Ecology*, 28, 861-877.

511 Haddou, Y., Mancy, R., Matthiopoulos, J., Spatharis, S., & Dominoni, D. M. (2022). Widespread  
512 extinction debts and colonization credits in United States breeding bird communities. *Nature Ecology*  
513 *& Evolution*, 6(3), 324-331.

514 Hurtt, G. C., Chini, L., Sahajpal, R., Frohking, S., Bodirsky, B. L., Calvin, K., ... & Zhang, X. (2020).  
515 Harmonization of global land use change and management for the period 850–2100 (LUH2) for  
516 CMIP6. *Geoscientific Model Development*, 13(11), 5425-5464.

517 Jiguet, F., Gregory, R. D., Devictor, V., Green, R. E., Voríšek, P., Van Strien, A., & Couvet, D.

518 (2010). Population trends of European common birds are predicted by characteristics of their climatic  
519 niche. *Global Change Biology*, 16(2), 497-505.

520 Johnson, T. F., Beckerman, A. P., Childs, D. Z., Webb, T. J., Evans, K. L., Griffiths, C. A., ... &  
521 Freckleton, R. P. (2024). Revealing uncertainty in the status of biodiversity change. *Nature*, 1-7.

522 Karger, D. N., Conrad, O., Böhrner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... & Kessler, M.  
523 (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, 4(1), 1-20.

524 Karger, D., Schmatz, D. R., Dettling, G., & Zimmermann, N. E. (2019). High resolution monthly  
525 precipitation and temperature timeseries for the period 2006-2100. *arXiv*, arXiv-1912.

526 Keller, V., Herrando, S., Vorišek, P., Franch, M., Kipson, M., Milanese, P., ... & Foppen, R. P. B.  
527 (2020). European breeding bird atlas 2: distribution, abundance and change.

528 Kim, H., McComb, B. C., Frey, S. J., Bell, D. M., & Betts, M. G. (2022). Forest microclimate and  
529 composition mediate long-term trends of breeding bird populations. *Global Change Biology*, 28(21),  
530 6180-6193.

531 Kitzes, J., & Harte, J. (2015). Predicting extinction debt from community patterns. *Ecology*, 96(8),  
532 2127-2136.

533 Kuussaari, M., Bommarco, R., Heikkinen, R. K., Helm, A., Krauss, J., Lindborg, R., ... & Steffan-  
534 Dewenter, I. (2009). Extinction debt: a challenge for biodiversity conservation. *Trends in ecology &*  
535 *evolution*, 24(10), 564-571.

536 Lajeunesse, A., & Fourcade, Y. (2023). Temporal analysis of GBIF data reveals the restructuring of  
537 communities following climate change. *Journal of Animal Ecology*, 92(2), 391-402.

538 Lalechère, E., Jabot, F., Archaux, F., & Deffuant, G. (2017). Non-equilibrium plant metapopulation  
539 dynamics challenge the concept of ancient/recent forest species. *Ecological modelling*, 366, 48-57.

540 Lalechère, E., Jabot, F., Archaux, F., & Deffuant, G. (2018). Projected regional forest plant  
541 community dynamics evidence centuries-long effects of habitat turnover. *Journal of Vegetation*

542 *Science*, 29(3), 480-490.

543 Lalechère, E., Archaux, F., & Jabot, F. (2019). Relative importance of landscape and species  
544 characteristics on extinction debt, immigration credit and relaxation time after habitat turnover.  
545 *Population Ecology*, 61(4), 383-395.

546 Lalechère, E., & Bergès, L. (2022). Importance of climate-induced tree species composition changes  
547 in forecasting the amount of reachable habitat for forest birds. *Diversity and Distributions*, 28(9),  
548 1781-1794.

549 Lalechère, E., Monnet, J. M., Breen, J., & Fuhr, M. (2024). Assessing the potential of remote sensing-  
550 based models to predict old-growth forests on large spatiotemporal scales. *Journal of Environmental*  
551 *Management*, 351, 119865.

552 Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J., & Grenouillet, G. (2020).  
553 Species better track climate warming in the oceans than on land. *Nature Ecology & Evolution*, 4(8),  
554 1044-1059.

555 Liao, Z., Peng, S., & Chen, Y. (2022). Half-millennium evidence suggests that extinction debts of  
556 global vertebrates started in the Second Industrial Revolution. *Communications Biology*, 5(1), 1311.

557 Liu, C., White, M., & Newell, G. (2013). Selecting thresholds for the prediction of species occurrence  
558 with presence-only data. *Journal of Biogeography*, 40(4), 778-789.

559 Lowe, E. B., Iuliano, B., Gratton, C., & Ives, A. R. (2022). ‘Scalescape’: an R package for estimating  
560 distance-weighted landscape effects on an environmental response. *Landscape Ecology*, 37(7), 1771-  
561 1785.

562 Miguet, P., Fahrig, L., & Lavigne, C. (2017). How to quantify a distance-dependent landscape effect  
563 on a biological response. *Methods in Ecology and Evolution*, 8(12), 1717-1724.

564 Nagelkerke, K. C., Verboom, J., Van Den Bosch, F., & Van De Wolfshaar, K. (2002). Time lags in  
565 metapopulation responses to landscape change. In *Applying landscape ecology in biological*



566 conservation (pp. 330-354). Springer, New York.

567 Perring, M. P., Bernhardt-Römermann, M., Baeten, L., Midolo, G., Blondeel, H., Depauw, L., ... &  
568 Verheyen, K. (2018). Global environmental change effects on plant community composition  
569 trajectories depend upon management legacies. *Global Change Biology*, 24(4), 1722-1740.

570 Richard, B., Dupouey, J. L., Corcket, E., Alard, D., Archaux, F., Aubert, M., ... & Lenoir, J. (2021).  
571 The climatic debt is growing in the understorey of temperate forests: Stand characteristics matter.  
572 *Global Ecology and Biogeography*, 30(7), 1474-1487.

573 Ridding, L. E., Newton, A. C., Keith, S. A., Walls, R. M., Diaz, A., Pywell, R. F., & Bullock, J. M.  
574 (2021). Inconsistent detection of extinction debts using different methods. *Ecography*, 44(1), 33-43.

575 Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., ... & Marra,  
576 P. P. (2019). Decline of the North American avifauna. *Science*, 366(6461), 120-124.

577 Rumpf, S. B., Hülber, K., Wessely, J., Willner, W., Moser, D., Gattringer, A., ... & Dullinger, S.  
578 (2019). Extinction debts and colonization credits of non-forest plants in the European Alps. *Nature*  
579 *Communications*, 10(1), 4293.

580 Sabatini, F. M., Burrascano, S., Keeton, W. S., Levers, C., Lindner, M., Pötschner, F., ... &  
581 Kuemmerle, T. (2018). Where are Europe's last primary forests?. *Diversity and Distributions*, 24(10),  
582 1426-1439.

583 Sandel, B. (2015). Towards a taxonomy of spatial scale-dependence. *Ecography*, 38(4), 358-369.

584 Thompson, K. L., Chase, J. M., Remelgado, R., & Meyer, C. (2023). The interacting effects of climate  
585 and land-use/land-cover changes on ecological communities. *bioRxiv*, 2023-12.

586 Tehrani, N. A., Naimi, B., & Jaboyedoff, M. (2021). Modeling current and future species distribution  
587 of breeding birds as regional essential biodiversity variables (SD EBVs): A bird perspective in Swiss  
588 Alps. *Global Ecology and Conservation*, 27, e01596.

589 Tilman, D., May, R. M., Lehman, C. L., & Nowak, M. A. (1994). Habitat destruction and the


590 extinction debt. *Nature*, 371(6492), 65-66.

591 Tobias, J. A., Sheard, C., Pigot, A. L., Devenish, A. J., Yang, J., Sayol, F., ... & Schleuning, M. (2022).  
592 AVONET: morphological, ecological and geographical data for all birds. *Ecology Letters*, 25(3), 581-  
593 597.

594 Wenger, S. J., & Olden, J. D. (2012). Assessing transferability of ecological models: an  
595 underappreciated aspect of statistical validation. *Methods in Ecology and Evolution*, 3(2), 260-267.

596 Zurell, D., Franklin, J., König, C., Bouchet, P. J., Dormann, C. F., Elith, J., ... & Merow, C. (2020).  
597 A standard protocol for reporting species distribution models. *Ecography*, 43(9), 1261-1277.

598 Zurell, D., Schifferle, K., Herrando, S., Keller, V., Lehikoinen, A., Sattler, T., & Wiedenroth, L.  
599 (2024). Range and climate niche shifts in European and North American breeding birds.  
600 *Philosophical Transactions of the Royal Society B*, 379(1902), 20230013.



602

603


604

605

606

607


Figure 1: Example of approaches and limits to model current species distributions in the context of gradual environmental changes (B, C) and how to circumvent these limits by using weighting functions that describe how the effects of past environmental factors decrease as we go back far into the past (A).



608

609 Figure 2: Decrease, as we go back in time (from the onset of 2015 for which  $\Delta t = 0$ ), of the effects  
 610 of past land use and climate predictors on current bird species distribution in Europe. The grey curves  
 611 depict the half-Gaussian weighting function ( $W$ ) of the estimates for each species. The mean  
 612 weighting functions (solid curve) and the mean temporal extent of the time lags (vertical dashed line)  
 613 are depicted in green. The temporal extent of a time lag is the date at which we must go back into the  
 614 past to account for 95 % of the effect that a given time series of past environmental conditions has on  
 615 the current distribution of a species. For each predictor, the number of species considered varies  
 616 slightly (77 species, 69 species, 63 species and 60 species for primary forest, secondary forest,  
 617 precipitation and temperature, respectively) because we did not consider time lag parameters leading

618 to temporal extents that were larger than or equal to the maximum possible temporal extent available  
619 for the predictors (see Material and methods). The density plots represent the distribution of the  
620 temporal extents of the time lags.  
621



622

623 Figure 3: Stacked species distributions providing a richness map of current forest bird communities  
 624 predicted by non-equilibrium species distribution models (SDMs, A) and the associated extinction  
 625 debt (negative values, B) or immigration credit (positive values, B). In panel B, the colors depict the  
 626 difference between the species richness as predicted by stacking all species binary maps based on the  
 627 non-equilibrium SDM from that obtained by the equilibrium SDM. The extinction debt and the  
 628 immigration credit are expressed in percent of the local, within a grid cell, total number of species  
 629 predicted by equilibrium SDMs. The spatial resolution on the maps is 50 km.

630

631 **Supplementary Information**

632

633 Appendix S1: The bird species list.


634 Using the European Breeding Bird Atlas (EBBA2), we selected the 92 species of birds whose habitat  
 635 is classified as either forest or woodland according to Tobias et al. (2022). The use of a threshold of  
 636 a minimum number of presences was unnecessary because the proportion of occupied EBBA2's  
 637 meshes was sufficient for all the 92 selected species (minimum proportion > 0.10 that is at least 508  
 638 meshes out of 5079 for the whole study area).

639 Table S1: Species list.

640

<i>Acanthis flammea</i>	<i>Falco subbuteo</i>	<i>Phylloscopus bonelli</i>
<i>Accipiter gentilis</i>	<i>Ficedula albicollis</i>	<i>Phylloscopus collybita</i>
<i>Accipiter nisus</i>	<i>Ficedula hypoleuca</i>	<i>Phylloscopus sibilatrix</i>
<i>Aegithalos caudatus</i>	<i>Ficedula parva</i>	<i>Phylloscopus trochiloides</i>
<i>Aegolius funereus</i>	<i>Fringilla coelebs</i>	<i>Phylloscopus trochilus</i>
<i>Anthus trivialis</i>	<i>Fringilla montifringilla</i>	<i>Picoides tridactylus</i>
<i>Asio otus</i>	<i>Garrulus glandarius</i>	<i>Picus canus</i>
<i>Bombycilla garrulus</i>	<i>Glaucidium passerinum</i>	<i>Picus viridis</i>
<i>Bonasa bonasia</i>	<i>Hieraaetus pennatus</i>	<i>Poecile montanus</i>
<i>Carduelis carduelis</i>	<i>Hippolais icterina</i>	<i>Poecile palustris</i>
<i>Certhia brachydactyla</i>	<i>Hippolais polyglotta</i>	<i>Prunella modularis</i>
<i>Certhia familiaris</i>	<i>Jynx torquilla</i>	<i>Pyrrhula pyrrhula</i>
<i>Chloris chloris</i>	<i>Lanius collurio</i>	<i>Regulus ignicapilla</i>
<i>Ciconia nigra</i>	<i>Leiopicus medius</i>	<i>Regulus regulus</i>
<i>Circaetus gallicus</i>	<i>Lophophanes cristatus</i>	<i>Scolopax rusticola</i>
<i>Clanga pomarina</i>	<i>Loxia curvirostra</i>	<i>Serinus serinus</i>
<i>Coccothraustes coccothraustes</i>	<i>Loxia pytyopsittacus</i>	<i>Sitta europaea</i>
<i>Columba oenas</i>	<i>Lullula arborea</i>	<i>Spinus spinus</i>
<i>Columba palumbus</i>	<i>Luscinia megarhynchos</i>	<i>Streptopelia turtur</i>
<i>Coracias garrulus</i>	<i>Lyrurus tetrix</i>	<i>Strix aluco</i>
<i>Corvus corax</i>	<i>Merops apiaster</i>	<i>Strix uralensis</i>
<i>Cuculus canorus</i>	<i>Milvus migrans</i>	<i>Sylvia atricapilla</i>
<i>Cyanistes caeruleus</i>	<i>Muscicapa striata</i>	<i>Sylvia borin</i>
<i>Dendrocopos leucotos</i>	<i>Nucifraga caryocatactes</i>	<i>Tetrao urogallus</i>
<i>Dendrocopos major</i>	<i>Oriolus oriolus</i>	<i>Troglodytes troglodytes</i>
<i>Dendrocopos syriacus</i>	<i>Otus scops</i>	<i>Turdus iliacus</i>
<i>Dryobates minor</i>	<i>Parus major</i>	<i>Turdus merula</i>
<i>Dryocopus martius</i>	<i>Passer montanus</i>	<i>Turdus philomelos</i>
<i>Emberiza rustica</i>	<i>Periparus ater</i>	<i>Turdus pilaris</i>
<i>Erithacus rubecula</i>	<i>Pernis apivorus</i>	<i>Turdus viscivorus</i>
<i>Falco columbarius</i>	<i>Phoenicurus phoenicurus</i>	

641 Appendix S2: Spatial and temporal patterns of land uses and climate.  
642




643

644 Figure S1: Spatial patterns of land uses and climate in 2015.

645





646

647 Figure S2: Temporal patterns of land uses and climate conditions until 2015. Values are averaged  
 648 over the whole study area. Time series begin in 852 and in 1852 for land uses and climate conditions,  
 649 respectively. The historical data time series were pooled with the projections of a business-as-usual  
 650 scenario (SSP2, RCP 4.5) that begins after 2005, or after 2015, for the CHELSA and the LUH2 time  
 651 series, respectively (Karger et al., 2019; Hurtt et al., 2020).


652

653 Appendix S3: Comparison between simulated and inferred time lag parameters.

654

655 To illustrate the capability of our non-equilibrium species distribution modelling (SDM) approach to  
656 capture the slope estimates ( $\beta_j$ ) and the time lag ( $\gamma_j$ ) parameters of a lagging response to a given  
657 predictor variable  $j$ , we performed a simulation exercise by focusing on multiple reference species.  
658 First, the slope and the time-lag parameters (Eq. 1) we inferred from the bird species dataset (EBBA2),  
659 using our non-equilibrium SDM approach, were considered as the reference or the target (*i.e.*, the  
660 “true parameters”, Fig. S3) to rely on realistic parameter combinations. For each predictor, the  
661 number of species considered varies slightly (77 species, 69 species, 63 species and 60 species for  
662 primary forest, secondary forest, precipitation and temperature, respectively) because we did not  
663 consider time lag parameters leading to temporal extents that were larger than or equal to the  
664 maximum possible temporal extent available for the predictors (see Material and methods). Then, for  
665 the sake of comparison between our simulations and the empirical observations, we used the exact  
666 same set of environmental predictors (*i.e.*, long-term time series of changes in forest cover and climate  
667 conditions at a yearly resolution) and we focused on the exact same study area (*i.e.*, Europe) as for  
668 the bird data. For each combination of true parameters (*i.e.*, each bird reference species), we  
669 generated 30 replicates of 5079 predicted occurrences (across the entire study area), using the  
670 predicted probability of presence of each species as obtained from the non-equilibrium SDM fitted  
671 with the empirical data. From these simulated set of occurrences available for each of the reference  
672 species, we fitted 30 replicates of non-equilibrium SDMs from which we extracted the estimated  
673 slope and time-lag parameters (30 sets for each reference species). Finally, we used a linear regression  
674 to evaluate both the accuracy and the uncertainty of the estimated parameters using the median values  
675 of the 30 replicates for a given species. We found that the true slope parameters (those we used to  
676 perform the simulations), for the four predictor variables, were retrieved by our non-equilibrium  
677 inference procedure with a very large accuracy and a very small uncertainty ( $R^2 \geq 0.99$ , Fig. S4). This


678 was also true for the time-lag parameter associated with primary forests ( $R^2 = 0.97$ , Fig. S4), while  
 679 the accuracy and the uncertainty associated to the time-lag parameters of the other predictor variables  
 680 were smaller and larger, respectively ( $R^2 \geq 0.7$ , Fig. S4).



681

682 Figure S3: Summary of the procedure used to assess the capacity of the non-equilibrium modelling  
 683 approach to recover the original parameter values set in the model formula (Eq. 1). The procedure is  
 684 depicted for all reference species and one covariate  $j$ , for which we know the true values associated  
 685 with the slope ( $\beta_j$ ) and the time-lag ( $\gamma_j$ ) parameters. The accuracy and the uncertainty of the  
 686 parameters, inferred from the non-equilibrium approach, are assessed from a linear regression line (in  
 687 blue), the 95 % confidence intervals (in gray), the coefficient of determination ( $R^2$ ), and by comparing  
 688 the regression line with the 1:1 line (in red).

689




690

691 Figure S4: Relationship between the true and the inferred slope ( $\beta_j$ ) and time lag ( $\gamma_j$ ) parameters. The  
 692  $R^2$  value is the coefficient of determination, the blue line is a regression line with confidence intervals  
 693 in gray, and the red line is the 1:1 line.

694


696



697

698 Figure S5: Distribution of the estimates of the predictors for the equilibrium and the non-  
699 equilibrium models.


700



701

702 Figure S6: Comparison between the slope parameters of the predictors ( $\beta_j$ ) for the equilibrium and  
 703 the non-equilibrium models. The larger the time lags and the more the discrepancies between the  
 704 two models.

705



706

707 Figure S7: Validity of the non-equilibrium models according to the distribution (A) and the spatial  
 708 autocorrelation (B) of the residual errors. The residual errors (A) were calculated with the R *resid*  
 709 function and pooled here among all the non-equilibrium species distribution models (SDMs), for  
 710 clarity purpose. The Moran's *I* indices were calculated by distance classes for each species and then  
 711 pooled (B). The pooled residual errors followed a normal distribution centered on zero (mean = 0.000,  
 712 SD = 0.365, Student's *t* = 0.000, *p*-value > 0.05) such as for the pooled distribution of the Moran's *I*  
 713 index (mean = 0.003, SD = 0.14; Student's *t* = 0.893, *p*-value > 0.05).