



Rapid correction of chronic hyperglycemia and bone remodeling, warning against overdoing

Dured Dardari, Beatrice Segurens

► To cite this version:

Dured Dardari, Beatrice Segurens. Rapid correction of chronic hyperglycemia and bone remodeling, warning against overdoing. World Journal of Diabetes, 2024, 15 (9), pp.1858-1861. <10.4239/wjd.v15.i9.1858>. <hal-04909896>

HAL Id: hal-04909896

<https://hal.science/hal-04909896v1>

Submitted on 24 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-ND 4.0 - Attribution - No Derivative Works - International License



Rapid correction of chronic hyperglycemia and bone remodeling, warning against overdoing

Dured Dardari, Beatrice Segurens

Specialty type: Endocrinology and metabolism

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade A

Novelty: Grade A

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Wu ZJ

Received: January 29, 2024

Revised: June 4, 2024

Accepted: July 2, 2024

Published online: September 15, 2024

Processing time: 210 Days and 23.1 Hours



Dured Dardari, Department of Diabetology, Centre Hospitalier Sud Francilien, Corbeil Essonne 91100, France

Beatrice Segurens, Université Paris-Saclay, CEA, Centre National de Recherche en Génomique Humaine (CNRGH), Evry Courcouronnes, France

Corresponding author: Dured Dardari, PhD, Academic Editor, Doctor, Department of Diabetology, Centre Hospitalier Sud Francilien, No. 40 Avenue Serge Dassault, Corbeil Essonne 91100, France. dured.dardari@chsf.fr

Abstract

It is widely recognized that chronic hyperglycemia decreases bone quality, although little is known about the impact of the rapid correction of chronic hyperglycemia on the quality of bone remodeling. This spotlight article explores this correlation by focusing on the stages of bone remodeling linked to glucose levels.

Key Words: Bone remodeling; Chronic hyperglycemia; Intensive hyperglycemia treatment; Diabetes complication; Rapid correction of hyperglycemia

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The protein osteocalcin involved in bone remodeling is affected by glucose metabolism and variation. Osteoprotegerin and receptor activator of NF- κ B ligand also involved in bone modeling are equally sensitive to glucose variation. Bone remodeling is impaired when glucose levels are reduced and only not only when the blood glucose threshold is exceeded.

Citation: Dardari D, Segurens B. Rapid correction of chronic hyperglycemia and bone remodeling, warning against overdoing. *World J Diabetes* 2024; 15(9): 1858-1861

URL: <https://www.wjgnet.com/1948-9358/full/v15/i9/1858.htm>

DOI: <https://dx.doi.org/10.4239/wjd.v15.i9.1858>

INTRODUCTION

The protein osteocalcin (OC) involved in bone remodeling is affected by glucose metabolism and variation.

Osteoprotegerin (OPG) and receptor activator of NF- κ B ligand also involved in bone modeling are equally sensitive to glucose variation.

Bone remodeling is impaired when glucose levels are reduced and only not only when the blood glucose threshold is exceeded.

HOW RAPID CORRECTION OF CHRONIC HYPERGLYCAEMIA AFFECTS BONE REMODELING

It has been suggested that bone contributes to the overall physiology of the body as an endocrine hormone[1]. To maintain its activity, bone is constantly remodeled by osteoblasts (OB) and osteoclasts[2]. During the various stages of OB differentiation, different biomarkers such as the OC are involved. Secreted by OB, OC is a bone component and biomarker of bone mineralization. Nevertheless, there are several shortcomings regarding the idea of OC as an endocrine hormone[3]. OC increases the insulin sensitivity of the target organs by stimulating glucose uptake by the muscles, increasing adiponectin production by adipose tissue, reducing lipid accumulation and inflammation in the liver, and promoting insulin secretion by the pancreas. The latter action is directly related to glucose metabolism[3]. This demonstrates the link between bone remodeling and glucose metabolism *via* the impact of glucose metabolism on OC. Patients with hyperglycemia have low levels of OC[4,5]. If hyperglycemia is corrected rapidly, the level of OC will also rise, thus accelerating the bone remodeling process.

Furthermore OPG is a soluble glycoprotein produced mainly by OB, which inhibits osteoclast genesis by preventing the binding of the receptor activator of nuclear factor- κ B ligand (RANKL) to its receptor RANK[6]. However, RANKL is an osteoclast differentiation factor produced by OB, which triggers osteoclast genesis by binding to RANK, a membrane receptor expressed by osteoclast precursors[7]. In a study[8], OPG was inhibited by the correction of hyperglycemia, twenty-two patients with newly diagnosed type 1 diabetes were treated with insulin, with a drop in HbA1c from 11.1% (98 mmol/mol) at diagnosis to 6.2% (44 mmol/mol) after 6 months. The plasma OPG level in the patients before treatment was 3.1 ng/L, and after 6 months treatment it was decreased to 2.6 ng/L ($P < 0.001$) but was still higher than in 28 healthy controls (2.1 ng/L). Elevated levels of OPG have previously been shown in patients with both type 1 and type 2 diabetes and have been associated to hyperglycemia[9,10]. We would therefore conclude that rapidly reducing hyperglycaemia inhibits OPG and increases RANKL.

In a different vein, OB use glucose during differentiation *via* both oxidative phosphorylation and aerobic glycolysis [11]. However, it is also known that hyperglycemia plays a role in OB differentiation, whereas OB have insulin receptors. The binding of insulin to its OB receptor induces bone formation and the production of OC as well as a decrease in the expression of OPG[11]. It can therefore be concluded that the lack of insulin induced by hyperglycemia reduces the OB differentiation. However, insulin can stimulate the secretion of OC. OB that produce proteins also express an insulin receptor. Following its binding to its receptor, insulin not only induces bone formation but also decreases the expression of OPG, which inhibits the differentiation and activation of osteoclasts, the cells responsible for the resorption of the bone matrix.

Glucose levels, glucose metabolism, and insulin seem play a role in every stage of bone remodeling: They activate OC, reduce OPG, and differentiate OB. While previous studies have focused on the role played by hyperglycemia in the disruption and imbalance of bone remodeling, few have focused on the possible impact of hyperglycemia on bone remodeling, as presented in Figure 1, which illustrates how the correction of hyperglycemia impacts bone remodeling. However, in our focus we have not put the spotlight on the role of RANK due to the lack of studies on the topic.




Figure 1 How the correction of hyperglycemia impacts bone remodeling. OC: Osteocalcin; OPG: Osteoprotegerin; OB: Osteoblasts; RANKL: Receptor

activator of nuclear factor- κ B ligand.

CONCLUSION

This spotlight highlights the impact of the rapid correction of hyperglycemia on bone remodeling. In the literature, this impact has rarely been described. It has been suggested that certain profiles of patients living with chronic hyperglycemia should undergo a bone health assessment prior to the correction of chronic hyperglycemia. For example, simple bone densitometry can guide physicians or give clues about the state of the bone structure. This type of low-cost exploration can reduce the risk of bone diseases whose mechanism is linked to the alteration of bone modeling such as osteoporosis or Charcot's neuroarthropathy[12]. Finally, it is possible that the anti-RANKL therapies used in certain diseases such as osteoporosis could be a means of preventing the perturbation of bone modelling for people living with chronic hyperglycaemia and for whom a rapid reduction in this hyperglycaemia is planned in a short-term perspective.

ACKNOWLEDGEMENTS

The authors acknowledge Natacha Vitrat Gip Genopole Genopole Campus 1 20, Rue Henri Desbruères F-91030 Evry, FRANCE.

FOOTNOTES

Author contributions: Dardari D and Segurens B wrote the protocol, produced the statistics, collected the data, wrote and reviewed the manuscript; Dardari D and Segurens B are the guarantors of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis; All authors approved the final version of the manuscript.

Conflict-of-interest statement: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: France

ORCID number: Dured Dardari 0000-0002-7172-4300.

S-Editor: Li L

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 Veldhuis-Vlug AG, Fliers E, Bisschop PH. Bone as a regulator of glucose metabolism. *Neth J Med* 2013; **71**: 396-400 [PMID: 24127499]
- 2 Weivoda MM, Bradley EW. Macrophages and Bone Remodeling. *J Bone Miner Res* 2023; **38**: 359-369 [PMID: 36651575 DOI: 10.1002/jbmr.4773]
- 3 Lee NK, Sowa H, Hinoi E, Ferron M, Ahn JD, Confavreux C, Dacquin R, Mee PJ, McKee MD, Jung DY, Zhang Z, Kim JK, Mauvais-Jarvis F, Ducy P, Karsenty G. Endocrine regulation of energy metabolism by the skeleton. *Cell* 2007; **130**: 456-469 [PMID: 17693256 DOI: 10.1016/j.cell.2007.05.047]
- 4 Kanazawa I, Yamaguchi T, Yamauchi M, Yamamoto M, Kurioka S, Yano S, Sugimoto T. Serum undercarboxylated osteocalcin was inversely associated with plasma glucose level and fat mass in type 2 diabetes mellitus. *Osteoporos Int* 2011; **22**: 187-194 [PMID: 20165834 DOI: 10.1007/s00198-010-1184-7]
- 5 Im JA, Yu BP, Jeon JY, Kim SH. Relationship between osteocalcin and glucose metabolism in postmenopausal women. *Clin Chim Acta* 2008; **396**: 66-69 [PMID: 18657532 DOI: 10.1016/j.cca.2008.07.001]
- 6 Schoppet M, Preissner KT, Hofbauer LC. RANK ligand and osteoprotegerin: paracrine regulators of bone metabolism and vascular function. *Arterioscler Thromb Vasc Biol* 2002; **22**: 549-553 [PMID: 11950689 DOI: 10.1161/01.atv.0000012303.37971.da]
- 7 Yasuda H. RANKL, a necessary chance for clinical application to osteoporosis and cancer-related bone diseases. *World J Orthop* 2013; **4**: 207-217 [PMID: 24147256 DOI: 10.5312/wjo.v4.i4.207]
- 8 Xiang GD, Sun HL, Zhao LS, Hou J, Yue L, Xu L. [Changes of osteoprotegerin before and after insulin therapy in type 1 diabetic patients]. *Zhonghua Yi Xue Za Zhi* 2007; **87**: 1234-1237 [PMID: 17686254]
- 9 Rasmussen LM, Tarnow L, Hansen TK, Parving HH, Flyvbjerg A. Plasma osteoprotegerin levels are associated with glycaemic status, systolic

- blood pressure, kidney function and cardiovascular morbidity in type 1 diabetic patients. *Eur J Endocrinol* 2006; **154**: 75-81 [PMID: 16381994 DOI: 10.1530/eje.1.02049]
- 10 **Browner WS**, Lui LY, Cummings SR. Associations of serum osteoprotegerin levels with diabetes, stroke, bone density, fractures, and mortality in elderly women. *J Clin Endocrinol Metab* 2001; **86**: 631-637 [PMID: 11158021 DOI: 10.1210/jcem.86.2.7192]
- 11 **Medeiros C**, Wallace JM. High glucose-induced inhibition of osteoblast like MC3T3-E1 differentiation promotes mitochondrial perturbations. *PLoS One* 2022; **17**: e0270001 [PMID: 35714142 DOI: 10.1371/journal.pone.0270001]
- 12 **Dardari D**. Trends in the pathophysiology of Charcot neuroarthropathy. *Trends Endocrinol Metab* 2023; **34**: 61-62 [PMID: 36528439 DOI: 10.1016/j.tem.2022.12.003]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

