

Arbovirus Epidemics as Global Health Imperative, Africa, 2023

Salifou Bangoura, Alpha-Kabinet Keita, Maladho Diaby, Sidikiba Sidibé, Frederic Le Marcis, Saidouba Camara, Stéphanie Maltais, Kadio J.J.O. Kadio, Eric D'ortenzio, Alioune Camara, et al.

▶ To cite this version:

Salifou Bangoura, Alpha-Kabinet Keita, Maladho Diaby, Sidikiba Sidibé, Frederic Le Marcis, et al.. Arbovirus Epidemics as Global Health Imperative, Africa, 2023. Emerging Infectious Diseases, 2025, 31 (2), 10.3201/eid3102.240754. hal-04915856

HAL Id: hal-04915856 https://hal.science/hal-04915856v1

Submitted on 28 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EMERGING INFECTIOUS DISEASES[®]

Disclaimer: Early release articles are not considered as final versions. Any changes will be reflected in the online version in the month the article is officially released.

Volume 31, Number 2–February 2025

Online Report

Arbovirus Epidemics as Global Health Imperative, Africa, 2023

Salifou T. Bangouraz, Alpha-Kabinet Keita, Maladho Diaby, Sidikiba Sidibé, Frederic Le-Marcis, Saidouba C. Camara, Stéphanie Maltais, Kadio J.J.O. Kadio, Eric D'Ortenzio, Alioune Camara, Eric Delaporte, Alexandre Delamou, Philippe Vanhems, Michèle Ottmann, Nagham Khanafer, and Abdoulaye Touré

Author affiliation: Gamal Abdel Nasser University, Conakry, Guinea (S.T. Bangoura, A.-K. Keita, M. Diaby, S. Sidibé, K.J.J.O. Kadio, A. Camara, A. Delamou, A. Touré); Centre de Recherche et de Formation en Infectiologie de Guinée, Conakry (S.T. Bangoura, A.-K. Keita, M. Diaby, F. Le-Marcis, S.C. Camara, K.J.J.O. Kadio, A. Camara, A. Touré); TransVIHMI, Université de Montpellier-INSERM-IRD, Montpellier, France (A.-K. Keita, F. Le-Marcis, E. Delaporte); Ecole Normale Supérieure de Lyon, Lyon, France (F. Le-Marcis); University of Ottawa School of International Development and Global Studies, Ottawa, Ontario, Canada (S. Maltais); ANRS Maladies infectieuses émergentes, Inserm, Paris, France (E. D'Ortenzio); AP-HP, Hôpital Bichat, Service de maladies infectieuses et tropicales, Paris (E. D'Ortenzio); African Centre of Excellence in the Prevention and Control of Communicable Diseases, Gamal Abdel Nasser University, Conakry (A. Delamou); National Center for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea (A. Delamou); Centre International de Recherche en Infectiologie, Inserm, Lyon (P. Vanhems, M. Ottmann, N. Khanafer); Infection Control Unit, Hôpital Edouard Herriot, Hospices Civils de Lyon, Lyon (P. Vanhems, N. Khanafer) Suggested citation for this article

Abstract

Arboviruses represent a major cause of illness in Africa and have the potential to trigger widespread epidemics. We present data on arbovirus epidemics in Africa in 2023 and demonstrate the need for global public health authorities to intensify efforts in the surveillance and control of arbovirus diseases. Data were collected from the World Health Organization Weekly Bulletin on Outbreaks and Other Emergencies, Africa Centers for Disease Control and Prevention Weekly Event Based Surveillance Report, and other online sources. In 2023, a total of 7 arboviruses were responsible for 29 outbreaks across 25 countries in Africa, 22 of which occurred in West Africa; the outbreaks resulted in 19,569 confirmed cases and 820 deaths. Arbovirus epidemics in Africa pose a threat not only to public health within the continent but also globally, underscoring the urgent need for substantial investment in arbovirus surveillance, research, and preparedness capacities in Africa to prevent and respond to health crises effectively.

Arthropodborne viral diseases represent a major global health challenge because of their capacity to cause explosive outbreaks and induce severe, potentially lifethreatening clinical conditions, including encephalopathy, meningoencephalitis, myelitis, and symptoms of Guillain-Barré syndrome (<u>1,2</u>). Dengue virus (DENV) is the most prevalent arbovirus; >7.6 million cases had been reported to the World Health Organization (WHO) as of April 30, 2024, including 3.4 million confirmed cases, >16,000 severe cases, and >3,000 deaths ($\underline{3}$), imposing substantial economic burdens on many tropical and subtropical countries ($\underline{4}, \underline{5}$).

The frequency and scale of outbreaks caused by these arboviruses, particularly those transmitted by Aedes mosquitoes, are rising globally, driven by the intersection of ecologic, economic, and social factors (<u>6,7</u>). In response, the WHO launched the Global Arbovirus Initiative to "raise the global alarm on the risk epidemics of arboviruses and the potential risk of pandemics" (<u>6,8</u>). This initiative focuses on risk monitoring; pandemic prevention, preparedness, detection, and response; and the development of a coalition of partners (6). However, the capacity of countries in Africa to respond to arboviral threats is a concern, because most lack the necessary infrastructure and resources (laboratory equipment, trained personnel, and funding) to conduct adequate surveillance of arthropods and the viruses they transmit, whereas performing diagnostic tests to differentiate among viruses presents substantial challenges (<u>9–12</u>). As a result, most arbovirus infections go undiagnosed until epidemics emerge, causing severe health, social, and economic consequences (<u>13–15</u>). Furthermore, seroprevalence studies indicate that both endemic and epidemic transmission of arboviruses occurs regularly across Africa (2,16-18). In this context, where arbovirus infections circulate frequently in low- and middle-income countries global demographic, societal, and environmental trends of the 21st Century (19,20). A recent example is Zika virus (ZIKV), which swiftly transitioned from obscurity to a WHO Public Health Emergency of International Concern (21,22). That potential underscores the necessity for substantial investment in the arbovirus surveillance, research, and preparedness capacities of countries in Africa to effectively prevent and respond to future public health threats. In 2023, multiple arbovirus epidemics occurred in Africa, frequently occurring simultaneously and transcending borders (7,23,24). In this review, we present data on arbovirus epidemics reported in Africa in 2023, emphasizing the need for global public health authorities to take steps toward an equitable distribution of health efforts and resources that could enhance both local and global health security.

Methods

We compiled a database of arbovirus-related human epidemics reported in Africa during 2023, sourcing information from the WHO Weekly Bulletin on Outbreaks and Other Emergencies and the Africa Centers for Disease Control and Prevention Weekly Event-Based Surveillance Report. In addition, we consulted various online resources related to infectious diseases, including ProMED-mail (<u>http://www.promedmail.org</u> 2) and the International Society for Infectious Diseases (http://www.isid.org 2). For cases reported on ProMED-mail, we verified the source of information, specifically checking whether information was in the form of an official statement from public health authorities. We used a standardized form to extract and enter the data into an Excel spreadsheet (Microsoft, https://www.microsoft.com]). The dataset encompassed the country where the epidemic occurred, the type of arbovirus, the reporting date of the first cases, the number of suspected cases, the number of confirmed cases, the number of deaths, and the data sources. The country's notification date was used to remove duplicates. We conducted a descriptive analysis of the data to summarize the findings. We categorized arbovirus infection cases into 2 groups: suspected cases, which included all reports lacking laboratory confirmation, and confirmed cases, which consisted of all reports with biological verification through PCR or IgM ELISA.

<u>Top</u>

Figure. Geographic distribution of confirmed arbovirus cases reported

across Africa, 2023.


Results

In total, 29 arboviral outbreaks were reported across 25 countries in Africa in 2023; of those, 22 occurred in West Africa (Figure). Seven distinct arboviruses were responsible for the outbreaks: DENV in 17 countries; yellow fever virus (YFV) in 9 countries; chikungunya virus (CHIKV) in 4 countries; Crimean-Congo hemorrhagic fever virus (CCHFV), Rift Valley fever virus (RVFV), and West Nile virus (WNV) each in 3 countries; and ZIKV in 2 countries. Senegal recorded the highest number of outbreaks linked to arboviruses (DENV, CHIKV, ZIKV, YFV, CCHFV, and WNV), followed by Namibia (CHIKV, CCHFV, and WNV) and Mali (DENV, CHIKV, and ZIKV) (Figure).

In 2023, a total of 19,569 confirmed cases and 820 deaths were reported. Most infections were attributed to DENV, which accounted for 18,697 confirmed cases and 771 deaths. Burkina Faso faced an unprecedented epidemic, reporting 17,125 confirmed DENV cases and 688 deaths. Mali followed with 808 confirmed cases and 34 deaths, then Ethiopia with 272 cases and 17 deaths and Senegal with 254 confirmed cases but no fatalities. Nine countries (Cameroon, Côte d'Ivoire, Central African Republic, Gabon, Guinea, Nigeria, Republic of Congo, Senegal, and Uganda) reported a total of 104 confirmed yellow fever cases, resulting in 39 deaths. The highest

number of cases was recorded in Cameroon (49 cases, 6 deaths), Nigeria (21 cases, 21 deaths), Congo (15 cases, 2 deaths), and the Central African Republic (13 cases, 6 deaths). CHIKV infections were documented in 4 countries, with most cases reported in Burkina Faso (351 confirmed cases, 1 death) and Senegal (338 confirmed cases, no deaths) (Table).

Confirmed cases of CCHFV infection were documented in Senegal (8 cases, 2 fatalities), Mauritania (3 cases, no fatalities), and Namibia (1 fatal case). RVFV infection also led to several confirmed cases and deaths, notably in Uganda (27 cases, 4 deaths), whereas isolated cases were reported in both Niger and Mauritania. Tunisia registered the highest number of WNV infections, with 10 confirmed cases. Mali and Senegal together reported 24 confirmed cases of ZIKV infection, with no associated fatalities (Table)



ISSN: 1080-6059

📕 An official website of the United States government Here's how you know 🗸

Search

The repeated and concurrent arbovirus epidemics across Africa raise concerns about the effectiveness and sustainability of current control and surveillance strategies in the region. This situation highlights the pressing need for comprehensive investments in national health systems to enhance clinical and biologic surveillance and to strengthen control and prevention efforts against these diseases.

West Africa has been disproportionately impacted by multiple, concurrent arbovirus outbreaks. This finding aligns with previous studies (10,25), raising concerns about the region's capacity to conduct effective surveillance of arboviruses and their vectors, and to detect, prevent, and respond to epidemics. The conclusions of the survey assessing the capacity of health systems to prevent, detect, and respond to epidemics of arbovirus diseases in the 47 countries of the WHO Regional Office for Africa revealed that the West Africa region had inadequate capacity for virologic surveillance and routine vector surveillance and control (12). Moreover, the region's diverse vegetation, favorable climate, and presence of both wild and domestic vertebrate hosts create an ideal environment for vectors (<u>1,10,26,27</u>). Furthermore, models predicting the global distribution of arbovirus diseases highlight West Africa as a high-risk area because of its heavy rainfall and optimal temperature conditions for both endemic and epidemic transmission (<u>4,28</u>). As a consequence, adopting innovative diagnostic and surveillance methods, such as genomic sequencing, will be critical to enhance early detection of both endemic and emerging arboviruses, assess viral strains' epidemic potential, evaluate intervention effectiveness, and map transmission networks across humans, animals, and vectors (20,29). For example, in Nigeria, metagenomic sequencing was used to identify an ongoing yellow fever outbreak and its etiology and inform real-time public health actions, resulting in accurate and timely disease management and control (<u>30</u>). The ability of genomic sequencing to rapidly identify and characterize reemerging viruses offers a valuable opportunity for enhancing public health surveillance. Furthermore, fostering crosscountry collaboration and networking is crucial for harmonizing laboratory diagnostics and strengthening epidemic preparedness. Without such coordinated efforts, isolated interventions might be inadequate or entirely ineffective (9,11).

DENV was the most frequently reported infection; major epidemics occurred in Burkina Faso, Mali, Sudan, and Senegal. DENV is the most prevalent arbovirus globally; Africa accounts for 16% of all infections, positioning it as the second most affected region after Asia (31,32). Since the 1960s, DENV has remained a leading cause of illness across all African subregions (32); outbreaks have been documented in numerous countries, such as Tanzania (33–35), Mozambique (36), Ghana (37,38), Sudan (13), Benin (39), Burkina Faso (40), and Côte d'Ivoire (41). DENV is not only of considerable medical importance but also imposes a substantial economic burden, estimated in billions of dollars annually (<u>42,43</u>).

YFV remains a public health challenge in Africa, despite the availability of a highly effective vaccine that induces strong and long-lasting immunity (<u>44,45</u>). Numerous cases, some fatal, have been documented across the continent. The persistence of yellow fever in Africa is largely attributable to insufficient vaccination coverage; the disease is most prevalent in tropical sub-Saharan Africa, where it is preventable primarily through vaccination (46). According to WHO/United Nations Children's Fund estimates on national immunization coverage, regional coverage stood at only 48% in 2021, far below the 80% threshold needed to achieve herd immunity against yellow fever (<u>47,48</u>). This shortfall was likely caused by the concurrent outbreaks of COVID-19, Ebola virus disease, measles, and cholera, which disrupted routine vaccination services, restricted movement, and reduced healthcare access. Therefore, countries must take steps to strengthen national immunization programs, alongside implementing supplementary vaccination campaigns in high-risk areas. Those campaigns should include catch-up campaigns targeting vulnerable populations to sustain high vaccination coverage. Such efforts must be coupled with implementation research to ensure equitable vaccine access. Given that implementation research is often context-specific, bolstering local capacity to identify challenges and barriers to adequate coverage would benefit all nations (49).

The findings of this study contribute valuable insights into the ongoing circulation of arboviruses in Africa and underscore the critical need to enhance surveillance efforts for these viruses in both vectors and human populations. They also reinforce the importance of integrated strategies for epidemic prevention and response, including vector control, expanding vaccination coverage, promoting the use of mosquito nets, and implementing education and communication initiatives aimed at fostering sustainable behavioral changes.

The first limitation of this study is that data came mainly from official public health sources, but numbers could be underestimated because of the weakness of healthcare systems in surveillance and the clinical overlap between arbovirus infections and endemic diseases such as malaria, which limits access to diagnostic testing. In addition, healthcare workers' limited knowledge about arboviruses could lead to misdiagnosis and affect the data, as demonstrated by a recent study in public health facilities in Conakry, Guinea, where only 1% of healthcare workers had adequate knowledge of arboviruses (50). Moreover, this study examined data collected over a year, but it did not analyze long-term trends in arbovirus epidemics. Furthermore, the data used in this study were sourced from open-access platforms. As a result, their completeness and accuracy depend on the availability of reports from the surveillance systems of each country. Therefore, caution must be exercised in interpreting the findings, and comparisons between countries should be avoided. Repeated serologic surveys could serve as a valuable population-based surveillance tool to track trends in arbovirus infections, offering vital information for shaping public health policies, guiding vaccine deployment strategies, and identifying regions at elevated risk for future outbreaks (<u>51</u>).

The primary goal of global health, as promoted by international public health organizations, is to ensure equitable distribution of health resources and knowledge across all regions, including Africa (52). However, funding is often tied to specific priorities and agendas, making research programs in Africa vulnerable to fluctuations in financial support and the shifting interests of northern partners, leaving critical areas of research underfunded (53).

Today, the threat posed by arboviruses to global health and economic security is widely acknowledged (54). Not only are arbovirus outbreaks increasing, but the risk for a pandemic looms (<u>54</u>). This concern was emphasized by the WHO's Director for Pandemic and Epidemic Diseases during the launch of the Global Arbovirus Initiative, who noted that "the next pandemic could very likely be caused by a new arbovirus, and we already have some signals that the risk is increasing" (<u>6,8</u>).

Despite substantial global financial investments in vaccines, access and usage remain major challenges in Africa. According to the list of vaccines authorized by the US Food and Drug Administration and the WHO, vaccines exist for DENV, YFV, tick-borne encephalitis virus, Japanese encephalitis virus, and CHIKV (55,56). Yet, the YFV vaccine is the only one currently available in Africa. Africa is one of the regions most affected by these arboviruses and has limited resources for their prevention, monitoring, and control, but those diseases continue to be neglected on the continent, despite receiving substantial attention and resources in other regions (57–60).

Making equitable and sustainable investments in health and economic systems and infrastructure is imperative to enhance the quality and efficiency of health services in Africa, effectively preventing future global health emergencies. This effort involves strengthening laboratory and medical staff capacities by providing access to innovative and rapid diagnostic techniques (such as serologic and molecular multiplex tests, commercial kits for the rapid detection of certain arboviruses), particularly in areas endemic for arboviruses representing major health risks. Increasing access to testing could considerably improve the early diagnosis of cases, the speed of epidemic management, and consequently, the treatment of patients (<u>61</u>). Continued support from countries in Africa is needed to develop and implement surveillance and control programs for arbovirus vectors adapted to local contexts; to better understand the genetic diversity of species, their distribution, and their abundance; and to develop more effective and sustainable tools and strategies for the control of arboviruses and vectors. Furthermore, funding research in Africa on known and new arboviruses is essential for understanding their ecologies and the interactions between the components of vector systems (vectors, hosts, pathogens, and environment) and the evolution of these interactions, including the mechanisms of adaptation to new environments. Such investments represent a long-term commitment to building public health capacity and global health security (<u>62</u>), and the development of research agendas for global health initiatives should take into account the national needs and priorities of countries, in consultation with local stakeholders in the intervention contexts (52). Adopting a holistic, multi-sectoral, and collaborative approach that addresses the unique challenges of combating arboviruses in Africa is also essential. This collaborative effort will not only mitigate immediate epidemics but will also establish a resilient and adaptive framework for addressing future outbreaks (<u>63</u>).

<u>Top</u>

Conclusions

This study demonstrates that arboviruses are actively circulating in Africa, resulting in severe epidemics and fatalities across various regions. Currently, no specific treatments are available, and access to effective vaccines remains limited. The primary strategies for effectively combating arboviruses include robust surveillance and control of vector populations, alongside research initiatives aimed at enhancing vaccine coverage and accessibility. In addition, strengthening personal and community preventive measures such as the use of mosquito nets and the management of water-containing containers is essential. Those efforts are anticipated to yield substantial benefits within the framework of global health.

Sustained support for arbovirus surveillance and research activities in Africa is critical not only for pandemic preparedness but also for enhancing overall health resilience. Furthermore, regional and cross-border collaboration should be established in alignment with international health regulations to develop adequate capacities for preventing arbovirus diseases. Forming global alliances is essential to consolidate resources and strengthen capabilities related to arbovirus surveillance and response.

Dr. Bangoura is a pharmacist epidemiologist, PhD student, and researcher at the Centre de Recherche et de Formation en Infectiologie de Guinée. Her primary research interest is reemerging infectious diseases, specifically arboviruses.

<u>Top</u>

Acknowledgment

We thank the WHO Regional Office for Africa and Africa Centers for Disease Control and Prevention for the availability of the data. We also thank Foromo Timothée Béavogui for editing the manuscript.

- 1. Braack L, Gouveia de Almeida AP, Cornel AJ, Swanepoel R, de Jager C. Mosquito-borne arboviruses of African origin: review of key viruses and vectors. Parasit Vectors. 2018;11:29. DOI 🛛 PubMed 🗗 Google Scholar
- 2. Power GM, Vaughan AM, Qiao L, Sanchez Clemente N, Pescarini JM, Paixão ES, et al. Socioeconomic risk markers of arthropod-borne virus (arbovirus) infections: a systematic literature review and meta-analysis. BMJ Glob Health. 2022;7:e007735. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 3. World Health Organization. Dengue—global situation [cited 2024 Sep 9]. https://www.who.int/emergencies/disease-outbreak-news/item/2024-DON518
- 4. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. Nature. 2013;496:504–7. DOI PubMed I Google Scholar
- 5. Seyler T, Hutin Y, Ramanchandran V, Ramakrishnan R, Manickam P, Murhekar M. Estimating the burden of disease and the economic cost attributable to chikungunya, Andhra Pradesh, India, 2005-2006. Trans R Soc Trop Med Hyg. 2010;104:133–8. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 6. World Health Organization. Launch of the Global Arbovirus Initiative [cited 2024 Jan 13].
- https://www.who.int/news-room/events/detail/2022/03/31/default-calendar/global-arbovirus-initiative
- 7. World Health Organization. Disease outbreak news: geographical expansion of cases of dengue and chikungunya beyond the historical areas of transmission in the Region of the Americas [cited 2024 Feb 19]. https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON448
- 8. Balakrishnan VS. WHO launches global initiative for arboviral diseases. Lancet Microbe. 2022;3:e407. DOI 🖄 PubMed 🖄 Google Scholar 🖄
- 9. Braack L, Wulandhari SA, Chanda E, Fouque F, Merle CS, Nwangwu U, et al. Developing African arbovirus networks and capacity strengthening in arbovirus surveillance and response: findings from a virtual workshop. Parasit Vectors. 2023;16:129. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 10. Buchwald AG, Hayden MH, Dadzie SK, Paull SH, Carlton EJ. Aedes-borne disease outbreaks in West Africa: A call for enhanced surveillance. Acta Trop. 2020;209:105468. DOI 🛛 PubMed 🗗 Google Scholar
- 11. Dadzie SK, Akorli J, Coulibaly MB, Ahadji-Dabla KM, Baber I, Bobanga T, et al.; WAASuN contributors. Building the capacity of West African countries in Aedes surveillance: inaugural meeting of the West African Aedes Surveillance Network (WAASuN). Parasit Vectors. 2022;15:381. DOI 🖄 PubMed 🖄 Google Scholar 🖄
- 12. World Health Organiazation (WHO) Department of Control of Neglected Tropical Diseases, WHO Regional Office for Africa, TDR. Assessing African country capacities to prevent, detect and respond to arboviral disease outbreaks [cited 2023 Aug 10]. https://tdr.who.int/newsroom/news/item/11-12-2021-assessing-african-country-capacities-to-prevent-detect-and-respond-to-arboviral-disease-outbreaks
- 13. Elaagip A, Alsedig K, Altahir O, Ageep T, Ahmed A, Siam HA, et al. Seroprevalence and associated risk factors of Dengue fever in Kassala state, eastern Sudan. PLoS Negl Trop Dis. 2020;14:e0008918. DOI Z PubMed Z Google Scholar Z
- 14. Hill SC, Vasconcelos J, Neto Z, Jandondo D, Zé-Zé L, Aguiar RS, et al. Emergence of the Asian lineage of Zika virus in Angola: an outbreak investigation. Lancet Infect Dis. 2019;19:1138–47. DOI 🖄 PubMed 🖄 Google Scholar
- 15. Vairo F, Aimè Coussoud-Mavoungou MP, Ntoumi F, Castilletti C, Kitembo L, Haider N, et al.; On Behalf Of The Pandora-Id-Net Consortium Chikungunya Outbreak Group Taskforce. Chikungunya outbreak in the Republic of the Congo, 2019—epidemiological, virological and entomological findings of a south-north multidisciplinary taskforce investigation. Viruses. 2020;12:1020. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 16. Gebremariam TT, Schallig HDFH, Kurmane ZM, Danquah JB. Increasing prevalence of malaria and acute dengue virus coinfection in Africa: a meta-analysis and meta-regression of cross-sectional studies. Malar J. 2023;22:300. DOI 2 PubMed 2 Google Scholar 2
- 17. Kayange N, Hau DK, Pain K, Mshana SE, Peck R, Gehring S, et al. Seroprevalence of dengue and chikungunya virus infections in children living in sub-Saharan Africa: systematic review and meta-analysis. Children (Basel). 2023;10:1662. DOI 2 PubMed 2 Google Scholar 2
- 18. Mwanyika GO, Mboera LEG, Rugarabamu S, Ngingo B, Sindato C, Lutwama JJ, et al. Dengue virus infection and associated risk factors in Africa: a systematic review and meta-analysis. Viruses. 2021;13:536. DOI 2 PubMed 2 Google Scholar 2
- 19. Musso D, Rodriguez-Morales AJ, Levi JE, Cao-Lormeau VM, Gubler DJ. Unexpected outbreaks of arbovirus infections: lessons learned from the Pacific and tropical America. Lancet Infect Dis. 2018;18:e355–61. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 20. Wallau GL, Abanda NN, Abbud A, Abdella S, Abera A, Ahuka-Mundeke S, et al.; Global Arbovirus Researchers United. Arbovirus researchers unite: expanding genomic surveillance for an urgent global need. Lancet Glob Health. 2023;11:e1501–2. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 21. Şahiner F. [Global spread of Zika virus epidemic: current knowledges and uncertainties] [in Turkish]. Mikrobiyol Bul. 2016;50:333–51. DOI 🛽 PubMed 🗹 Google Scholar
- 22. Wilder-Smith A, Gubler DJ, Weaver SC, Monath TP, Heymann DL, Scott TW. Epidemic arboviral diseases: priorities for research and public health. Lancet Infect Dis. 2017;17:e101–6. DOI 🛛 PubMed 🗗 Google Scholar
- 23. Santé publique France. Chikungunya, dengue, and Zika—enhanced surveillance data in metropolitan France in 2023 [in French] [cited 2024 Feb 19]. https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-a-transmission-vectorielle/chikungunya/articles/donnees-en-francemetropolitaine/chikungunya-dengue-et-zika-donnees-de-la-surveillance-renforcee-en-france-metropolitaine-en-2023
- 24. World Health Organization. Dengue—global situation. 2023 Dec 21 [cited 2024 Jan 26].
- https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON498
- 25. Agboli E, Zahouli JBZ, Badolo A, Jöst H. Mosquito-associated viruses and their related mosquitoes in West Africa. Viruses. 2021;13:891. DOI 🛽 PubMed 🗹 Google Scholar
- 26. Weetman D, Kamgang B, Badolo A, Moyes CL, Shearer FM, Coulibaly M, et al. Aedes mosquitoes and Aedes-borne arboviruses in Africa: current and future threats. Int J Environ Res Public Health. 2018;15:220. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 27. Zahouli JBZ, Koudou BG, Müller P, Malone D, Tano Y, Utzinger J. Effect of land-use changes on the abundance, distribution, and host-seeking behavior of Aedes arbovirus vectors in oil palm-dominated landscapes, southeastern Côte d'Ivoire. PLoS One. 2017;12:e0189082. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 28. Bogoch II, Brady OJ, Kraemer MUG, German M, Creatore MI, Brent S, et al. Potential for Zika virus introduction and transmission in resource-limited countries in Africa and the Asia-Pacific region: a modelling study. Lancet Infect Dis. 2016;16:1237–45. DOI 🖄 PubMed 🖄 Google Scholar
- 29. Pronyk PM, de Alwis R, Rockett R, Basile K, Boucher YF, Pang V, et al. Advancing pathogen genomics in resource-limited settings. Cell Genomics. 2023;3:100443. DOI 🛛 PubMed 🖾 Google Scholar 🖾
- 30. Ajogbasile FV, Oguzie JU, Oluniyi PE, Eromon PE, Uwanibe JN, Mehta SB, et al. Real-time metagenomic analysis of undiagnosed fever cases unveils a yellow fever outbreak in Edo State, Nigeria. Sci Rep. 2020;10:3180. DOI 🛛 PubMed 🗹 Google Scholar 🖄
- 31. Centers for Disease Control and Prevention. Public health matters blog: unveiling the burden of dengue in Africa [cited 2024 Apr 18]. https://blogs.cdc.gov/publichealthmatters/2015/07/unveiling-the-burden-of-dengue-in-africa
- 32. Diallo D, Diouf B, Gaye A, NDiaye EH, Sene NM, Dia I, et al. Dengue vectors in Africa: A review. Heliyon. 2022;8:e09459. DOI 🖄 PubMed 🖄 Google Scholar 🖄
- 33. Chipwaza B, Sumaye RD, Weisser M, Gingo W, Yeo NKW, Amrun SN, et al. Occurrence of 4 dengue virus serotypes and chikungunya virus in Kilombero Valley, Tanzania, during the dengue outbreak in 2018. Open Forum Infect Dis. 2020;8:ofaa626. DOI 🖄 PubMed 🖄 Google Scholar 🖄
- 34. Vairo F, Mboera LEG, De Nardo P, Oriyo NM, Meschi S, Rumisha SF, et al. Clinical, virologic, and epidemiologic characteristics of dengue outbreak, Dar es Salaam, Tanzania, 2014. Emerg Infect Dis. 2016;22:895–9. DOI 🖄 PubMed 🖄 Google Scholar 🖄
- 35. Ward T, Samuel M, Maoz D, Runge-Ranzinger S, Boyce R, Toledo J, et al. Dengue data and surveillance in Tanzania: a systematic literature review. Trop Med Int Health. 2017;22:960–70. DOI 🛛 PubMed 🗹 Google Scholar 🗹
- 36. Mugabe VA, Ali S, Chelene I, Monteiro VO, Guiliche O, Muianga AF, et al. Evidence for chikungunya and dengue transmission in Quelimane, Mozambique: Results from an investigation of a potential outbreak of chikungunya virus. PLoS One. 2018;13:e0192110. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 37. Amoako N, Duodu S, Dennis FE, Bonney JHK, Asante KP, Ameh J, et al. Detection of dengue virus among children with suspected malaria, Accra, Ghana. Emerg Infect Dis. 2018;24:1544–7. DOI 🛛 PubMed 🗹 Google Scholar 🗠
- 38. Manu SK, Bonney JHK, Pratt D, Abdulai FN, Agbosu EE, Frimpong PO, et al. Arbovirus circulation among febrile patients at the greater Accra Regional Hospital, Ghana. BMC Res Notes. 2019;12:332. DOI D PubMed D Google Scholar
- 39. Allanonto V, Yanogo P, Sawadogo B, Akpo Y, Noudeke ND, Saka B, et al. Investigation of dengue cases in the departments of Atlantic, Littoral and Ouémé, Bénin, April–July 2019 [in French] [cited 2023 Jan 7]. https://www.afenet-journal.net/content/series/4/3/5/full HYPERLINK "https://doi.org/10.37432/jieph.supp.2021.4.3.03.5"
- 40. Ridde V, Agier I, Bonnet E, Carabali M, Dabiré KR, Fournet F, et al. Presence of three dengue serotypes in Ouagadougou (Burkina Faso): research and public health implications. Infect Dis Poverty. 2016;5:23. DOI 🛛 PubMed 🖄 Google Scholar
- 41. Ano AKMN, Coulibaly D, Bénie VJ, Akani BC, Douba A, Ahoussou EM, et al. Epidemiological situation of dengue fever in Côte d'Ivoire from 2017 to 2020 [in French]. Rev Epidemiol Sante Publique. 2022;70:S146. DOI Z Google Scholar Z
- 42. Castro MC, Wilson ME, Bloom DE. Disease and economic burdens of dengue. Lancet Infect Dis. 2017;17:e70–8. DOI 🛽 PubMed 🗗 Google Scholar 🗗
- 43. Shepard DS, Undurraga EA, Halasa YA, Stanaway JD. The global economic burden of dengue: a systematic analysis. Lancet Infect Dis. 2016;16:935–41. DOI 🗹 PubMed I Google Scholar
- 44. Beck AS, Barrett ADT. Current status and future prospects of yellow fever vaccines. Expert Rev Vaccines. 2015;14:1479–92. DOI 2 PubMed 2 Google Scholar
- 45. Schnyder JL, de Jong HK, Bache BE, Schaumburg F, Grobusch MP. Long-term immunity following yellow fever vaccination: a systematic review and metaanalysis. Lancet Glob Health. 2024;12:e445–56. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 46. Centers for Disease Control and Prevention. Areas with risk of yellow fever virus transmission in Africa [cited 2024 Jan 30]. https://www.cdc.gov/yellowfever/maps/africa.html
- 47. World Health Organization. Yellow fever—African region (AFRO) [cited 2024 Jan 30]. https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON431
- 48. World Health Organization. WHO/UNICEF estimates of national immunization coverage [cited 2024 Apr 22].
- https://www.who.int/teams/immunization-vaccines-and-biologicals/immunization-analysis-and-insights/global-monitoring/immunization-coverage/who-unicefestimates-of-national-immunization-coverage
- 49. Maher D, Aseffa A, Kay S, Tufet Bayona M. External funding to strengthen capacity for research in low-income and middle-income countries: exigence, excellence and equity. BMJ Glob Health. 2020;5:e002212. DOI 🛛 PubMed 🖄 Google Scholar
- 50. Bangoura ST, Hounmenou CG, Sidibé S, Camara SC, Mbaye A, Olive MM, et al. Exploratory analysis of the knowledge, attitudes and perceptions of healthcare workers about arboviruses in the context of surveillance in the Republic of Guinea. PLoS Negl Trop Dis. 2023;17:e0011814. DOI 🛛 PubMed 🖄 Google Scholar 🖄
- 51. Nwaiwu AU, Musekiwa A, Tamuzi JL, Sambala EZ, Nyasulu PS. The incidence and mortality of yellow fever in Africa: a systematic review and meta-analysis. BMC Infect Dis. 2021;21:1089. DOI 🛛 PubMed 🗹 Google Scholar
- 52. Bain LE, Adeagbo OA, Avoka CK, Amu H, Memiah P, Ebuenyi ID. Identifying the conundrums of "global health" in the Global North and Global South: a case for Sub-Saharan Africa. Front Public Health. 2024;12:1168505. DOI 2 PubMed 2 Google Scholar 2
- 53. Caelers D, Okoth D. Research funding in Africa: navigating sustainability and shifting perspectives [cited 2024 Feb 4]. https://www.nature.com/articles/d44148-023-00360-4
- 54. Girard M, Nelson CB, Picot V, Gubler DJ. Arboviruses: A global public health threat. Vaccine. 2020;38:3989–94. DOI 🛽 PubMed 🖄 Google Scholar 🖄
- 55. US Food and Drug Administration. Vaccines licensed for use in the United States [cited 2024 Feb 9].
- https://www.fda.gov/vaccines-blood-biologics/vaccines/vaccines-licensed-use-united-states
- 56. World Health Organization. Vaccine-preventable diseases [cited 2024 Feb 9]. https://www.who.int/teams/immunization-vaccines-and-biologicals/diseases 🖸
- 57. Gainor EM, Harris E, LaBeaud AD. Uncovering the burden of dengue in Africa: considerations on magnitude, misdiagnosis, and ancestry. Viruses. 2022;14:233. DOI ☑ PubMed ☑ Google Scholar ☑
- 58. Were F. The dengue situation in Africa. Paediatr Int Child Health. 2012;32(Suppl 1):18–21. DOI 🛽 PubMed 🖄 Google Scholar 🖄
- 59. Head MG, Brown RJ, Newell ML, Scott JAG, Batchelor J, Atun R. The allocation of USdollar;105 billion in global funding from G20 countries for infectious disease research between 2000 and 2017: a content analysis of investments. Lancet Glob Health. 2020;8:e1295–304. DOI 🛛 PubMed 🖄 Google Scholar
- 60. Kading RC, Cohnstaedt LW, Fall K, Hamer GL. Emergence of arboviruses in the United States: the boom and bust of funding, innovation, and capacity. Trop Med Infect Dis. 2020;5:96. DOI PubMed Google Scholar
- 61. Migné CV, Beck C, Gonzalez G, Lecollinet S, Moutailler S. [Which tools for monitoring emerging arboviruses within their mammalian hosts and arthropod vectors?] [in French]. Virologie. 2021;25:12–28. DOI 🛛 PubMed 🖄 Google Scholar
- 62. Moser F, Habtemariam MK, Forland F, Djoudalbaye B. A new precedent in international health cooperation from African public health leaders. Lancet Glob Health. 2023;11:e828–9. DOI 🖄 PubMed 🖄 Google Scholar
- 63. Côrtes N, Lira A, Prates-Syed W, Dinis Silva J, Vuitika L, Cabral-Miranda W, et al. Integrated control strategies for dengue, Zika, and Chikungunya virus infections. Front Immunol. 2023;14:1281667. DOI 🖄 PubMed 🖄 Google Scholar

Figure

Figure. Geographic distribution of confirmed arbovirus cases reported across Africa, 2023.

Table

Table. Distribution of arbovirus cases in humans, Africa, 2023

<u>Top</u>

<u>Top</u>

Suggested citation for this article: Bangoura ST, Keita A-K, Diaby M, Sidibé S, Le-Marcis F, Camara SC, et al. Arbovirus epidemics as global health imperative, Africa, 2023. Emerg Infect Dis. 2025 Feb [date cited]. https://doi.org/10.3201/eid3102.240754

DOI: 10.3201/eid3102.240754

Original Publication Date: January 21, 2025

Table of Contents – Volume 31, Number 2—February 2025

EID Search Options

Advanced Article Search – Search articles by author and/or keyword.

Articles by Country Search – Search articles by the topic country.

Q <u>Article Type Search</u> – Search articles by article type and issue.

				<u>Top</u>
Comments				
Please use the form below to submit correspondence to the authors or contact them at the following address:				
Salifou Talassone Bangoura, Centre de Recherche et de Formation en Infectiologie de Guinée (CERFIG), Gamal Abdel Nasser University, Conakry, 6629P, Guinea				
Return Address				
Send To				
• Authors O Editors				
Comments				
	li			
10000 character(s) remaining.				
Send				
				<u>Top</u>
Page created: January 06, 2025				
Page updated: January 21, 2025				
Page reviewed: January 21, 2025				
The conclusions, findings, and opinions expressed by authors contributing to this journal do not necessarily reflect the official position of the U.S. Department of Health and Human Services, the Pub	lic Health S€	ervice, the Cent	ers for Diseas	e Control and
Prevention, or the authors' affiliated institutions. Use of trade names is for identification only and does not imply endorsement by any of the groups named above.				
About CDC Contact Us 800-232-4636	in		P	

U.S. Department of Health & Human Services Accessibility External Links Privacy Policies No Fear Act FOIA Nondiscrimination OIG Vulnerability Disclosure Policy Public Health Publications USA.gov