



HAL
open science

Editorial: Broadening our view on nucleic acid sensing: novel sensors, signaling pathways, and involvement in non-infectious diseases

Lise Chauveau, Nadine Laguette, Natalia Sampaio

► To cite this version:

Lise Chauveau, Nadine Laguette, Natalia Sampaio. Editorial: Broadening our view on nucleic acid sensing: novel sensors, signaling pathways, and involvement in non-infectious diseases. *Frontiers in Immunology*, 2023, 14, pp.3. 10.3389/fimmu.2023.1266732 . hal-04918462

HAL Id: hal-04918462

<https://hal.science/hal-04918462v1>

Submitted on 29 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED AND REVIEWED BY
Francesca Granucci,
University of Milano-Bicocca, Italy

*CORRESPONDENCE

Lise Chauveau

✉ lise.chauveau@cnrs.fr

Nadine Laguette

✉ nadine.laguette@cnrs.fr

Natalia G. Sampaio

✉ natalia.sampaio@hudson.org.au

RECEIVED 25 July 2023

ACCEPTED 28 July 2023

PUBLISHED 11 August 2023

CITATION

Chauveau L, Laguette N and Sampaio NG (2023) Editorial: Broadening our view on nucleic acid sensing: novel sensors, signaling pathways, and involvement in non-infectious diseases. *Front. Immunol.* 14:1266732. doi: 10.3389/fimmu.2023.1266732

COPYRIGHT

© 2023 Chauveau, Laguette and Sampaio. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Editorial: Broadening our view on nucleic acid sensing: novel sensors, signaling pathways, and involvement in non-infectious diseases

Lise Chauveau^{1*}, Nadine Laguette^{2*} and Natalia G. Sampaio^{3,4*}

¹Institut de Recherche en Infectiologie de Montpellier (IRIM) - CNRS UMR 9004, Université de Montpellier, Montpellier, France, ²IGMM, Université de Montpellier, CNRS, Montpellier, France, ³Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research, Clayton, VIC, Australia, ⁴Department of Molecular and Translational Sciences, School of Clinical Sciences, Monash University, Clayton, VIC, Australia

KEYWORDS

nucleic acid sensing, pattern recognition receptor (PRR), PAMPs and DAMPs, self/nonself recognition, sterile inflammation, non-infectious disease

Editorial on the Research Topic

Broadening our view on nucleic acid sensing: novel sensors, signaling pathways, and involvement in non-infectious diseases

Nucleic acid sensing is a crucial part of the innate immune system, and nucleic acid sensors are part of a class of receptors broadly called Pattern Recognition Receptors (PRRs). PRRs were initially studied as part of the immune response to pathogens. The concept stated that hosts need receptors to broadly sense incoming pathogens in a non-specific manner and trigger the activation of cells required for mounting pathogen-specific adaptive immune responses. In line with this core concept, PRRs recognize Pathogen Associated Molecular Patterns (PAMPs), which consist of parts of the invading pathogens such as their nucleic acid genomes. The binding of a PRR to a PAMP induces a signaling cascade in the infected cell, leading to the production of cytokines, including interferons, that are secreted into the extracellular milieu. These cytokines have multiple effects such as promoting resistance to infection in the neighboring cells and recruiting immune cells crucial for the adaptive response. However, how PRRs distinguish host nucleic acids (self) from pathogen-derived nucleic acids (non-self) has long been under investigation. Moreover, PRRs can be activated in sterile conditions (i.e. in the absence of pathogens), owing to the existence of Danger-Associated Molecular Patterns (DAMPs) that arise during infectious or non-infectious pathologies, and can include self-nucleic acids. Identifying the nature of these PRR-activating self-nucleic acids is an area of ongoing research that could inform on the mechanisms of self/non-self recognition. New PRRs are also still being discovered, and roles of PRRs beyond the production of cytokines are being reported. The field of nucleic acid sensing is therefore expanding at multiple levels, and this Research Topic aims to broaden our view on this complex area of research.

Regulating unusual self-nucleic acid sensing in sterile inflammation

In sterile inflammation, all nucleic acids detected by PRRs are self-nucleic acids. This suggests that some self-nucleic acids are more prone to inducing an immune response than others and the mechanisms regulating this recognition are poorly understood. In this Research Topic, [Gutierrez-Chamorro et al.](#) studied how SAMHD1, a deoxynucleotide triphosphate hydrolase, regulates innate immune responses via the RNA sensors MDA5 and RIG-I in ovarian cancer. Their clinical data suggest that stratifying patients according to SAMHD1 expression in the tumor could inform on ovarian cancer clinical outcomes, thereby helping to orient patient care. Furthermore, [Straub and Sampaio](#) review how RNA sensors can detect unusual self-RNA in the absence of pathogens, and the identification of the endogenous ligands involved. They further discuss the implications for non-infectious diseases and how these mechanisms can be targeted therapeutically. Moreover, nucleic acid sensing is not restricted to animals and considerably less is known about DNA sensing in plants, where non-self DNA can originate from pathogens or other plants. [Vega-Munoz et al.](#) investigated the role of the ATM and ATR DNA damage response proteins in self vs non-self DNA sensing in plants in sterile conditions. They show that these proteins are required for part of the self/non-self response normally activated by pathogens or minor physical damage, but another mechanism is responsible for activation of the wound response to intensive tissue damage.

New pathways for propagation of nucleic acid signaling

Production of cytokines and interferons in response to nucleic acid sensing is thought to occur within the same cell that detected the PAMPs/DAMPs and, in this model, these molecules are responsible for propagating the signal. However, recent studies challenged this model in the cGAS-STING DNA sensing pathway. In this Research Topic, [Blest and Chauveau](#) review the different pathways used by cGAMP, the second messenger in the cGAS-STING pathway, to travel between cells. In this case, sensing of DNA occurs in one cell but production of cytokines and interferons can happen in neighboring or even distant cells, thereby expanding the range of the innate immune response. They detail how these mechanisms have been described in various pathologies, in the presence or absence of pathogens, and how these could be targeted for therapies.

New consequences of nucleic acid sensing

Beyond the production of cytokines and interferons, nucleic acid sensing can result in a range of effects that are currently being uncovered. In this Research Topic, [Aubé et al.](#) review how DNA sensors are involved in the formation of Neutrophil extracellular traps (NETs), a neutrophil-specific response that can trap

pathogens, thereby limiting infection, but can also be detrimentally induced during sterile inflammation. As NETs themselves contain DNA, the authors further describe the DNA sensors that are activated in this pathogenic inflammatory response. In addition, some of these new immune functions of nucleic acid sensing have been attributed to other known proteins. On this Research Topic, [Cornec and Poirier](#) review recent findings on the existence of RNA interference in mammalian cells, especially in stem cells. There, Dicer proteins sense self-RNAs, particularly from transposable elements, and degrade them to avoid unwanted effects of these RNAs on gene expression and genome stability.

Implications of nucleic acid sensing in non-infectious diseases

Due to the increasing evidence that nucleic acid sensing pathways can detect self-nucleic acids, there is a growing interest in identifying how these pathways play a role in non-infectious diseases. In this Research Topic, [He et al](#) review the implications of cytoplasmic DNA sensing in various lung pathogenesises, including inflammatory diseases and cancer. Knowledge of the different mechanisms of self-DNA release into the cytoplasm, the PRRs involved in DNA detection, and the downstream pathways activated, could inform innovative therapies. Moreover, [Fetter et al](#) review the importance of inflammasome activation in both inflammatory and auto-immune skin diseases.

General conclusion

Overall, these articles and reviews highlight novel findings of nucleic acid sensing at multiple levels: nucleic acid detection, propagation of the signal, molecular consequences and broad implications for non-infectious diseases. This Research Topic expands our view of this complex field, and we hope will spark further research and innovation in this topic.

Author contributions

LC: Writing – original draft, Writing – review & editing. NL: Writing – review & editing. NS: Writing – review & editing.

Funding

LC is funded by Agence Nationale de Recherche sur le SIDA et les Hépatites virales (ANRS: ECTZ134113 and ECTZ134139). Work in NL's laboratory is currently funded by LA LIGUE pour la recherche contre le cancer (ARNs therapeutics, AAPARN 2021.LCC/JuF), Agence Nationale de Recherche sur le SIDA et les Hépatites virales (ANRS: ECTZ117448), la fondation ARC, La Région Languedoc Roussillon, the CNRS and the Université de Montpellier (Plan d'investissements France 2030, LabUM EpiGenMed, I-SITE). NS is funded by an mRNA Victoria

Activation Program Grant from the Government of Victoria, Australia.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.