



HAL
open science

COARSE DISTANCE FROM DYNAMICALLY CONVEX TO CONVEX

Julien Dardennes, Jean Gutt, Vinicius Ramos, Jun Zhang

► **To cite this version:**

Julien Dardennes, Jean Gutt, Vinicius Ramos, Jun Zhang. COARSE DISTANCE FROM DYNAMICALLY CONVEX TO CONVEX. 2025. <hal-04920068>

HAL Id: hal-04920068

<https://hal.science/hal-04920068v1>

Preprint submitted on 29 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

COARSE DISTANCE FROM DYNAMICALLY CONVEX TO CONVEX

JULIEN DARDENNES, JEAN GUTT, VINICIUS G. B. RAMOS, AND JUN ZHANG

ABSTRACT. Chaidez and Edtmair have recently found the first examples of dynamically convex domains in \mathbb{R}^4 that are not symplectomorphic to convex domains (called symplectically convex domains), answering a long-standing open question. In this paper, we discover new examples of such domains without referring to Chaidez-Edtmair's criterion in [3]. We also show that these domains are arbitrarily far from the set of symplectically convex domains in \mathbb{R}^4 with respect to the coarse symplectic Banach-Mazur distance by using an explicit numerical criterion for symplectic non-convexity.

1. INTRODUCTION AND MAIN RESULTS

1.1. **Introduction.** Convex domains in \mathbb{R}^{2n} have very strong symplectic rigidity properties, including the existence of closed orbits on its boundary in many cases [18, 12], the celebrated strong Viterbo conjecture that claims the equality of all the normalized symplectic capacities, as well as its interesting consequences on the systolic inequality between symplectic capacity and volume [22] and its implication to Mahler conjecture in convex geometry [1]. However, geometric convexity is not preserved by symplectic transformations.

The main motivation for this paper is the desire to understand **the symplectic analogue of convexity**. Hofer, Wysocki and Zehnder observed that the characteristic flow on the boundary of a convex set satisfies a certain index condition [12], that they called *dynamical convexity*, which is preserved by symplectomorphisms. For a long time, it was speculated that if a domain is dynamically convex, then it is symplectomorphic to a convex domain. Chaidez and Edtmair have shown [3] that there exist dynamically convex domains that are not symplectomorphic to convex ones in dimension 4. A recent work in [8] discovers more such examples which have been then easily applied to Chaidez and Edtmair's generalization of their criterion in higher dimensions [2].

In this paper we answer the following quantitative question in dimension 4.

Question 1. *How far can dynamically convex domains be away from convex domains?*

Let \mathcal{C}_4 be the set of domains in \mathbb{R}^4 which are symplectomorphic to convex domains, called symplectically convex, and let \mathcal{D}_4 denote the set of dynamically convex domains as first introduced in [12]. It follows from the same paper that $\mathcal{C}_4 \subset \mathcal{D}_4$.

A non-linear symplectic analogue of the Banach-Mazur distance (8) was first suggested by Ostrover and Polterovich and further developed by [21, 20]. It is defined as follows. For $U, V \subset \mathbb{R}^4$, let

$$(1) \quad d_c(U, V) = \inf \left\{ \log \lambda \geq 0 \mid \frac{1}{\lambda}U \hookrightarrow V \hookrightarrow \lambda U \right\}$$

where \hookrightarrow represents a symplectic embedding via some symplectomorphism ϕ of \mathbb{R}^4 . According to [21, Definition 1.3] where d_c is named as the coarse symplectic Banach-Mazur distance, we briefly call d_c the *coarse distance* in this paper. Following what is usually done in any metric space, given \mathcal{A}, \mathcal{B} two collections of subsets of \mathbb{R}^4 , we define the distance from \mathcal{A} to \mathcal{B} as:

$$(2) \quad \sigma_{d_c}(\mathcal{A}, \mathcal{B}) = \sup_{U \in \mathcal{A}} \inf_{V \in \mathcal{B}} d_c(U, V).$$

Note that $\sigma_{d_c}(-, -)$ is not symmetric in general. For instance, if $\mathcal{A} \subset \mathcal{B}$, then $\sigma_{d_c}(\mathcal{A}, \mathcal{B}) = 0$ while $\sigma_{d_c}(\mathcal{B}, \mathcal{A})$ could be large. Moreover, $\sigma_{d_c}(-, -)$ satisfies a monotonicity property, namely, $\sigma_{d_c}(\mathcal{A}, \mathcal{B}') \leq \sigma_{d_c}(\mathcal{A}, \mathcal{B})$ if $\mathcal{B} \subset \mathcal{B}'$. In what follows, for simplicity, let us denote σ_{d_c} by σ .

The main result of this paper is to answer Question 1 under the distance d_c . It turns out that the d_c -distance from the set of dynamically convex domains to the set of symplectically convex domains is unbounded.

Theorem 1.1. $\sigma(\mathcal{D}_4, \mathcal{C}_4) = +\infty$.

Along with the proof of Theorem 1.1, we discover a family of dynamically convex domains X_{Ω_p} (see (3)), parametrized by $p \in (0, 1]$, that are not symplectomorphic to convex ones when p is sufficiently small.

Novelty: The novelty of the toric domain X_{Ω_p} in the proof of Theorem 1.1 consists of the following four points:

(i) It is the first family of dynamically convex domains whose symplectic non-convexity can be verified *without* referring to Chaidez-Edtmair’s criterion.

(ii) The verification is only based on the classical machinery - ECH capacities (see [13]), where a concrete estimation on how small p can be so that X_{Ω_p} is not symplectically convex is obtained (see Remark 1.3). Note that an estimation on p could also be obtained via Chaidez-Edtmair’s criterion (See Remark 1.1).

(iii) It shows a successful metric-geometrical application, based on a symplectic version of the classical John’s ellipsoid theorem (see Section 2) that quantitatively characterizes the property of being symplectomorphic to convex domains. In fact, our second main result Theorem 1.4 illustrates that this metric geometry approach is “independent” of Chaidez-Edtmair’s criterion.

(iv) It indicates a new direction going to infinity in the large-scale geometry of the pseudo-metric space that consists of all the star-shaped domains in \mathbb{R}^4 equipped with d_c (see Remark 1.3).

Remark 1.1 (About the point (ii) above). *Soon after the first version of this paper appears to the public, Oliver Edtmair informed us that, by chasing the arguments in [3], one was able to obtain an explicit estimate of C in the criterion (7), though not presented in [3]. As a consequence, based on a careful calculation of $c_{\text{CE}}(X_{\Omega_p})$, this upper bound also helps to estimate, in a relatively precisely manner, the parameter p in X_{Ω_p} for symplectic non-convexity.*

1.2. d_c -large toric domains. To state our first main result, towards the proof of Theorem 1.1, let us recall some notations from the symplectic toric geometry. Let $\mu : \mathbb{R}^{2n} \rightarrow [0, \infty)$ be the standard moment map, i.e.,

$$\mu(z_1, \dots, z_n) = (\pi|z_1|^2, \dots, \pi|z_n|^2).$$

A toric domain is a set of the form $X_\Omega = \mu^{-1}(\Omega)$, where Ω is the closure of a non-empty open set. We say that X_Ω is star-shaped if the origin is in the interior of X_Ω and $r \cdot X_\Omega \subset X_\Omega$ for $0 < r < 1$. Given a star-shaped toric domain X_Ω , let $\partial_+ \Omega$ denote the closure of $\partial\Omega \cap (0, \infty)^n$. In dimension 4, $\partial_+ \Omega$ is a curve that connects the x and the y axes. We say that a star-shaped toric domain is monotone if $\partial_+ \Omega$ is the graph of a decreasing function. Let \mathcal{T}_4 and \mathcal{M}_4 denote the sets consisting of the star-shaped toric domains and of the monotone toric domains, respectively. It was shown in [9, Theorem 1.7] that for every $U \in \mathcal{M}_4$, all the normalized symplectic capacities of U coincide. The set \mathcal{M}_4 is quite a large set including all

convex and concave toric domains, see [9, Definition 2.1, 2.2], for example. In [9, Proposition 1.8], it was shown that $\mathcal{M}_4 = \mathcal{T}_4 \cap \mathcal{D}_4$.

The following Figure 1 illustrates the schematic relations between \mathcal{T}_4 , \mathcal{C}_4 , \mathcal{M}_4 , and \mathcal{D}_4 , where \mathcal{M}_4 is the shaded region.




FIGURE 1. Relations between \mathcal{T}_4 , \mathcal{C}_4 , \mathcal{M}_4 , and \mathcal{D}_4 .

As mentioned above, the relation $\mathcal{C}_4 \subset \mathcal{D}_4$ is given by [12]. It is strict due to examples that were recently found by [3] and [8]. So for the proof of Theorem 1.1, it suffices to find a family of monotone toric domains that are arbitrary far from all elements of \mathcal{C}_4 .


Now consider the following toric domain denoted by $X_{\Omega_p} \in \mathcal{M}_4$, where Figure 2 shows the moment image Ω_p . Explicitly, for $p \in (0, 1]$, let

$$(3) \quad X_{\Omega_p} := \left\{ (z_1, z_2) \in \mathbb{C}^2 \mid (\pi|z_1|^2)^p + (\pi|z_2|^2)^p < 1 \right\}.$$

Define $d_c(X_{\Omega_p}, \mathcal{E}_4) = \inf_{E \in \mathcal{E}_4} d_c(X_{\Omega_p}, E)$, where \mathcal{E}_4 consists of symplectic ellipsoids in \mathbb{R}^4 defined by (11). Here is a main result.

Theorem 1.2. *For toric domain X_{Ω_p} defined as above, we have the following estimation for $p < \frac{1}{5}$,*

$$(4) \quad d_c(X_{\Omega_p}, \mathcal{E}_4) \geq \frac{1}{8} \log \left(\frac{g(p)}{1 + \log 4 + \log g(p)} \right)$$

FIGURE 2. X_{Ω_p} for $p \in (0, 1]$.

where $g(p) = 2^{\frac{2}{p}-2} \text{Vol}_{\mathbb{R}^4}(X_{\Omega_p})$. In particular, when p satisfies the condition that

$$(5) \quad \frac{g(p)}{1 + \log 4 + \log g(p)} \geq 256,$$

then X_{Ω_p} is dynamically convex but not symplectically convex.

The proof of the second conclusion of Theorem 1.2 directly comes from Proposition 2.1, while the first conclusion needs a sophisticated argument based on ECH capacities, given in Section 1.1.

Remark 1.3. From the second conclusion of Theorem 1.2, one can estimate $p \in (0, 1]$ so that X_{Ω_p} is symplectically non-convex. Explicitly, from the inductive relation that $g(\frac{1}{k+1}) = \frac{2k+2}{2k+1} g(\frac{1}{k})$ for any positive integer k , one can show that for $p \leq \frac{1}{62460059}$, the condition (5) is satisfied and such X_{Ω_p} is not symplectically convex. Here, we emphasize that little effort was made in the proof of Theorem 1.2 to improve the estimation (5), where the estimation from Chaidez-Edtmair's criterion (7) mentioned in Remark 1.1 potentially results in a better estimation on p .

Remark 1.2. For two domains $U, V \subset \mathbb{R}^4$, we call them symplectically equivalent if $d_c(U, V) = 0$. Note that if U is symplectomorphic to V , then they are symplectically equivalent by the definition of d_c . Potentially, the symplectic equivalence relation is weaker (that is, more general) than being symplectomorphic. Back to X_{Ω_p} , even though the criterion (7) applies to verify its symplectic non-convexity,

our Theorem 1.2 in fact shows that they are not even symplectically equivalent to any convex domains when p is sufficiently small.

Proof of Theorem 1.1. (Assuming Theorem 1.2)

In proof of Theorem 1.2 in Section 2, we show that $g(p)$ increases to $+\infty$ when p goes to 0 so do the lower bound in (4). Then the desired conclusion results from the following triangular inequality,

$$d_c(X_{\Omega_p}, \mathcal{E}_4) \leq d_c(X_{\Omega_p}, \mathcal{C}_4) + \sigma(\mathcal{C}_4, \mathcal{E}_4)$$

and Proposition 2.1. □

Remark 1.3. *Consider the following pseudo-metric space*

$$(\mathcal{S}_4, d_c) = (\{\text{star-shaped domains in } \mathbb{R}^4\}, d_c).$$

From a perspective of the coarse geometry that focuses on large-scale geometrical phenomena, one could ask how many linearly independent directions in (\mathcal{S}_4, d_c) that go to infinity, usually called the rank of a quasi-flat in (\mathcal{S}_4, d_c) . The higher the rank is or the more directions there are, the richer star-shaped domains in \mathbb{R}^4 will be in terms of d_c . Immediately from the relation $\mathcal{E}_4 \subset \mathcal{S}_4$ together with Proposition 2.1, one concludes that there are at least 2 such directions, given by \mathcal{C}_4 consisting of all the symplectically convex domains.

Moreover, the classical example, 1-finger shape from Hermann in [11], shows that there exists a third direction. Here, by Theorem 1.2 and a quick observation via the comparison of Lagrangian capacity c_{Lag} (see, for instance, [5, 17]) of X_{Ω_p} and Hermann's example, the family X_{Ω_p} for $p \in (0, 1]$ indicates the existence of a new direction going to infinity in (\mathcal{S}_4, d_c) .

1.3. Differences in two criteria. As mentioned above, before this paper, the only technique that was known for proving that dynamically convex domains are not symplectically convex was Chaidez–Edtmair's criterion based on the Ruelle invariant [14]. It turns out that this criterion and the metrical-geometric approach used in this paper are quite different in essence as we now explain. To state the next result, let us introduce a notation: for any star-shaped domain $X \subset \mathbb{R}^4$ (where ∂X is a contact manifold with the contact 1-form given by $\lambda = \lambda_{\text{std}}|_{\partial X}$), define

$$(6) \quad c_{\text{CE}}(X) := \frac{\text{Ru}(\partial X) \cdot (\text{min period of a Reeb orbit of } \partial X)}{\text{Vol}(\partial X)},$$

where $\text{Ru}(\partial X)$ is the Ruelle invariant of ∂X with respect to the contact 1-form λ above (for more details, see [14]) and $\text{Vol}(\partial X)$ is the contact volume computed from the volume form $\lambda \wedge d\lambda$. Then the criterion from [3] says that there exist constants $0 < c \leq C$ such that if X is symplectically convex, then one has the following numerical constraint,

$$(7) \quad c \leq c_{\text{CE}}(X) \leq C.$$

The following result shows that the Ruelle invariant based criterion (7) and d_c provide two different perspectives to study symplectic non-convexity.

Theorem 1.4. *There exist sequences of star-shaped toric domains $\{X_{\Omega_1^k}\}$ and $\{X_{\Omega_2^k}\}$ in \mathbb{R}^4 such that*

$$\lim_{k \rightarrow \infty} \frac{c_{\text{CE}}(X_{\Omega_1^k})/c_{\text{CE}}(X_{\Omega_1^1})}{d_c(X_{\Omega_1^k}, X_{\Omega_1^1})} = 0,$$

$$\lim_{k \rightarrow \infty} \frac{c_{\text{CE}}(X_{\Omega_2^k})/c_{\text{CE}}(X_{\Omega_2^1})}{d_c(X_{\Omega_2^k}, X_{\Omega_2^1})} = \infty.$$

Remark 1.5. *Since d_c reflects the changes of symplectic capacities, Theorem 1.4 also implies that c_{CE} and symplectic capacities are independent to each other in general. It would be interesting to explore other numerical characterizations of toric domains (cf. recent work from Hutchings [15]).*

Remark 1.6 (Informed by Oliver Edtmair). *The non-toric examples constructed in [3] with small and large c_{CE} in fact lie in arbitrarily small neighborhoods of the round ball with respect to the coarse distance d_c . This shows another evidence of the independence between c_{CE} and d_c .*

Acknowledgement. The second author was partially supported by the ANR LabEx CIMI (grant ANR-11-LABX-0040) within the French State Programme ‘‘Investissements d’Avenir’’. The first and second author were partially supported by the ANR COSY (ANR-21-CE40-0002) grant. The third author was partially supported by NSF grant DMS-1926686, FAPERJ grant JCNE E-26/201.399/2021 and a Serrapilheira Institute grant. The fourth author was supported by USTC Research Funds of the Double First-Class Initiative. We are grateful for communications with Michael Hutchings, particularly concerning the second family of

examples appearing in Theorem 1.4. We also thank the hospitality of the conference - Persistence Homology in Symplectic and Contact Topology - held in Albi, France and organized by the second author, which provides an opportunity to a close collaboration between authors. Finally, we are grateful for Oliver Edtmair's crucial comments on the first version of this paper.

2. SYMPLECTIC JOHN'S ELLIPSOID THEOREM

The classical John's ellipsoid theorem [16] says that convex domains in \mathbb{R}^n are close to ellipsoids. Explicitly, denote by $\text{Conv}(\mathbb{R}^n)$ the convex domains in \mathbb{R}^n . A quantitative comparison between any two convex domains inside $\text{Conv}(\mathbb{R}^n)$ is given by the so-called Banach-Mazur distance [19],

$$(8) \quad d_{\text{BM}}(U, V) := \inf \left\{ \log \lambda \geq 0 \mid \begin{array}{l} \exists A \in \text{GL}(n), u, v \in \mathbb{R}^n \text{ such that} \\ \frac{1}{\lambda}(U + u) \subset A(V + v) \subset \lambda(U + u) \end{array} \right\}$$

where $\cdot + u$ and $\cdot + v$ denote translations while $\frac{1}{\lambda} \cdot$ and $\lambda \cdot$ stand for dilations. It is easily verified that d_{BM} defines a pseudo-metric on $\text{Conv}(\mathbb{R}^n)$ and $d_{\text{BM}}(U, V) = 0$ if and only if U is an affine transformation of V .

Since d_{BM} is defined up to affine transformations in \mathbb{R}^n , consider $\text{Ell}(n) := \{\text{ellipsoids in } \mathbb{R}^n\}$. Then John's ellipsoid theorem [16] says that for any $U \in \text{Conv}(n)$, we have

$$(9) \quad d_{\text{BM}}(U, \text{Ell}(n)) := \inf_{E \in \text{Ell}(n)} d_{\text{BM}}(U, E) \leq \frac{1}{2} \log n$$

in particular, finite.

Recall that $\sigma = \sigma_{d_c}$, defined by (2), is a symplectic version of the Banach-Mazur distance with respect to the coarse distance d_c . Recall also that \mathcal{E}_4 consists of all the symplectic ellipsoids in \mathbb{R}^4 .

Proposition 2.1 (Symplectic John's ellipsoid theorem). $\sigma(\mathcal{C}_4, \mathcal{E}_4) \leq \log 2$.

Proof. Up to symplectomorphism, let $U \in \mathcal{C}_4$ be a convex domain of \mathbb{R}^4 , by the classical John's ellipsoid theorem (9), there exists an ellipsoid $E \subset \mathbb{R}^4$ such that

$$(10) \quad E \subset U \subset o + 4 \cdot (E - o)$$

where o is the center of E . Note that this E is not necessarily a symplectic ellipsoid. However, by Williamson's theorem on standard forms for symplectic ellipsoids, there exists a (linear) symplectomorphism $\phi \in \text{Sp}(4)$ such that $\phi(E) \in$

\mathcal{E}_4 . Without loss of generality, let us assume that $\phi(E) = E(a, b)$ which is defined by

$$(11) \quad E(a, b) := \left\{ (z_1, z_2) \in \mathbb{C}^2 \mid \frac{\pi|z_1|^2}{a} + \frac{\pi|z_2|^2}{b} \leq 1 \right\}$$

for some $0 < a \leq b$. Then, since the shifting (by o) is also a symplectomorphism, the relation (10) implies that $E(a, b) \hookrightarrow U \hookrightarrow 4E(a, b)$. By a rescaling, one gets

$$\frac{1}{2}E(4a, 4b) \hookrightarrow U \hookrightarrow 2E(4a, 4b).$$

Therefore, by the definition (1), we have $d_c(U, \mathcal{E}_4) \leq \log 2$. \square

Remark 2.2. *Let us clarify the notation of rescaling $\alpha E(a, b)$ for $\alpha \in \mathbb{R}_{>0}$ that appears in the proof of Proposition 2.1. Here is the definition,*

$$\alpha E(a, b) := \left\{ (\alpha z_1, \alpha z_2) \in \mathbb{C}^2 \mid \frac{\pi|z_1|^2}{a} + \frac{\pi|z_2|^2}{b} \leq 1 \right\}$$

for $\alpha \in \mathbb{R}_{>0}$. In particular, we have an identification that $\alpha E(a, b) = E(\alpha^2 a, \alpha^2 b)$. This implies that $E(a, b) \hookrightarrow U$ if and only if $\frac{1}{\sqrt{\alpha}}E(\alpha a, \alpha b) \hookrightarrow U$.

Remark 2.3. *Proposition 2.1 holds in any $2n$ -dimensional case. More explicitly, we have $d_c(\mathcal{C}_{2n}, \mathcal{E}_{2n}) \leq \frac{1}{2} \log(2n)$.*

Remark 2.4. *The monotonicity of $\sigma(-, -)$ implies that*

$$\sigma(\mathcal{C}_4, \mathcal{M}_4) \leq \log 2$$

since $\mathcal{E}_4 \subset \mathcal{M}_4$. However, for any symplectically convex domain $U \in \mathcal{C}_4$, it is not clear how to approximate U via non-ellipsoids domains $V \in \mathcal{M}_4$. Nevertheless, since \mathcal{M}_4 contains substantially more elements than \mathcal{E}_4 , it is naturally to expect that the upper bound of $\sigma(\mathcal{C}_4, \mathcal{M}_4)$ is much lower than $\log 2$.

Interestingly, this suggests a metric-geometrical approach to prove the strong Viterbo conjecture. Namely if $\sigma(\mathcal{C}_4, \mathcal{M}_4) = 0$, then the strong Viterbo conjecture holds, due to [9, Theorem 1.7]. In fact, from the main result of a recent work [7, Theorem 1.2], the same conclusion holds in higher dimensions if $\sigma(\mathcal{C}_{2n}, \mathcal{M}_{2n}) = 0$ for $n \geq 3$.

At last, let us emphasize that this metric-geometrical approach is sensitive to the distance picked to define σ . Due to communications with M. Hutchings [15], if one replaces d_c by a rather similar pseudo-metric called symplectic Banach-Mazur distance d_{SBM} that has been studied in [10, 21, 20], then surprisingly $\sigma_{d_{\text{SBM}}}(\mathcal{C}_4, \mathcal{M}_4) >$

0. This does not immediately disprove the conjecture $\sigma(\mathcal{C}_4, \mathcal{M}_4) = 0$ above as $\sigma_{d_{\text{SBM}}}(-, -) \geq \sigma(-, -)$.

3. PROOF OF THEOREM 1.2

Denoted by c_k^{ECH} the k -th ECH capacity for $k \in \mathbb{N}$. For more background on ECH capacities, see [13]. Also, as explained in Section 1.2, it suffices to prove the first conclusion of Theorem 1.2. For simplicity, the 4-dimensional volume $\text{Vol}_{\mathbb{R}^4}(-)$ is denoted by $\text{vol}(-)$.

Proof. Suppose that $E(a, b) \hookrightarrow X_{\Omega_p} \hookrightarrow \lambda E(a, b)$ for some $E(a, b)$ and $\lambda \geq 1$. We assume without loss of generality that $a \leq b$. Note that Ω_p is a concave toric domain. For each $k \in \mathbb{N}$, let $x_k(p)$ denote the x -intercept of the line of slope $-\frac{1}{k}$, which is tangent to the curve $x^p + y^p = 1$. It follows from a straight-forward computation that

$$x_k(p) = (1 + k^{\frac{p}{p-1}})^{-\frac{1}{p}} + k(1 + k^{\frac{p}{1-p}})^{-\frac{1}{p}}.$$

Consider a certain truncation of Ω_p , defined by

$$X_k(p) = \{(x, y) \in [0, \infty)^2 \mid x^p + y^p \leq 1 \text{ and } \max(x, y) \leq x_k(p)\}$$

which is shown in Figure 3. Let (w_1, w_2, \dots) be the weight decomposition of Ω_p




FIGURE 3. $X_k(p)$ for $p \in (0, 1]$.

(see the corresponding definition in [4, Section 1.3]). Let $k = \lfloor b/a \rfloor$.

$$(12) \quad ka = c_k(E(a, b)) \leq c_k(X_{\Omega_p}) = c_k\left(\bigsqcup_{i=1}^k B(w_i)\right).$$

From [14, Equation (3.9)], we obtain

$$(13) \quad c_k \left(\bigsqcup_{i=1}^k B(w_i) \right) \leq 2 \sqrt{k \cdot \text{vol} \left(\bigsqcup_{i=1}^k B(w_i) \right)}.$$

It follows from the definition of the weight decomposition that $\bigsqcup_{i=1}^k B(w_i) \subset X_k(p)$. So

$$(14) \quad \begin{aligned} \text{vol} \left(\bigsqcup_{i=1}^k B(w_i) \right) &\leq \text{vol}(X_k(p)) = 2^{-\frac{2}{p}} + 2 \int_{2^{-\frac{1}{p}}}^{x_k(p)} (1-x^p)^{\frac{1}{p}} dx \\ &= 2^{-\frac{2}{p}} \left(1 + 2 \int_1^{2^{\frac{1}{p}} x_k(p)} (2-u^p)^{\frac{1}{p}} du \right). \end{aligned}$$

We note that the integrand in the last integral above increases as $p \rightarrow 0$ and it converges pointwise to $\frac{1}{u}$. Moreover it is readily verified that $2^{\frac{1}{p}} x_k(p)$ also increases as $p \rightarrow 0$ and it converges to $2\sqrt{k}$. So

$$(15) \quad 2 \int_1^{2^{\frac{1}{p}} x_k(p)} (2-u^p)^{\frac{1}{p}} du \leq 2 \int_1^{2\sqrt{k}} \frac{1}{u} du = 2 \log(2\sqrt{k}) = \log 4 + \log k.$$

Combining (12), (13), (14) and (15), we obtain $ka \leq 2 \cdot 2^{-\frac{1}{p}} \sqrt{k(1 + \log 4 + \log k)}$. So

$$(16) \quad a \leq 2 \cdot 2^{-\frac{1}{p}} \sqrt{\frac{1 + \log 4 + \log k}{k}}.$$

Since $X_{\Omega_p} \hookrightarrow \lambda E(a, b)$, we have

$$(17) \quad 2 \cdot 2^{-\frac{1}{p}} = c_1^{\text{ECH}}(X_{\Omega_p}) \leq \lambda^2 c_1^{\text{ECH}}(E(a, b)) = \lambda^2 a.$$

It follows from (16) and (17) that $a \leq \lambda^2 a \sqrt{\frac{1 + \log 4 + \log k}{k}}$. So

$$(18) \quad \lambda^4 \geq \frac{k}{1 + \log 4 + \log k}.$$

Note that this estimation is not enough to deduce the requested large d_c -distance conclusion since when p changes, the constant k (so that the corresponding ellipsoid $E(a, b)$ with $k = \lfloor b/a \rfloor$ that embeds into X_{Ω_p}) may change.

Now, from (16), we also obtain the following estimations,

$$\begin{aligned}
a &\leq \sqrt{\frac{(1 + \log 4 + \log k) \operatorname{vol}(X_{\Omega_p})}{k \cdot g(p)}} \\
&\leq \sqrt{\frac{(1 + \log 4 + \log k) \operatorname{vol}(\lambda E(a, b))}{k \cdot g(p)}} \\
&= \sqrt{\frac{(1 + \log 4 + \log k) \lambda^4 ab}{2k \cdot g(p)}} \\
&\leq a \sqrt{\frac{(1 + \log 4 + \log k) \lambda^4 (k + 1)}{2k \cdot g(p)}} \\
&\leq a \sqrt{\frac{(1 + \log 4 + \log k) \lambda^4}{g(p)}}
\end{aligned}$$

where the final step comes from the estimation $\frac{k+1}{2k} \leq 1$ for all $k \in \mathbb{N}$. So, we get

$$(19) \quad \lambda^4 \geq \frac{g(p)}{1 + \log 4 + \log k}.$$

From (18) and (19) it follows that

$$(20) \quad \lambda^4 \geq \frac{\max(g(p), k)}{1 + \log 4 + \log k}.$$

From a change of variables we obtain

$$\begin{aligned}
g(p) &= 2^{\frac{2}{p}-2} \operatorname{vol}(X_{\Omega_p}) = 2^{\frac{2}{p}-2} \int_0^1 (1 - x^p)^{\frac{1}{p}} dx \\
&= \frac{1}{4} \int_0^{2^{\frac{1}{p}}} (2 - u^p)^{\frac{1}{p}} du.
\end{aligned}$$

One observes that $\lim_{p \rightarrow 0} (2 - u^p)^{\frac{1}{p}} = \frac{1}{u}$. Therefore, $g(p)$ increases as $p \rightarrow 0$ and $\lim_{p \rightarrow 0} g(p) = \infty$. Moreover,

$$g\left(\frac{1}{5}\right) = \frac{\Gamma(1 + 5)^2}{\Gamma(1 + 2 \cdot 5)} \cdot 2^{2 \cdot 5 - 2} = \frac{5!^2}{10!} \cdot 2^8 = \frac{64}{63} > 1.$$

So for $p < \frac{1}{5}$, it follows that $\lfloor g(p) \rfloor \geq 1$. In this case,

$$\begin{aligned}
\min_k \frac{\max(g(p), k)}{1 + \log 4 + \log k} &= \min \left(\frac{g(p)}{1 + \log 4 + \log \lfloor g(p) \rfloor}, \frac{\lfloor g(p) \rfloor + 1}{1 + \log 4 + \log(\lfloor g(p) \rfloor + 1)} \right) \\
&\geq \frac{g(p)}{1 + \log 4 + \log g(p)}.
\end{aligned}$$

Taking the infimum of over all triples (λ, a, b) , we obtain

$$f(p)^4 \geq \frac{g(p)}{1 + \log 4 + \log g(p)}$$

where $f(p) = \inf\{\lambda \geq 1 \mid E(a, b) \hookrightarrow X_{\Omega_p} \hookrightarrow \lambda E(a, b) \text{ for some } a, b > 0\}$. Then one can show that $f(p)$ is linked to the coarse distance d_c by

$$d_c(X_{\Omega_p}, \mathcal{E}_4) = \frac{1}{2} \log f(p)$$

thus we obtain (4), proving the desired conclusion. \square

Here, we point out that if we “linearize” the profile curve in Ω_p of X_{Ω_p} considered in the proof of Theorem 1.2 above, denoted by Ω_p^{lin} and shown in Figure 4, then following conclusion holds.




FIGURE 4. A linear approximation of X_{Ω_p} from Ω_p^{lin} .

Proposition 3.1. *For any $p \in (0, 1]$, we have $d_c(X_{\Omega_p^{\text{lin}}}, \mathcal{C}_4) \leq \frac{1}{2} \log 3$.*

Proof. On the one hand, by inclusion, the ellipsoid $E(1, \frac{2^{-1/p}}{1-2^{-1/p}})$ shown by the bold edges in Figure 4 embeds inside $X_{\Omega_p^{\text{lin}}}$. On the other hand, by Lemma 4.5 in [9], $X_{\Omega_p^{\text{lin}}} \hookrightarrow P(1, 2^{-1/p+1})$. Since $P(1, 2^{-1/p+1}) \subset (3 - 2^{-1/p+1})E(1, \frac{2^{-1/p}}{1-2^{-1/p}})$ trivially, one gets the following relation,

$$E\left(1, \frac{2^{-\frac{1}{p}}}{1 - 2^{-\frac{1}{p}}}\right) \hookrightarrow X_{\Omega_p^{\text{lin}}} \hookrightarrow 3E\left(1, \frac{2^{-\frac{1}{p}}}{1 - 2^{-\frac{1}{p}}}\right)$$

This implies that

$$d_c(X_{\Omega_p^{\text{lin}}}, \mathcal{E}_4) \leq \frac{1}{2} \log 3$$

which completes the proof. \square

Remark 3.2. Here is a way to explain the essential difference between Theorem 1.1 and Proposition 3.1, by simply comparing the respective volumes, where

$$\frac{\text{Vol}_{\mathbb{R}^4}(X_{\Omega_p^{\text{lin}}})}{\text{Vol}_{\mathbb{R}^4}(X_{\Omega_p})} \rightarrow +\infty$$

as $p \rightarrow 0$.

4. PROOF OF THEOREM 1.4

Proof. The proof is given by two different examples.

Example One: Let us consider the strangulation operation on $B^4(1)$ or for brevity $B(1)$. By definition, it cuts off a part of $B^4(1)$, which is, on the level of the moment image, a thin triangle symmetric to the diagonal $y = x$. See Figure 5, where $\epsilon(\delta)$ is proportional to δ , Denote the resulting toric domain by B_δ . Then




FIGURE 5. A strangulation on $B(1)$, resulting a toric domain B_δ .

obviously we have $B_\delta \hookrightarrow B(1)$ simply by inclusion. Now, cut through the diagonal and taking closures, we obtain two toric domains B_δ^- and B_δ^+ (which are in fact symplectomorphic to each other by symmetry). Apply the affine transformation

$$\begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix},$$

then the moment image of B_δ^- transfers to a region E_T where the corresponding toric domain (still denoted by) E_T is an example of so-called *truncated ellipsoid* defined in [21]. More explicitly, by an elementary calculation, it is a truncated ellipsoid from $E\left(1, \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right)$, with truncated parameters δ and $\frac{\frac{1}{2}+\epsilon(\delta)-\delta}{2\epsilon(\delta)}$. These parameters are denoted by ϵ and β respectively in [21]. For our purpose, let us denote

$$(21) \quad \beta(\delta) := \frac{\frac{1}{2} + \epsilon(\delta) - \delta}{2\epsilon(\delta)}.$$

Since affine transformations on moment images induce symplectomorphisms on the corresponding toric domains, we know that

$$(22) \quad (E_T)^- \sqcup (E_T)^+ \simeq B_\delta^- \sqcup B_\delta^+ \hookrightarrow B_\delta$$

where $(E_T)^-$ and $(E_T)^+$ stand for two copies of the truncated ellipsoid above. One of the more striking properties of truncated ellipsoids is Corollary 5.3 in [21], which proves that

$$(23) \quad d_c\left(E_T^-, E\left(1, \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right)\right) \leq 2 \log\left(\frac{1+\beta(\delta)}{\beta(\delta)}\right) := \log C(\delta).$$

Hence, with all the notations above, we have

$$\begin{aligned} \frac{1}{C(\delta)} B\left(\frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) &\hookrightarrow \frac{1}{C(\delta)} B\left(\frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) \sqcup \frac{1}{C(\delta)} B\left(\frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) \\ &\hookrightarrow \frac{1}{C(\delta)} E\left(1, \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) \sqcup \frac{1}{C(\delta)} E\left(1, \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) \\ &\hookrightarrow (E_T)^- \sqcup (E_T)^+ \hookrightarrow B_\delta \end{aligned}$$

where the first embedding is just the inclusion; the second embedding comes from the Gromov width of an ellipsoid; the third embedding comes from (23); the last embedding is just (22). To sum up, we obtain the following embedding relations,

$$B\left(\frac{1}{C(\delta)^2} \cdot \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)}\right) \hookrightarrow B_\delta \hookrightarrow B(1).$$

Then when $\delta \rightarrow 0$, we have

$$\epsilon(\delta) \rightarrow 0, \text{ so } \frac{1+2\epsilon(\delta)}{2-4\epsilon(\delta)} \rightarrow \frac{1}{2}; \quad \beta(\delta) \rightarrow +\infty, \text{ so } C(\delta) \rightarrow 1.$$

Therefore, $d_c(B(1), B_\delta) \rightarrow \log \sqrt{2}$ as $\delta \rightarrow 0$. In particular, $B(1)$ and B_δ do not differ much in terms of any symplectic capacity. However, by the argument in [8, Section 3.1], we know $c_{\text{CE}}(B_\delta) \rightarrow 0$ as $\delta \rightarrow 0$.

Example Two: Let us consider the strain operation on $B^4(99)$ or for brevity $B(99)$. By definition, we add two small “tail” triangles along each axis on the moment image of $B(99)$. See Figure 6, where small triangles are shaded, they are symmetric with respect to $y = x$, and the horizontal one has height 1 and base 99. Denote the resulting domain by X . Note that




FIGURE 6. A strain on $B(99)$, resulting a toric domain X .

$$\text{Vol}_{\mathbb{R}^4}(X) = \frac{99^2}{2} + 2 \cdot \frac{99}{2} < \frac{100^2}{2} = \text{Vol}_{\mathbb{R}^4}(B(100)).$$

Then obviously we have $B(99) \hookrightarrow X$. Now, we claim that $X \hookrightarrow B(100)$. Since X is concave toric, $B(100)$ is convex toric, and the weight sequence of X is

$$(w_1, w_2, \dots) = (99, \underbrace{1, \dots, 1}_{198\text{-many}})$$

such required embedding can be transferred to a ball-packing problem by [6, Theorem 2.1]. In fact, we claim that

$$B(99) \cup \underbrace{B(1) \cup \dots \cup B(1)}_{199\text{-many}} \hookrightarrow B(100).$$

Indeed, the shaded tail triangles in X (for instance, the horizontal one) can be divided into small triangles where each is equivalent to the triangle with vertices $(98, 0), (99, 0), (98, 1)$, up to affine transformation. For an illustration, see the left picture in Figure 7. Then by further affine transformations, we can move the




FIGURE 7. Packing the tail triangles of X into $B(100)$.

“start” triangle into $B(100) \setminus B(99)$, on the level of their moment image denoted by $\Delta(100) \setminus \Delta(99)$. The right picture in Figure 7 shows how the small triangle, shaded one with vertices $(100, 0), (101, 0), (98, 1)$, is placed inside $\Delta(100) \setminus \Delta(99)$, precisely located at the shaded small triangle with vertices $(99, 1), (100, 1), (99, 2)$. Since

$$\text{Area}_{\mathbb{R}^2}(\Delta(100) \setminus \Delta(99)) = \frac{100^2}{2} - \frac{99^2}{2} = \frac{199}{2} = 199 \cdot \text{Area}_{\mathbb{R}^2}(\text{small } \Delta),$$

such procedure can be inductively carried out and pack 199-many small Δ into $\Delta(100) \setminus \Delta(99)$. This corresponds to a packing of 199-many $B(1)$ into $B(100) \setminus B(99)$. Thus we obtain the desired claim.

Note that the same argument works identically if we replace the tail triangle in X by one with the furthest vertex being $(k, 0)$ (but with volume fixed). Denote the resulting toric domain by X_k , then we complete the proof by the argument in [8, Section 3.2] where $c_{\text{CE}}(X_k) \rightarrow \infty$ when $k \rightarrow \infty$. \square

REFERENCES

1. Shiri Artstein-Avidan, Roman Karasev, and Yaron Ostrover, *From symplectic measurements to the Mahler conjecture*, *Duke Math. J.* **163** (2014), no. 11, 2003–2022. MR 3263026
2. Julian Chaidez and Oliver Edtmair, *Convexity and the Ruelle invariant in higher dimensions*, arXiv preprint, arXiv: 2205.00935.
3. ———, *3D convex contact forms and the Ruelle invariant*, *Invent. Math.* **229** (2022), no. 1, 243–301. MR 4438355
4. Keon Choi, Michael Hutchings, Daniel Cristofaro-Gardiner, David Frenkel, and Vinicius Gripp Barros Ramos, *Symplectic embeddings into four-dimensional concave toric domains*, *Journal of Topology* **7** (2014), no. 4, 1054–1076.
5. K. Cieliebak and K. Mohnke, *Punctured holomorphic curves and Lagrangian embeddings*, *Invent. Math.* **212** (2018), no. 1, 213–295. MR 3773793
6. Dan Cristofaro-Gardiner, *Symplectic embeddings from concave toric domains into convex ones*, *Journal of Differential Geometry* **112** (2019), no. 2, 199–232.
7. Dan Cristofaro-Gardiner and Richard Hind, *On the agreement of symplectic capacities in high dimension*, arXiv preprint, arXiv: 2307.12125.
8. Julien Dardennes, Jean Gutt, and Jun Zhang, *Symplectic non-convexity of toric domains*, *Communications in Contemporary Mathematics*, <https://doi.org/10.1142/S0219199723500104> (2023).
9. Jean Gutt, Michael Hutchings, and Vinicius G. B. Ramos, *Examples around the strong Viterbo conjecture*, *J. Fixed Point Theory Appl.* **24** (2022), no. 2, Paper No. 41, 22. MR 4413022
10. Jean Gutt and Michael Usher, *Symplectically knotted codimension-zero embeddings of domains in \mathbb{R}^4* , *Duke Math. J.* **168** (2019), no. 12, 2299–2363. MR 3999447
11. David Hermann, *Non-equivalence of symplectic capacities for open sets with restricted contact type boundary*, Can be found at https://www.imo.universite-paris-saclay.fr/~biblio/pub/1998/abs/ppo1998_32.html.
12. Helmut Hofer, Krzysztof Wysocki, and Eduard Zehnder, *The dynamics on three-dimensional strictly convex energy surfaces*, *Ann. of Math. (2)* **148** (1998), no. 1, 197–289. MR 1652928
13. Michael Hutchings, *Quantitative embedded contact homology*, *J. Differential Geom.* **88** (2011), no. 2, 231–266. MR 2838266
14. ———, *ECH capacities and the Ruelle invariant*, *J. Fixed Point Theory Appl.* **24** (2022), no. 2, Paper No. 50, 25. MR 4441526
15. ———, *Private communications*, 2023.
16. Fritz John, *Extremum problems with inequalities as subsidiary conditions*, *Studies and Essays Presented to R. Courant on his 60th Birthday, January 8, 1948*, Interscience Publishers, Inc., New York, 1948, pp. 187–204. MR 0030135
17. Miguel Pereira, *Equivariant symplectic homology, linearized contact homology and the lagrangian capacity*, PhD thesis, University of Augsburg, May 2022.

18. Paul H. Rabinowitz, *Periodic solutions of Hamiltonian systems*, Comm. Pure Appl. Math. **31** (1978), no. 2, 157–184. MR 467823
19. Mark Rudelson, *Distances between non-symmetric convex bodies and the MM^* -estimate*, Positivity **4** (2000), no. 2, 161–178. MR 1755679
20. Vukašin Stojisavljević and Jun Zhang, *Persistence modules, symplectic Banach-Mazur distance and Riemannian metrics*, Internat. J. Math. **32** (2021), no. 7, Paper No. 2150040, 76. MR 4284596
21. Michael Usher, *Symplectic Banach-Mazur distances between subsets of \mathbb{C}^n* , J. Topol. Anal. **14** (2022), no. 1, 231–286. MR 4411106
22. Claude Viterbo, *Metric and isoperimetric problems in symplectic geometry*, J. Amer. Math. Soc. **13** (2000), no. 2, 411–431. MR 1750956

Email address: `julien.dardennes@math.univ-toulouse.fr`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TOULOUSE, 118 ROUTE DE NARBONNE,
31062 TOULOUSE CEDEX 9, FRANCE

Email address: `jean.gutt@math.univ-toulouse.fr`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TOULOUSE, 118 ROUTE DE NARBONNE,
31062 TOULOUSE CEDEX 9, FRANCE AND INU CHAMPOLLION, PLACE DE VERDUN, 81000
ALBI, FRANCE

Email address: `vgbramos@impa.br`

INSTITUTO DE MATEMÁTICA PURA E APLICADA, ESTRADA DONA CASTORINA, 110, RIO
DE JANEIRO - RJ -BRASIL, 22460-320

Email address: `jzhang4518@ustc.edu.cn`

THE INSTITUTE OF GEOMETRY AND PHYSICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY
OF CHINA, 96 JINZHAI ROAD, HEFEI ANHUI, 230026, CHINA