



HAL
open science

3D models related to the publication: A 50-million-year-old, three-dimensionally preserved bat skull supports an early origin for modern echolocation

Jacob Mougoust, Maëva Judith Orliac

► **To cite this version:**

Jacob Mougoust, Maëva Judith Orliac. 3D models related to the publication: A 50-million-year-old, three-dimensionally preserved bat skull supports an early origin for modern echolocation. *MorphoMuseum*, 2023, 9 (4), pp.e217. <10.18563/journal.m3.217>. <hal-04934115>

HAL Id: hal-04934115

<https://hal.science/hal-04934115v1>

Submitted on 7 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

3D models related to the publication: A 50-million-year-old, three-dimensionally preserved bat skull supports an early origin for modern echolocation

Jacob Mougoust^{1*}, Maëva Judith Orliac¹

¹Institut des Sciences de l'Évolution de Montpellier (UMR 5554, CNRS, UM, IRD, EPHE), c.c. 064, Université Montpellier, place Eugène Bataillon, F-34095 Montpellier Cedex.

*Corresponding author: jacob.mougoust@gmail.com

Abstract

The present 3D Dataset contains 3D models of the cranium surface and of the bony labyrinth endocast of the stem bat *Vielasia sigei*. They are used by (Hand et al., 2023) to explore the phylogenetic position of this species, to infer its laryngeal echolocating capabilities, and to eventually discuss chiropteran evolution before the crown clade diversification.

Keywords: Bony labyrinth, Chiroptera, Cranium, Eocene, Paleontology

Submitted:2023-09-27, published online:2023-10-19. <https://doi.org/10.18563/journal.m3.217>

INTRODUCTION

Bats are one of the most speciose and ecologically diverse mammalian orders (Burgin et al., 2018; Simmons and Cirranello, 2023). They are additionally morphologically highly divergent in their skeletons and skulls: they are the only mammals to use powered flight and many species use echolocation calls (Teeling et al., 2016; Anderson and Ruxton, 2020). It is yet undetermined whether flight and echolocation appeared and evolved together (Speakman, 2001) or whether one was acquired before the other (Norberg and Rayner, 1987; Teeling et al., 2005; Simmons et al., 2008). Paleontological evidences of flight acquisition have been demonstrated in stem bats, a paraphyletic assemblage of early-diverging bats (Simmons et al., 2008; Smith et al., 2012; Rietbergen et al., 2023); echolocation ability has been inferred in such forms, but more tentatively (Simmons and Geisler, 1998; Simmons et al., 2008; Veselka et al., 2010). On the other hand, ontogenetic observations recently highlighted diverging developmental patterns to obtain a similar structure, suggesting distinct origins of distinct adaptations toward a similar ability (Nojiri et al., 2021). Hence, echolocation ability might have been acquired once, before the crown bat clade diversification, or in parallel across crown and stem bats (Anderson and Ruxton, 2020). Hand et al., (2023) describe a new early Eocene stem bat from French Quercy Phosphorites, *Vielasia sigei*, with unprecedented material including the oldest uncrushed, three-dimensionally preserved, bat cranium yet found (Fig. 1). Accompanied by several dental, cranial, and postcranial specimens representing at least 23 individuals, this cranium helps in placing *V. sigei* outside modern bats in a total evidence tip-dating Bayesian phylogeny. Furthermore, the *in situ* preservation of one petrosal allows for reconstructing the endocast of its right bony labyrinth (Fig. 2). It is of high anatomical and morphometric similarity to that of other, crown, laryngeally-echolocating bats. The results suggest that *V. sigei* is a stem bat that used a similar kind of laryngeal echolocation as the one used by most modern bats, supporting

Inv nr.	Description
UM VIE-250_M3#1269	External surface of the cranium
UM VIE-250_M3#1270	Virtual endocast of the right bony labyrinth

Table 1. List of models of the holotype of *Vielasia sigei* (Inv. nr: UM VIE-250). Collection: Institut des Sciences de l'Évolution de Montpellier (ISEM), Université de Montpellier, Montpellier, France.

the hypothesis of a unique origin of this trait for crown bats.

METHODS

Three dimensional data acquisition from fossils was performed using the μ CT facility (EasyTom 150 μ CT) at the University of Montpellier (MRI, ISEM) with 90 kV and 80 μ A at a resolution of 8.93 μ m. A numeric binning has been performed prior to the segmentation, and the models were extracted from a μ CT dataset with a resolution of 17.86 μ m. Segmentation was performed using Avizo @ 9.3 (Thermo Fisher Scientific-FEI) and its "threshold", "lasso", and "brush" tools; visualization was performed using MorphoDig@ (Lebrun, 2018). Of a single physical specimen (UM VIE-250) housed in the Université de Montpellier collections of paleontology, two virtual specimens have been created: the external surface of the cranium (Fig. 1, Table 1), and the endocast of the right bony labyrinth (Fig. 2, Table 1). Further works are being conducted for internal structures of this specimen; therefore, the cranium surface has been filled, automatically (filling holes in all three orthogonal plans) and manually (filling the braincase and ducts around the nasal cavity, and unfilling holes and pits in the hard palate, around the pterygoid wings, and in the teeth).

ACKNOWLEDGEMENTS

We warmly thank Anne-Lise Charruault who successfully performed the μ CT-scanning of this extremely delicate specimen, a mandatory prerequisite for this study. We are grateful to the




Figure 1. Digitized external surface of the holotype of *Vielasia sigei* (UM VIE-250) in dorsal (A), ventral (C), left (B), and right (D) views. Scale bar corresponds to 5 mm.




Figure 2. Virtual endocast of the right bony labyrinth of the holotype of *Vielasia sigei* (UM VIE-250) in dorsal (A), ventral (D), anterior (B), posterior (E), lateral (C), and medial (F) views. The orientation of the bony labyrinth follows that of the corresponding cranium. Scale bar corresponds to 1 mm..

MRI platform member of the national infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04, «Investments for the future»), the labex CEMEB (ANR-10-LABX-0004) and NUMEV (ANR-10-LABX-0020). This work was financially supported by a PhD thesis funding from the French Ministry of Higher Education, Research, and Innovation and by the ANR program ENLIVEN (ANR-22-CE2-0014-01) headed by M. J. Orliac.

BIBLIOGRAPHY

- Anderson, S.C., Ruxton, G.D., 2020. The evolution of flight in bats: a novel hypothesis. *Mammal Review*. 50, 426–439. <https://doi.org/10.1111/mam.12211>.
- Burgin, C.J., Colella, J.P., Kahn, P.L., Upham, N.S., 2018. How many species of mammals are there? *Journal of Mammalogy*. 99, 1–14. <https://doi.org/10.1093/jmammal/gyx147>.
- Hand, S.J., Maugoust, J., Beck, R.M.D., Orliac, M.J., 2023. A 50-million-year-old, three-dimensionally preserved bat skull supports an early origin for modern echolocation. *Current Biology*. <https://doi.org/10.1016/j.cub.2023.09.043>
- Lebrun, R., 2018. MorphoDig, an open-source 3D freeware dedicated to biology.
- Nojiri, T., Wilson, L.A.B., López-Aguirre, C., Tu, V.T., Kuratani, S., Ito, K., Higashiyama, H., Son, N.T., Fukui, D., Sadier, A., Sears, K.E., Endo, H., Kamihori, S., Koyabu, D., 2021. Embryonic evidence uncovers convergent origins of laryngeal echolocation in bats. *Current Biology*. 31, 1353–1365.e3. <https://doi.org/10.1016/j.cub.2020.12.043>.
- Norberg, U.M.L., Rayner, J., 1987. Ecological morphology and flight in bats (Mammalia; Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*. 316, 335–427. <https://doi.org/10.1098/rstb.1987.0030>.
- Rietbergen, T.B., van den Hoek Ostende, L.W., Aase, A., Jones, M.F., Medeiros, E.D., Simmons, N.B., 2023. The oldest known bat skeletons and their implications for Eocene chiropteran diversification. *PLOS ONE*. 18, e0283505. <https://doi.org/10.1371/journal.pone.0283505>.
- Simmons, N.B., Cirranello, A.L., 2023. Bat species of the world: a taxonomic and geographic database [WWW Document]. URL <https://batnames.org/> (accessed 4.17.23).
- Simmons, N.B., Geisler, J.H., 1998. Phylogenetic relationships of *Icaronycteris*, *Archaeonycteris*, *Hassianycteris*, and *Palaeochiropteryx* to extant bat lineages, with comments on the evolution of echolocation and foraging strategies in Microchiroptera. *Bulletin of the American Museum of Natural History*. 1998, 1–182.
- Simmons, N.B., Seymour, K.L., Habersetzer, J., Gunnell, G.F., 2008. Primitive Early Eocene bat from Wyoming and the evolution of flight and echolocation. *Nature*. 451, 818–821. <https://doi.org/10.1038/nature06549>.
- Smith, T., Habersetzer, J., Simmons, N.B., Gunnell, G.F., 2012. Systematics and paleobiogeography of early bats. In: Gunnell, G.F., Simmons, N.B. (Eds.), *Evolutionary History of Bats*. Cambridge University Press, Cambridge, pp. 23–66. <https://doi.org/10.1017/CBO9781139045599.003>.
- Speakman, J.R., 2001. The evolution of flight and echolocation in bats: another leap in the dark. *Mammal Review*. 31, 111–130. <https://doi.org/10.1046/j.1365-2907.2001.00082.x>.
- Teeling, E.C., Jones, G., Rossiter, S.J., 2016. Phylogeny, Genes, and Hearing: Implications for the Evolution of Echolocation in Bats. In: Fenton, M.B., Grinnell, A.D., Popper, A.N., Fay, R.R. (Eds.), *Bat Bioacoustics, Springer Handbook of Auditory Research*. Springer New York, New York, NY, pp. 25–54. https://doi.org/10.1007/978-1-4939-3527-7_2.
- Teeling, E.C., Springer, M.S., Madsen, O., Bates, P.J.J., O'Brien, S.J., Murphy, W.J., 2005. A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science*. 307, 580–584. <https://doi.org/10.1126/science.1105113>.
- Veselka, N., McErlain, D.D., Holdsworth, D.W., Eger, J.L., Chhem, R.K., Mason, M.J., Brain, K.L., Faure, P.A., Fenton, M.B., 2010. A bony connection signals laryngeal echolocation in bats. *Nature*. 463, 939–942. <https://doi.org/10.1038/nature08737>.