



**HAL**  
open science

## **Sexual reproduction is controlled by successive transcriptomic waves in *Podospora anserina***

Frederique Bidard, Pierre Grognet, Gaëlle Lelandais, Sandrine Imbeaud, Marie-Hélène Muchielli, Robert Debuchy, Véronique Berteaux-Lecellier, Fabienne Malagnac

► **To cite this version:**

Frederique Bidard, Pierre Grognet, Gaëlle Lelandais, Sandrine Imbeaud, Marie-Hélène Muchielli, et al.. Sexual reproduction is controlled by successive transcriptomic waves in *Podospora anserina*. 2025. hal-04941892

**HAL Id: hal-04941892**

**<https://hal.science/hal-04941892v1>**

Preprint submitted on 12 Feb 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Sexual reproduction is controlled by**  
2 **successive transcriptomic waves in**  
3 ***Podospora anserina***

4

5 Frédérique Bidard<sup>1,2, §</sup>, Pierre Grognet<sup>3, §</sup>, Gaëlle Lelandais<sup>3, §</sup>, Sandrine  
6 Imbeaud<sup>4, 5</sup>, Marie-Hélène Muchielli<sup>4, 6</sup>, Robert Debuchy<sup>1, 3</sup>, Véronique  
7 Berteaux-Lecellier<sup>1, 7, #</sup> and Fabienne Malagnac<sup>3, #, \*</sup>.

8

9 <sup>1</sup> Univ Paris-Sud 11, CNRS, Institut de Génétique et Microbiologie UMR8621, F- 91405 Orsay,  
10 France.

11 <sup>2</sup> IFP Energies Nouvelles, 1 et 4 avenue de Bois-Préau, 92852Rueil-Malmaison, France

12 <sup>3</sup> Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC),  
13 91198, Gif-sur-Yvette, France

14 <sup>4</sup> CNRS, Centre de Génétique Moléculaire, FRE3144, GODMAP, 91190 Gif sur Yvette, France.

15 <sup>5</sup> present address : INSERM, Génomique Fonctionnelle des tumeurs solides, UMR U-674, IUH,  
16 Université Paris Descartes, 75010 Paris, France.

17 <sup>6</sup> present address : Institut de Sciences des Plantes - Paris-Saclay, Bâtiment 630, Avenue des  
18 Sciences, 91190 Gif-sur-Yvette, France

19 <sup>7</sup> present address: UMR250/9220 ENTROPIE (CNRS-IRD-UR-IFREMER-UNC), Promenade  
20 Roger-Laroque, Noumea Cedex, New Caledonia, France

21

22 **§ Contributed equally**

23 **# Contributed equally**

24 **\*Corresponding author**

25 Correspondence: [fabienne.malagnac@universite-paris-saclay.fr](mailto:fabienne.malagnac@universite-paris-saclay.fr)

26

27

28 **Abstract**

29 Despite the inherent challenge of finding suitable mating partners, most eukaryotes use sexual  
30 reproduction to produce offspring endowed with increase genetic diversity and fitness. The  
31 persistence of this mode of reproduction is a key question in evolutionary biology. Fungi offer  
32 valuable insights into this question, due to their diversity and short lifecycle. This study focuses  
33 on *Podospora anserina*, a pseudohomothallic ascomycete fungus that bypasses self-sterility by  
34 maintaining two compatible nuclei in one mycelium. We performed genome-wide gene  
35 expression profiling during ten stages of *P. anserina* sexual reproduction and identified five  
36 major expression patterns. Our expert annotation approaches identified differentially expressed  
37 genes related to secondary metabolite production, fungal vegetative incompatibility,  
38 programmed cell death, and epigenetic regulation. In addition, master transcriptional regulators  
39 and their target networks were uncovered. This study provides a comprehensive database for  
40 future functional genomics experiments and novel pathway characterization during sexual  
41 reproduction.

42

43 **Keywords:** transcriptome, time course, sexual differentiation, filamentous fungi

44 **Running head:** Transcriptome landscape during fungal sexual development

45

## 46 **Introduction**

47 Most eukaryotes rely on sexual reproduction to produce offspring. However, the  
48 maintenance of sex remains a fundamental question in evolutionary biology. The amazing  
49 diversity of fungi makes them of particular interest for addressing this question. Fungi are found  
50 in almost every terrestrial and marine ecosystems (Stajich et al. 2009; Breyer et Baltar 2023).  
51 Some play key roles in biogeochemical cycles, while others possess traits that enable them to  
52 be pathogenic to plants or animals (Iliev et al. 2024). Understanding the biology of fungi, and  
53 in particular why many of them can produce either asexual or meiotic products depending on  
54 environmental conditions (Dyer et Kück 2017; Wang et al. 2018), is crucial for fields as diverse  
55 as ecology, agronomy and human health.

56 For fungi, as for all eukaryotes, sexual reproduction has well-documented advantages. In  
57 particular, meiosis, the sex-specific division, reduces the accumulation of deleterious mutations  
58 and can link beneficial alleles that are subsequently fixed in populations by natural selection  
59 (Otto 2009). In harsh environments, sexual reproduction, as opposed to asexual reproduction,  
60 can lead to improved fitness of the offspring.

61 Fungal sexual reproduction relies on a wide range of developmental strategies and mating  
62 systems. These include both inbreeding, in which a single individual can self-fertilize, and  
63 obligate outbreeding, in which only genetically distinct compatible partners can mate. Some  
64 fungi, referred to as pseudohomothallic, have managed to retain the best of each strategie.  
65 Pseudohomothallism extends the diversity of reproductive systems evolved in fungi by  
66 selecting breeding systems that allow self-sterility to be bypassed by maintaining two  
67 populations of compatible nuclei in the same mycelium (Peraza-Reyes et Malagnac 2016).  
68 Early forward genetic screens and more recent transcriptomic studies in Sordariomycetes  
69 (Wang et al. 2014; Lütkenhaus et al. 2019; Kim et al. 2022) have led to the identification of  
70 large sets of genes involved in specialized developmental pathways that control the fertilization  
71 and the subsequent developmental stages (Wilson et al. 2019). Some of these species were  
72 heterothallic (self-incompatible, self-sterile (Wang et al. 2014; Lütkenhaus et al. 2019)), while  
73 others were homothallic (self-compatible, self-fertile (Teichert et al. 2012; Wang et al. 2019)).  
74 However, how the development of meiotic tissues within the fruiting body is controlled and  
75 which genes are critical in this process remains unknown.

76 In this study, we propose to approach this question from a different angle using the  
77 pseudohomothallic species *Podospora anserina*. This species is used as a model system for the  
78 study of the fruiting body development (Peraza-Reyes et Malagnac 2016) but a detailed analysis

79 of the cellular programs allowing its formation from fertilization to ascospore discharge was  
80 lacking. In this context, we performed a genome-wide gene expression profiling during the  
81 sexual reproduction of *P. anserina* at ten time points representing key developmental steps. To  
82 characterize the topology of differential gene expression profiles across sexual reproduction,  
83 we analyzed our transcriptomic data using two complementary approaches: *i*) a data-driven  
84 approach using k-means clustering and *ii*) a biological expertise-driven approach to mine the  
85 gene functions. The data-driven approach revealed five major patterns (waves) of differential  
86 expression during sexual development. The expert annotation approach has led to the  
87 identification of a number of interesting gene categories, including secondary metabolites and  
88 toxins encoding genes (Wang et al. 2022), genes responsible for fungal vegetative  
89 incompatibility programmed cell death (Chou, Hsu, et Leu 2022), and gene related to epigenetic  
90 effectors that interfere with messenger stability (RNAi/MSUD genes, (Gladyshev 2017)).  
91 Finally, we identified several master transcriptional regulators and draw their potential target  
92 gene network. Overall, our study has produced a comprehensive database that offers numerous  
93 candidates for functional genomics experiments. This database also paves the way for the  
94 characterization of new pathways during sexual reproduction.

## 95 **Materials and methods**

### 96 ***Podospora anserina* strains and methods**

97 *P. anserina* is a filamentous ascomycete whose life cycle and general methods have  
98 already been described (Esser, 1974; Rizet and Engelmann, 1949; Zickler et al., 1995). All  
99 strains used in this study were derived from the *S* strain. Growth and crosses of *S mat+* and *S*  
100 *mat-* strains were carried out on a minimal synthetic medium (M2 medium) at 27°C with  
101 constant light.

### 102 **Cytology**

103 Sexual cells were either fixed in 7.4% paraformaldehyde and processed for microscopy  
104 as previously described (Thompson-Coffe and Zickler, 1994) or directly observed in M2  
105 medium. When required, nuclei were stained with DAPI (4',6-diamidino-2-phenylindole, 1  
106 mg/mL; Boehringer Ingelheim). Observations were performed with a Zeiss Axioplan  
107 microscope, and images captured with a CDD Princeton camera system.

### 108 **Time course procedure and RNA isolation**

109 To facilitate biological materials collecting, *S* strains from both mating types were  
110 grown on cheesecloth (Sefar Nitex 03-48/31 Heiden) or cellophane sheet (cat#1650193,  
111 BioRad laboratories, Hercules) previously placed on Petri dishes containing M2 medium. *S*  
112 *mat*<sup>-</sup> strain has been used as male gamete donor and *S mat*<sup>+</sup> as female. The *S mat*<sup>+</sup> strains were  
113 grown on M2 medium plus cheesecloth except the T-24h (on cellophane sheet). For each ten  
114 points of the time course, the number of Petri dishes has been adjusted (five were used for T-  
115 24h, T0h, T6h, T18h, T24h, four for T30h, two for T42h, one for T54h and T96h) to collect 20  
116 to 100 mg of fertilization competent mycelium or/and perithecia. One to five Petri dishes *per*  
117 time course experiment were thus inoculated simultaneously with a growing *S mat*<sup>-</sup> strain and  
118 spermatia were collected by washing with 1.5 ml of H<sub>2</sub>O six days later. The collected spermatia  
119 (around 10<sup>4</sup> spermatia per ml) were pooled and then used to fertilize a corresponding number  
120 of Petri dishes of 4-day-old *S mat*<sup>+</sup> strain. The moment just prior the fertilization process was  
121 considered as the T0 (0 hours). The biological material was then collected 6, 12, 18, 24, 30, 42,  
122 54 and 96 hours after fertilization and an additional time point 24 hours prior fertilization was  
123 taken. Three biological replicates issued from three independent time course have been  
124 collected for all time points and two for T42. The RNAs pool used as a reference was assembled  
125 by combining equal amount of RNAs from various growth and crosses conditions and has  
126 already been described (Bidard *et al.*, 2010).

127 The biological material collected by scraping gently either cellophane or cheesecloth  
128 with a cover glass, was dried out with filter paper, weighted, frozen in liquid nitrogen and stored  
129 at -80°C. Total RNAs of *P. anserina* were extracted using RNeasy Plant Mini Kit (Qiagen),  
130 including, a grinding process using a Mikro-Dismembrator (Sartorius) and a DNase treatment.  
131 The quality and quantity of the total RNAs was determined by using a Nanodrop  
132 spectrophotometer (Nanodrop Technologies) and the Bioanalyzer 2100 system (Agilent  
133 Technologies) as previously described (Imbeaud *et al.*, 2005 (Bidard *et al.* 2010; Bidard *et al.*  
134 2011)).

### 135 **Gene expression array-based hybridization**

136 Transcriptome microarray experimental procedure, *i.e.* targets preparation,  
137 hybridization and washing, was done following the two-color microarray-based gene  
138 expression analysis instructions (version 5.0, February 2007) as described by the manufacturer  
139 (Agilent Technologies). One- $\mu$ g aliquots of total RNA were labelled using the Low RNA input  
140 fluorescent linear amplification (LRILAK) PLUS kit (Agilent Technologies); internal standards  
141 came from the Two-color RNA spike kit (Agilent technologies). The labelling efficiency and

142 the product integrity were checked as described by (Graudens et al., 2006). The reference cRNA  
143 was labelled with Cy5 and sample probes with Cy3. Then, Cy3- and Cy5-labeled targets were  
144 mixed and incubated on an Agilent microarray slide for 17 hours at 65°C, in a rotating oven (6  
145 rpm for a 1x44K and 10 rpm for a 4x44K array format), using an Agilent *in situ* hybridization  
146 kit. The slides were washed and then any traces of water were removed by centrifugation at 800  
147 rpm for 1 min (Bidard et al. 2010; Bidard et al. 2011).

#### 148 **Data acquisition and analysis**

149 Microarrays were scanned using an Agilent dual laser DNA microarray scanner (Agilent  
150 Technologies), model G2567AA, with 5- $\mu$ m resolution. The resulting 16-bit image files were  
151 analyzed in the *Feature Extraction* processing system (FE, v9.5.3). Spot and background  
152 intensities were extracted and normalized with the Feature Extraction software using the GE2-  
153 v4\_95\_Feb07 default protocol (Local background subtracted and LOWESS normalization).  
154 Preliminary array quality was assessed using Agilent control features as well as spike-in  
155 controls (Agilent 2-Color Spike-in Kit for RNA experiment). Subsequent flagging was done  
156 according to the GenePix Pro software (Molecular Devices Sunnyvale, CA, USA)  
157 nomenclature, including four levels of flags (good (100), bad (-100), not found (-50), moderate  
158 (0)) and raw data from the FE software ((Bidard et al. 2010; Bidard et al. 2011)).

159 The statistical analysis was conducted using the R environment and the MAANOVA  
160 (MicroArray ANalysis Of Variance) package (version 1.14.0) (Wu et al., 2003). It allows fitting  
161 an ANOVA model as proposed by Kerr et al (2001) and provides a way to consider multiple  
162 sources of variation in the microarray experiment. Technical biases were then corrected by  
163 including specific terms in the ANOVA model.

164 The differential analysis was performed on the whole of the replicated probes by excluding  
165 control spots. A F-test based on the James-Stein approach using a weighted combination of  
166 global and gene-specific variance estimates was used (Cui et al., 2003). P-values were  
167 computed by permutation analysis.

168 All p-values obtained were adjusted by the false discovery rate step-up method of  
169 Benjamini and Hochberg to account for multiple testing (Benjamini and Hochberg, 1995).  
170 Differentially expressed genes were defined as those whose adjusted p-value was statistically  
171 significant at a level of  $p < 0.05$ , with a fold change  $\geq 2$  or  $\leq -2$  and a  $A_{mean} \geq 6$  or a fold change  
172  $\geq 4$  or  $\leq -4$  and a  $A_{mean} \geq 4.5$

#### 173 **Classification**

174 Genes significantly differentially expressed in at least one of the comparisons and presenting a  
175 maximum value minus a minimal value  $>2.5$  after a median center transformation were  
176 classified (1798 genes). To this list, the mating-type transcription factor SMR1, SMR2 and  
177 FMR1 were added. Clusters analysis was computed from expression intensity measurements  
178 with the correlation coefficient as similarity measure. A hierarchical clustering (UPGMA  
179 method) and a K-means clustering to 15 classes were computed independently using Spotfire  
180 DecisionSite software package (Spotfire, Somerville, Mass.) with correlation coefficient as  
181 similarity measure on the 1798 regulated genes. Intersection of clusters obtained by the two  
182 clustering method results in 13 robust clusters with 1197 CDS classified. All microarray data  
183 is MIAME compliant. The raw data has been deposited in the MIAME compliant Gene  
184 Expression Omnibus database (Edgar et al. 2002) and is accessible through the GEO Series  
185 accession number GSE21659  
186 <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE93094>.

### 187 **Ortholog analysis**

188 Orthologous genes were searched between *P. anserina* and the other species *N. crassa*,  
189 *S. macrospora*, *C. globosom* and *T. reesei*. For this purpose, protein sequences were  
190 downloaded from Uniprot (<https://www.uniprot.org/>) under the accession numbers:  
191 UP000001805\_367110 (*N. crassa*), UP000001881\_771870 (*S. macrospora*),  
192 UP000001056\_306901 (*C. globosum*) and UP000008984\_431241 (*T. reesei*). The OrthoFinder  
193 algorithm (Emms et Kelly 2019) was used with default parameters to obtain “orthogroups”  
194 (Table S3), *i.e.* groups of genes that are good candidates to share a common ancestor between  
195 the compared species.

### 196 **Transcription factors network**


197 To search among all the transcriptional regulators for those that could be involved in  
198 setting up the observed transcriptional waves, we listed the candidate genes that encode  
199 transcription factors (TFs). This list includes 92 genes (table S7). Gene expression profiles for  
200 each of the TF genes were then retrieved from the microarray datasets and used to search for  
201 significant correlations with the expression profiles of all genes previously classified in  
202 transcriptional waves I, II, III, IV or V. We then considered the situations, in which the  
203 calculated correlation between the expression profiles of a TF gene and a gene classified in a  
204 transcriptional wave was above a predefined threshold (here 0.95). Our assumption is that such  
205 a high correlation value may reveal a potential link of transcriptional control, between the TF  
206 and another gene (referred to as “target gene”).



## 207 Results

### 208 High-resolution kinetics of transcription fluctuations during sexual reproduction

209 To ensure an accurate, high-density representation of transcriptional variation, we  
 210 collected RNA samples at ten key time points during the sexual reproductive cycle of *P.*  
 211 *anserina* (Fig. 1).




212  
 213 **Figure 1 - *P. anserina* developmental time course from fertilization to ascospore**  
 214 **maturation.** A. Schematic representation of the fertilization and the main steps of  
 215 zygotic development. A. Pictures of the main steps of sexual development. Sexual  
 216 reproduction began with the production of male (spermatium) and female  
 217 (ascogonium) gametes. A pheromone/receptor signaling system allows the female  
 218 gametes to attract specialized hyphae (trichogynes) that recognize and fuse with male  
 219 gametes of compatible mating type. This event initiates the multi-step process of  
 220 sexual development that begins with the formation of fructifications and ends with  
 221 the discharge of ascospores. The nucleus of the male gamete enters the trichogyne,

222 migrates along this specialized hypha, reaches the ascogonium and undergoes  
223 several mitotic divisions. Fertilization (T0) triggers the development of  
224 protoperithecia (T6 to T24), which recruit multiple layers of protective maternal  
225 hyphae to grow into perithecia (T24 to T45). The developing fructifications shelter  
226 the fertilized ascogonial cells (T24 to T30), which contain multiple haploid parental  
227 nuclei of either *mat+* or *mat-* genotype. Differentiation of paraphyses begins (T24)  
228 just before three-celled hook-shaped structures called croziers emerge from the  
229 ascogonial cells (T30 to T42). Croziers are dikaryotic because they contain two  
230 haploid nuclei derived from both parents. Karyogamy (T30 to T44) occurs in the  
231 upper cells of croziers. The resulting diploid zygote immediately undergoes meiosis  
232 (T30 to T44), producing four haploid nuclei, which then undergo mitosis prior to  
233 ascospore delineation (T54 to T96). Ascospores are formed around two non-sister  
234 nuclei within the developing asci. In rare cases, two ascospores form around a single  
235 haploid nucleus, resulting in five-spored asci. C. Time line of the sexual cycle of *P.*  
236 *anserina* with the steps from which RNAs have been extracted on the left.

237  
238 *P. anserina* is a pseudohomothallic fungus, namely natural isolates are self-fertile  
239 because they contain nuclei of each mating-type. Self-fertilization does not start at a specific  
240 time point and therefore natural isolates cannot be used for a time course analysis of sexual  
241 development. However, the genetic basis of the *P. anserina* mating system is heterothallism  
242 with two mating-type idiomorphs called *mat+* and *mat-* (Turgeon et Debuchy 2007). Typical  
243 ascospores contain *mat+* and *mat-* nuclei, but a few asci contain monokaryotic ascospores  
244 which produce self-sterile individuals. Those latter individuals were used in the experiments  
245 presented here. Fertilization was performed by spreading *mat-* male gametes on a *mat+* strain.  
246 This experimental design allows the time course to start at a well determined time point for  
247 synchronization purpose. Under this experimental condition almost all fertilization events  
248 occurred within the first six hours (Fig. S1A and S1B). However, despite this experimental  
249 design, RNA samples were extracted from biologically non-homogeneous tissues because *P.*  
250 *anserina* fructifications are composed of different cell types (Fig. 1A and 1B).

251 We measured the expression of 10,507 nuclear genes at ten time points which are  
252 designated as Tm24 for mycelium 24 h before fertilization and T0, T6, T12, T18, T24, T30,  
253 T42, T54 and T96 for time 0 to 96 h postfertilization, respectively (Fig. 1C, Table S1). We  
254 performed a Principal Component Analysis (PCA) on the complete dataset. More than 75% of  
255 the total variance was captured by only three principal components (PCs, Fig. 2A and 2B, Table  
256 S2).



257  
258  
259  
260  
261  
262  
263  
264

**Figure 2 - PCA analysis reveals a strong temporal effect on gene expression variations relevant to the key developmental steps in *P. anserina***

Principal Component Analysis (PCA) on the complete dataset (*i.e.* all genes and all time points). **A.** Axis 1 and 2 of the PCA explain 38% + 24 % = 62% of the variance. **B.** Axis 2 and 3 of the PCA explain 24% + 16 % = 40 % of the variance. Altogether, these first three axes of the PCA explain 78 % of the variance. **C.** Gene expression profiles for the 50 genes, which contribute the most to the definition of the first


265 (PCA1), second (PCA2) and third (PCA3) principal components, as shown in A and  
266 B. Detailed lists of these genes are given in Table S2. They were separated according  
267 to the type of regulation they experience during sexual development (positive or  
268 negative). Tm24, T0, T6, T12, T18, T24, T30, T42, T54 and T96 : time points of the  
269 kinetic were the biological samples were collected, described in Fig. 1.

270

271 With the exception of Tm24, we observed that all samples were distributed along the PC1 axis  
272 (38%) according to the sexual development time line (Fig. 2A). On PC2 axis, we found T30  
273 and T42 samples clustered apart from the others (24%). Interestingly, this period corresponds  
274 to the formation of the dikaryotic cells, which are the progenitors of fungal zygotic lineage (Fig.  
275 1). The peripheral position of the Tm24 sample (Fig. 2B) which appeared on the PC3 axis  
276 (16%), indicated that the gene expression profiles derived from vegetative mycelium (Tm24)  
277 were clearly different from those of all samples collected after fertilization. Overall, this  
278 unsupervised exploratory data analysis validated both the relevance of our experimental design  
279 and the robustness of our differential analysis (see below). The expression profiles for the 50  
280 genes whose expression measurements contributed the most to the calculation of the principal  
281 components 1, 2 and 3 are shown Figure 2C (and detailed Table S2). The data indicated that  
282 the greatest fold changes, both positive and negative, occurred after the formation of dikaryotic  
283 cells (>T30). This was also the case for axis 2, but only after the formation of the ascospores  
284 (>T54). We observed that the positive fold change values on axis 1 exhibited a higher dispersion  
285 after T42.

### 286 **Topology of global expression variation reveals five major transcriptional waves during** 287 **sexual development**

288 Using k-means clustering, we defined 13 representative expression patterns (Fig. S2),  
289 encompassing 1,186 genes. Further data outlined five successive transcriptional waves (Fig. 3  
290 and Table S1). Notably, they all shared a common pivot point (Fig. S3), corresponding to the  
291 mid-course of the kinetic (T30-T42). This observation indicated a pause in the regulation of  
292 transcriptional activity at the time of dikaryon formation. The 1,186 genes were not evenly  
293 distributed across the five waves (Fig. 3 and Table S1), resulting in two periods of intense and  
294 equal transcriptional fluctuation (waves II and IV) interspersed with three periods of moderate  
295 transcriptional fluctuation (waves I, III, and V).



296

297

298

### Figure 3 - Temporal transcriptional waves over the course of sexual development

299

300

301

302

303

304

305

306

A. Expression profiles of the five major successive transcriptional waves (numbered in Roman figures) that take place during *P. anserina* sexual reproduction (Bottom timeline, see panel B for zoom). Number of DE genes present in each wave is indicated below the corresponding graph. Waves II (early up/late down pattern, 31.51%) and IV (early down/late up pattern, 36.37%) accounted for 67.87% of them (33.98% and 37.42%, respectively), while wave I (early down pattern, 7.41%), wave III (early-mid up pattern, 9.62%) and wave V (late up pattern, 15.02%) contained only 32.13% of them, respectively.

307

308

B. Major developmental events according to the sexual reproduction timeline described in Figure 1.

309

310

311

312

313

314

The topology of the transcriptional waves therefore suggests that the genes of waves II and III are down-regulated after a peak of expression during protoperithecium development, while the genes of waves IV and V are up-regulated after perithecium development. These patterns are in accordance with the results obtained in *Neurospora crassa*, during fruiting body development (Wang et al. 2014).

315

316

### Evolutionary conservation and functional annotation of differentially expressed genes during sexual reproduction



317

318

319

Across the entire kinetic, we found 3,466 genes that were differentially expressed (DE, see Methods). They represent 33% of the total gene set for which gene expression measurements were monitored (Table S1, Fig. 4A), including 1,133 of the genes clustered in

320 the five transcriptional waves (more than 95% of the clustered genes, Table S1 and Fig. 3).  
 321 Overall, this analysis revealed that in *P. anserina*, one-third of the complete gene set is  
 322 transcriptionally regulated during sexual reproduction. Of this, one-third follows one of the five  
 323 transcriptional waves disclosed by the clustering analysis. To test for evolutive conservation of  
 324 these DE genes, we built group of orthologs, using four species of filamentous fungi, *i.e.* the  
 325 homothallic species *Chaetomium globosum* and *Sordaria macrospora* and the heterothallic  
 326 species *N. crassa* and *Trichoderma reesei*. A total of 2,957 DE genes (more than 85%) were  
 327 identified as belonging to an ortholog group shared with at least one of the four species (Table  
 328 S3). A core set of 1,496 DE genes is conserved in all four fungi (Fig. 4, Table S3).



329


330 **Figure 4 - Distribution of differentially expressed genes along the kinetics of**  
331 **sexual reproduction and their evolutive conservation**

332 **A.** Heatmap of pairwise comparisons of DE genes across the complete kinetics. For  
333 each pairwise comparison, the number of DE genes is expressed as a percentage  
334 relative to the total number of DE genes (3,466) and also as a color intensity given  
335 by the color key bar (right).

336 **B.** Venn diagram showing the conserved orthologs between the indicated fungal  
337 species. Cg: *C. globosum* (homothallic), Sm: *S. macrospora* (homothallic), Nc: *N.*  
338 *crassa* (heterothallic) and Tr: *T. reesei* (heterothallic).

339  
340 A pairwise comparison of the DE gene set of *P. anserina* with each gene set of the four  
341 fungi, showed that the number of conserved DE genes is consistent with fungal evolutionary  
342 history (*i.e.* closely related fungi share more DE genes than distantly related fungi). The  
343 remaining 506 genes (14.78%) were classified as orphan genes, with 247 (21.84%) belonging  
344 to the five transcriptional waves. Notably, the set of DE genes, including that of the five waves,  
345 were found to be enriched with orphan genes (p-values:  $2.55 \times 10^{-9}$  and  $4.60 \times 10^{-20}$  respectively).

346 We then search for functional annotation of the proteins encoded by the DE genes. We  
347 found that 1,279 (36.95%) of the 3,466 putative proteins encoded by the DE genes lacked Pfam  
348 annotation (Table S1, (Mistry et al. 2021)), which is significantly more (p-value:  $8.55 \times 10^{-9}$ )  
349 than in the complete list of *P. anserina*'s proteins. The lack of annotated conserved domains  
350 was further enriched (p-value:  $1.48 \times 10^{-13}$ ) in wave-embedded DE genes, as 515 out of these  
351 1,133 putative proteins (45.44%) lacked Pfam annotation. We also performed Gene Ontology  
352 (GO, (Ashburner et al. 2000; The Gene Ontology Consortium et al. 2023)) analysis on the DE  
353 genes (Table S4, Fig. 5A).



354

355

**Figure 5 - GO term and expert annotation categories enriched in the transcriptional waves.**

356

**A.** Descriptions of GO terms significantly enriched ( $p\text{-value} \leq 0.05$ ) in at least one

358 wave of the transcriptional waves. **B.** Enrichments of genes of interest in at least one

359 wave of the transcriptional waves. TF: transcription factors, SM: secondary

360 metabolites, HET: fungal specific vegetative incompatibility HET domain. Points

361 are scaled and colored according to the number of genes in each category for each

362 wave. No score is shown for zero genes.

363

364 In general, the enriched GO terms and Pfam domains (Table S4 and S5) described

365 intense catalytic (GO:0003824) and metabolic (GO:0008152) activities, associated with

366 binding (GO:0005488, PF00187). Accordingly, we observed enrichments in genes involved in



367 redox catalytic activities (GO:0003824, PF00067, PF03055, PF01565, PF00106, PF01494,  
368 PF00199, PF08240) and transporter activities (GO:0005215, Pfam PF07690.19). We also  
369 uncovered enrichments in DE genes involved in cellular differentiation programs more  
370 specifically related to sexual reproduction. They either deal with cellular anatomical entity  
371 (GO:0110165), localization (GO:0051179) or encode zinc-binding domain transcription factors  
372 (PF13695), necrosis inducing factor (PF14856) and TUDOR domain containing protein  
373 (PF11160). However, if the repertoire of GO terms (Table S4) and Pfam domains (Table S1  
374 and S5) generated by this study should provide some clues to select candidate genes for *in vivo*  
375 functional characterization, our analysis essentially showed that more than one third of the DE  
376 genes have no associated function yet.

### 377 **Searching for gene of interest, the expertise-driven approach**

378 We then performed a biological expertise-driven approach to mine the functions of the  
379 full set of DE genes (Table S6, Fig. 5B).

### 380 **Developmental processes at work**

381 Fertilization of *P. anserina* requires exposure to blue light, but the molecular basis of  
382 this signal transduction is still unknown. In *N. crassa*, the blue light signal is transduced by the  
383 photoreceptors WC-1 and WC-2 (Froehlich et al. 2002; He et al. 2002). In this study, we found  
384 five blue light responsive genes that were differentially expressed (Table S6A, Fig. S4A-B).  
385 Consistent with a role in fertilization three of them (including *Pa\_5\_6130/WC-1*) were  
386 expressed before meiosis, while the other two were expressed after meiosis, suggesting that  
387 blue light may also play a role during ascospore formation.

388 Orthologs of genes involved in plasmogamy in yeast models (*Pa\_3\_11320/Prm1* (Curto  
389 et al. 2014), *Pa\_3\_4770/CDC3* (Kurahashi, Imai, et Yamamoto 2002), *Pa\_5\_9510/Fig1*  
390 (Erdman et al. 1998), *Pa\_1\_1930/KEX2* (Julius et al. 1984), *Pa\_4\_6420/KAR2* (Rose, Misra,  
391 et Vogel 1989), *Pa\_7\_10810/SEC63* (Young et al. 2001) were found to be up-regulated until  
392 T42 and then downregulated (Table S56, Fig. S4C). Development of the zygotic lineage likely  
393 involves compartmentalization through active cell wall remodeling, as we found several DE  
394 genes encoding different types of cell wall proteins and growth factor receptors (Table S5). One  
395 of them was the expansin-like EEL2, involved in cell wall expansion, whose gene *Pa\_2\_310*  
396 was downregulated from karyogamy to ascospore maturation (T30-T96). This expression  
397 pattern was similar to that of the *C. globosum* homolog (*CHGG\_00523*) but divergent from that  
398 of the *N. crassa* homolog (*NCU04603*) (Hutchison et Glass 2010; Wang et al. 2019). Cell-cell  
399 interactions may also be tightly regulated, as exemplified by the strict downregulation (T0-T96)  
400 of *Pa\_1\_17480*, which encodes the adhesin protein MAD1 (Wang et St Leger 2007).

401 Since sexual development is rich in cell divisions, we searched for DE cell cycle related  
402 genes and found this category slightly enriched (p-value =  $2.9 \times 10^{-2}$ , Table S5B). In addition to  
403 the TOR pathway inhibitor *Pa\_3\_3700/FgFkbp12*, it included homologs of genes required for  
404 proper fructification formation (*Pa\_1\_17170/KIN3*, *Pa\_1\_23300/cel-2*, *Pa\_2\_6460/Pah1*,  
405 *Pa\_2\_890/asm-1*), axial budding (*Pa\_2\_290/axl2*) and genes encoding G-coupled receptors  
406 (*Pa\_4\_5350*, *Pa\_1\_590*) and ras-like protein (*Pa\_1\_19030/krev-1*). The expression profile of  
407 the *Pa\_2\_290/axl2* gene, encoding an integral plasma membrane protein, was of particular  
408 interest: it was down-regulated when the two haploid parental nuclei actively divide into  
409 multinucleated syncytial ascogonial cells (Fig. S4D).

410 As expected, we also found DE genes encoding cytoskeletal elements (Fig. S4E).  
411 Interestingly, *Pa\_5\_5720*, encoding a pyruvate decarboxylase (PDC) was found strongly up-  
412 regulated during ascospore maturation (wave V). This observation further enforces the  
413 hypothesis that pyruvate decarboxylase is involved as a structural protein (PDC-filaments)  
414 associated with the cytoskeleton during the sexual development of filamentous fungi  
415 (Thompson-Coffe et al. 1999). Conversely, *Pa\_4\_7260*, encoding the nucleoprotein Cik1,  
416 required for both karyogamy and mitotic spindle organization (Cottingham et al. 1999), was  
417 found down-regulated at the same time. Some of the cytoskeleton DE genes showed co-  
418 expression with the plasmogamy DE genes (Fig. S4F), making them good candidates to be  
419 involved in fusion morphogenesis processes.

420 Three genes essential for ascus development were also found DE (Fig. S4G, Table S5A):  
421 *P. anserina*'s orthologs of *Pa\_4\_9730/asd-1* (wave IV (Nelson, Merino, et Metzberg 1997)),  
422 *Pa\_4\_9750/asd-3* (Galagan et al. 2003) and *Pa\_2\_13270/con-6* (wave V (White et Yanofsky  
423 1993)). *P. anserina* does not form clonal propagules. However, the orthologs of genes required  
424 for conidiation in *N. crassa* and *Aspergillus nidulans* such as *Pa\_2\_13270/con-6*,  
425 *Pa\_4\_9730/asd-1*, *Pa\_4\_9750/asd-3*, *Pa\_1\_12720/Pyg1*, *Pa\_1\_30/TmpA* and *Pa\_0\_1540*  
426 were found to be up-regulated after meiosis, suggesting that genes primarily involved in  
427 conidiation, *i.e.* asexual reproduction, may have been recruited to participate in ascospore  
428 formation. On the contrary, early conidial development-2 *Pa\_7\_780/ecd-2* was down regulated  
429 during the first half of the kinetic (Sun et al. 2012).

### 430 **Producing energy**

431 The reproductive process is energy consuming and therefore the mobilization of carbon  
432 resources is one of the keys to its success. The genome of *P. anserina* contains 276 genes  
433 encoding CAZymes ((Espagne et al. 2008) and Table S5D). Among them, 141 were found DE  
434 during sexual development, showing an enrichment of this metabolic activity (p-value =

435  $4.36 \times 10^{-5}$ ). The expression profile of 42 CAZyme genes corresponded to one of the five  
436 transcriptional waves, especially waves II and IV. The remaining 99 showed contrasting  
437 expression patterns (Fig. S4 H-T), but up-regulation was enriched over down-regulation, as 62  
438 of them were up-regulated during the kinetic ( $\text{Chi}^2$ ,  $p\text{-val} < 0.001$ ). We also found that several  
439 CAZymes genes had some of the highest positive fold changes measured in this study (Table  
440 S1). In addition, three glycolysis enzymes genes (6-phosphofructo-2-kinase *Pa\_3\_6610*,  
441 fructose-bisphosphate aldolase *Pa\_0\_710*, and phosphoglycerate kinase *Pa\_6\_10610*) were up-  
442 regulated with a similar pattern of overexpression when asci begin to differentiate (Fig. S4U,  
443 Table S5E).

444 In *P. anserina*, long-chain fatty acids are shortened in peroxisomes then degraded in  
445 mitochondria. The *Pa\_6\_3500/PaPex7* gene, which encodes the peroxisomal-targeting-signal-  
446 2 receptor, is found to be up-regulated during ascospore maturation (T54-T96, Table S1 and  
447 Fig. S4V). This is in line with the defects in nuclear positioning of the post-meiotic asci  
448 observed in *pex7* knock-out mutants, which can result in either abnormal ascospore formation  
449 or to ascospores containing an abnormal number of nuclei (Bonnet et al. 2006). The  
450 *Pa\_1\_23790/PaFox2* gene (Table S1 and Fig. S4V), which encodes the peroxisomal  
451 multifunctional protein 2, a key enzyme in the fatty acid  $\beta$ -oxidation pathway, was found to be  
452 down-regulated during the early stages of sexual development (T12-T54) and up-regulated  
453 during ascospore maturation (T54-T96). This expression profile may explain the reduced  
454 acetyl-CoA availability within the ascospores that results in the pigmentation defect observed  
455 in *PaFox2* knock-out mutants (Boisnard et al. 2009).

#### 456 **Secondary metabolites: major actors of sexual development?**

457 Prokaryotic and eukaryotic micro-organisms produce a wide variety of secondary  
458 metabolites (SMs). These molecules are not intended to perform housekeeping functions, but  
459 rather are used as adaptive tools to cope with environmental fluctuations or to compete with  
460 other living organisms. The *P. anserina* genome contains 35 biosynthetic gene clusters (BGCs)  
461 ((Lamacchia et al. 2016), Table S5F, Fig. S4W), containing 474 genes. SM encoding genes  
462 were enriched among the complete set of DE genes ( $p\text{-value} = 8.31 \times 10^{-10}$ ) as 258 of them were  
463 found to be DE (33 out of 35 BGCs contained at least two DE genes). These genes were also  
464 significantly up-regulated ( $\text{Chi}^2$ ,  $p\text{-val} < 0.001$ ). BGCs N°1, 11, 18, 19, 24, 27 and 34 showed  
465 an expression profile topology (Fig. S4W) similar to the general pattern described in *N. crassa*  
466 (Wang et al. 2022), which was a coordinated downregulation from fertilization to meiosis  
467 followed by a coordinated upregulation during ascospore formation and maturation.

468 We also examined whether co-expression was shared by multiple SM genes within the  
469 34 BGCs. This was the case for the sterigmatocystin BGC, where 17 of the 23 genes were  
470 differentially expressed during sexual reproduction (Table S5F, cluster 32, Fig. S4X, (Slot et  
471 Rokas 2011)), including the transcription factor *PaAflr* (Shen et al. 2019). In addition, we  
472 identified four positive regulators of the sterigmatocystin BGC (Table S5F, Fig. S4Y,  
473 *Pa\_2\_7220*, *Pa\_2\_7240*, *Pa\_2\_7250*, *Pa\_2\_7260* (Bhetariya et al. 2011)), that showed the  
474 same expression pattern across sexual reproduction. Similarly, 13 of the 20 genes in the  
475 terrequinone BGC were differentially expressed (Fig. S4Z).

476 In addition, we identified several isolated genes from different BGCs that showed  
477 similar peak expression to each other but no co-expression with other genes from their  
478 respective BCG (Fig. S4AA-AB). These expression peaks of genes encoding mostly proteins  
479 of unknown function corresponded to two key moments in early development (T6 and T12),  
480 just before and during dikaryon formation. Taken together, our analyses further support the key  
481 role of SM in sexual reproduction and highlight the role of previously uncharacterized clusters  
482 and isolated SM genes.

### 483 **From regulation of biotic interactions to that of sexual development**

484 Genes encoding STAND proteins (Signal-transducing ATPases with Numerous  
485 Domains, (Leipe, Koonin, et Aravind 2004) have been described as involved in fungal biotic  
486 interactions with living organisms, either from the same species or from different species but  
487 mostly during vegetative growth. The *P. anserina* genome contains 34 genes encoding STAND  
488 proteins with a NACHT domain (PF05729), eight of which were found to be down-regulated  
489 either throughout the sexual cycle kinetics or prior to perithecium development (Table S5G,  
490 Fig. S4AC). This group includes *Pa\_3\_10930/nwd1* (Chevanne et al. 2010), *Pa\_7\_10370/sesB-*  
491 *like*, *Pa\_2\_7940/het-d* (Espagne et al. 2002) and *Pa\_1\_11380* (Bidard, Clavé, et Saupe 2013).  
492 Fungal interactions are also under the control of a programmed cell death called vegetative  
493 incompatibility that prevents hyphal fusion of two genetically distinct wild-type strains. Most  
494 of the effectors of this incompatibility reaction (IR) contain HET domains (PF06985).  
495 Strikingly, 59 of the 128 *het* genes annotated in the *P. anserina*'s genome were found to be DE,  
496 including *Pa\_4\_1190/hnwd1* and *Pa\_7\_3460/Pin-C* (Chevanne et al. 2010; Kaneko et al. 2006;  
497 Bidard, Clavé, et Saupe 2013). The majority of the DE *het* genes were down-regulated either  
498 throughout the complete sexual development or from dikaryon formation (Fig. S4AD), but  
499 seven of them were up-regulated during some steps of the kinetic, mostly after karyogamy (Fig.  
500 S4AE). In addition, three *het* genes showed a peculiar behavior (Fig. S4AF): *Pa\_5\_2540*  
501 expression oscillated during sexual reproduction, *Pa\_3\_5550* expression peaked during

502 protoperithecium development and *Pa\_4\_1800* expression was down-regulated until ascospore  
503 maturation and over-expressed during this process.

504 In the DE gene set, we also found gene families encoding toxins that can induced host  
505 cell death or effector-like proteins. All of the six genes encoding the NEP1-like proteins pore-  
506 forming toxin NPP1 (necrosis-inducing *Phytophthora* protein, (Ottmann et al. 2009) were  
507 down-regulated at least during the second half of the sexual development, i.e. before karyogamy  
508 (Fig. S4AG). Six out of the eight genes encoding Hce2 pathogen effectors (Stergiopoulos et al.  
509 2013) and one gene annotated as Ptu bug toxin encoding gene (*Pa\_6\_9450*) were also DE. In  
510 contrast to the *NPP1* genes, all of them showed up-regulation during the course of sexual  
511 development, five of them belonging to the wave IV pattern of expression (Fig S4AH).

### 512 **Epigenetic effectors at work in sexual reproduction**

513 In recent years, epigenetic effectors have been identified as key players in fructification  
514 morphogenesis (Carlier et al. 2021; Nowrousian 2022). Therefore, we investigated the  
515 expression patterns of genes encoding proteins that can either methylate cytosines, write or  
516 erase histone post-translational modifications or interfere with messenger RNA synthesis and  
517 stability.

518 *PaRid* and *PaDnmt5*, two of the three DNA methyltransferase encoding genes present  
519 in the *P. anserina* genome, were found to be DE during this kinetic. *PaRid* expression, which  
520 is essential for dikaryon formation (Grognet et al. 2019) peaked at T12, while *PaDnmt5*  
521 expression was increasingly up-regulated from T12 to T42 (Fig. S4AI). *Pa\_5\_2900*, which  
522 encodes a RING-type E3 ubiquitin transferase URF1-like, was found to be strongly up-  
523 regulated from T42. In mammals, this protein binds both hemi-methylated DNA and tri-  
524 methylated lysine 9 of histone H3 (H3K9me3), thereby bridging DNA methylation and  
525 chromatin modification (Foster et al. 2018).

526 *Pa\_7\_10020/KAT1* and *Pa\_1\_7460/Hos-2* encodes two chromatin modifiers with  
527 antagonistic enzymatic activities, respectively a histone acetyltransferase encoding gene (Tong  
528 et al. 2012) and a histone deacetylase (Wang, Kurdistani, et Grunstein 2002) respectively.  
529 However, these two genes showed co-expression over the entire kinetic, i.e. they were up-  
530 regulated from fertilization to karyogamy and then down-regulated (Fig. S4AJ). This pattern  
531 was in contrast to that of *Pa\_7\_5890* encoding the Sin3-associated polypeptide SAP18, which  
532 enhances transcriptional repression through histone deacetylation (Cheng et Bishop 2002). Two  
533 genes encoding sirtuins NAD(+)-dependent histone deacetylases were also found to be DE:  
534 *Pa\_4\_8570/PaSir2* showed a wave III expression pattern that resembled that of  
535 *Pa\_1\_19440/PaRid* (Fig. S4AI), while the gene encoding *Pa\_4\_7390/SIRT5* was down-

536 regulated throughout sexual reproduction. Of the 32 genes encoding SET domain proteins,  
537 seven were found to be DE (Fig. S4AK), although none has yet been shown to have histone  
538 lysine methyltransferase activity. Four genes encoding putative histone demethylase containing  
539 JmJC domains were also found DE (Fig. S4AL).

540 Promoter accessibility also depends on nucleosome positioning. The ATP-dependent  
541 chromatin remodeling protein SWI/SNF *Pa\_5\_4310/SMARCA3* (Oh et al. 2013) was one of the  
542 most highly up-regulated wave III genes (Fig. S4AM). Its transcription pattern is similar to that  
543 of *Pa\_1\_19440/PaRid* (see Fig. S4AI) and *Pa\_4\_8570/PaSir2* (Fig. S4AJ). Transcription  
544 requires a large number of proteins assembled into conserved complexes. Two genes  
545 (*Pa\_1\_17860* and *Pa\_6\_1460*) encoding components of the SAGA complex were found to be  
546 up-regulated from T6 to T54 with very similar expression patterns (Fig. S4 AM). However, we  
547 also found similarly co-regulated genes although encoding factors with antagonistic effects on  
548 transcription, such as the SWD2 subunit of the COMPASS complex (*Pa\_7\_11070*) and the  
549 extra sexcombs subunit of the Polycomb Repressive Complex 2 (*Pa\_3\_4080*).

550 There is no experimental evidence that post-transcriptional RNAi mechanisms Quelling  
551 (Romano et Macino 1992) and MSUD (Shiu et al. 2001) are functional in *P. anserina*. However,  
552 genes encoding essential components of Quelling (Fig. S4AN), the RNA-dependant RNA  
553 polymerase (RdRP) *Pa\_7\_4790/Qde1* (Cogoni et Macino 1999) and the dicer *Pa\_6\_6150/Dcl-1*  
554 (Yang, Ye, et Liu 2015), were downregulated from T18 and T30, respectively. On the  
555 contrary, expression of *Pa\_7\_9210* encoding the Qde-2 interacting exonuclease QIP was up-  
556 regulated from T6 to T30. *N. crassa*  $\Delta qde-1$  quelling mutant strains produced normal-sized  
557 perithecia but no ascospore (Wang et al. 2014). Four genes involved in MSUD were also found  
558 to be DE (Fig. S4AO). *Pa\_2\_9830/Sad-2* and *Pa\_5\_11380/Sad-5* were up-regulated whereas  
559 *Pa\_1\_19900/Sad-1*, encoding the RdRP and *Pa\_1\_16040/Sad-3*, encoding the helicase, were  
560 down-regulated. Interestingly, *Pa\_6\_11690/rrp-3*, which encodes an RdRP that could not be  
561 clearly assigned to Quelling or MSUD pathways in *N. crassa* (Borkovich et al. 2004), was co-  
562 expressed with the Quelling genes *Pa\_1\_19900/Sad-1* and *Pa\_1\_16040/Sad-3* in *P. anserina*.  
563 Knockout of the *N. crassa Sad-1* and *Sad-3* resulted in arrest of ascus development in meiotic  
564 prophase and production of normal-sized perithecia and young asci without ascospores,  
565 respectively (Wang et al. 2014). *Sad2* is essential to *N. crassa* fructification development (Wang  
566 et al. 2014). These results suggest that RNAi genes may be silenced during meiosis and  
567 sporulation, which may explain the apparent lack of MSUD in *P. anserina* but also that  
568 *Pa\_2\_9830/Sad-2* and *Pa\_5\_11380/Sad-5* might have been recruited to play a role during  
569 ascospore dormancy.

570 Repeated sequences present in both haploid parental genomes of *P. anserina*'s  
571 dikaryotic cells are subjected to cytosine-to-thymine mutations by repeat-induced point  
572 mutation (RIP, (Bouhouche et al. 2004)). Modified 5-methylcytosines are known to be hotspots  
573 for cytosine-to-thymine transitions, because, unlike unmethylated cytosines, spontaneous  
574 deamination of 5-methylcytosines cannot be repaired by uracil excision (Pearl 2000). To  
575 counteract this genetic threat, members of the AlkB alpha-ketoglutarate-dependent dioxygenase  
576 (PF13532) family oxidize 5-methylcytosines to restore undamaged bases (Bian et al. 2019).  
577 Two genes encoding AlkB-like enzymes are present in the *P. anserina* genome. The expression  
578 of *Pa\_1\_12280* was down-regulated throughout the kinetic, while *Pa\_1\_17260* was up-  
579 regulated during two segments of the kinetic: i) from Tm24 to T12, corresponding to the peak  
580 of expression of *PaRid* and ii) from T24 to T54, when *PaRid* is no longer expressed (Fig.  
581 S4AP). Notably, both genes showed a superimposable flat expression profile at the time of RIP.

582 In filamentous fungi, a conserved post-transcriptional A-to-I mRNA editing process  
583 recodes transcription units (Teichert et al. 2017; Blank-Landeshammer et al. 2019), mainly  
584 during fructification and ascospore morphogenesis. We identified seven DE genes encoding  
585 adenosine deaminase acting on tRNA (ADAT), a class of enzymes which are responsible for  
586 A-to-I mRNA editing. *Pa\_1\_21170*, a wave III gene, showed a peak of expression similar to  
587 that of *PaRid*, while *Pa\_3\_6600* and *Pa\_6\_7750* were up-regulated from perithecium  
588 development to ascospore discharge (Fig. S4AQ).

### 589 **Co-expression Network: In Search of Potential Transcription Factor Targets During** 590 **Sexual Reproduction**

591 We then focused on the 96 transcription factors (TFs) encoding genes that were found  
592 to be DE (Table S6I). The expression profile of each TF gene was used to search for significant  
593 correlations with the expression profiles of all genes previously classified as transcriptional  
594 waves I, II, III, IV or V. Using this strategy, we build a network of 2002 potential transcriptional  
595 links between 76 TFs and 683 target genes (Fig. 6 and Table S7A-C).





616 include the mating-type proteins FMR1 and SMR2, which are both required for the  
617 development of the perithecium after fertilization (Zickler et al. 1995; Arnaise et al. 2001).  
618 Physical interaction between FMR1 and SMR2 was demonstrated and proposed to be required  
619 for their post-fertilization functions (Arnaise et al. 1995; Turgeon et Debuchy 2007; Bidard et  
620 al. 2011). Our study shows that FMR1 and SMR2 have a similar expression profile throughout  
621 sexual reproduction (Fig. S4AR). Their co-expression networks revealed also an almost  
622 identical number of potential target genes (FMR1, n = 79 ; SMR2, n = 78) (Table S7). They  
623 share 62 of their 79 potential target genes (Table S7). These observations support the idea that  
624 they act together in identical regulatory pathways by forming a heterodimer (Arnaise, Debuchy,  
625 and Picard 1997). The third and last *mat*- mating-type gene, *SMR1*, does not belong to the  
626 HMG-box family. It encodes a HPG protein (Turgeon et Debuchy 2007), which has been  
627 proposed to provide a scaffold for physical interaction of mating-type proteins (Zheng et al.  
628 2013). This role is consistent with its presence in wave IV, along with *FMR1* and *SMR2* (Fig.  
629 S4AR). Critical genes for the development of ascogenous hyphae are likely to be found among  
630 target genes of FMR1, SMR2 and SMR1 (Fig. 6, Table S7A). The only non-mating-type HMG  
631 protein present in the network is Pa\_1\_11050/PaHMG4, related to Nhp6p and Hmo1p from *S.*  
632 *cerevisiae* (Ait Benkhali et al. 2013). Its deletion in *P. anserina* resulted in an increased number  
633 of male organs. The lack of effect on fructification development is consistent with the down-  
634 regulation of Pa\_1\_11050/PaHMG4 at T42-T96 (Fig. S4AS). Proper shaping of the developing  
635 fructifications requires the homeobox protein Pa\_2\_6460/Pah1 (Coppin et al. 2012). The  
636 expression of the corresponding gene was accordingly up-regulated from T0 to T42 (Fig.  
637 S4AS). In contrast, Pa\_3\_4650/Pah6 was down-regulated between T0 and T54 (Fig. S4AS),  
638 although no fructification defect was associated with its deletion.

639 The Zn(II)<sub>2</sub>Cys<sub>6</sub> class of regulators is one of the largest families found in fungi  
640 (MacPherson, Larochelle, et Turcotte 2006) and consequently they represented almost half of  
641 the TFs present in the network (Table S7A). Three of them are of particular interest and have  
642 been extensively studied because of their properties. Pa\_1\_10140/PaPro1 is a regulator of the  
643 expression of the mating-type and *IDC* genes, both required for female fertility (Gautier et al.  
644 2018). In *S. macrospora*, *Pro1* has been identified as a master regulator of fructification  
645 formation (Steffens et al. 2016; Teichert, Pöggeler, et Nowrousian 2020). Its temporal and  
646 spatial expression is consistent with the presence of Pa\_1\_10140/PaPro1 in wave III (Fig.  
647 S4AS). Pa\_7\_7000 was characterized in wave IV as a coregulator of FMR1, SMR2 and SMR1  
648 target genes (Fig. 6). Pa\_7\_7000 is revealed as one of the most prominent nodes in the TF  
649 network (Fig. 6 and Fig. S4AT), but has not yet been characterized in *P. anserina*. Pa\_7\_7000

650 has orthologs in the seven species of the *Podospora* complex (Boucher, Nguyen, et Silar 2017)  
651 and in some fungi closely related to this species complex (*Cercophora samala*, *Triangularia*  
652 *verruculosa* and *Apiosordaria backusii*). However, more distantly related species (*Podospora*  
653 *setosa*, *Podospora australis*, *Podospora appendiculata*, *Thermostelomyces thermophilus*,  
654 *Neurospora tetrasperma* and *Neurospora discreta*) contain a protein that shares some identity  
655 with Pa\_7\_7000, but they all lack the Zn(II)<sub>2</sub>Cys<sub>6</sub> domain and are therefore unlikely to be  
656 functional orthologs. Wave V contains Pa\_6\_11860/*Fluffy* (Bailey et Ebbole 1998) (Fig.  
657 S4AT), which is required in *N. crassa* for the formation of conidia. These vegetative spores are  
658 absent in *P. anserina*, suggesting that this gene has been recruited for ascospore formation in  
659 this fungus, as in others members of the *P. anserina* / *P. pauciseta* / *P. comata* species complex  
660 (Boucher, Nguyen, et Silar 2017).

### 661 **Comparison of mating-type target genes at different developmental stages**

662 A transcriptomic analysis of *P. anserina* was performed previously to determine the FMR1 and  
663 FPR1 target genes required for fertilization (Bidard et al. 2011). The analysis focused on the  
664 comparison of *mat*<sup>-</sup> and *mat*<sup>+</sup> mycelia competent for fertilization, namely the stage preceding  
665 the development of the perithecium. It was performed using the same methods as described  
666 here. The analysis identified 69 genes upregulated in a *mat*<sup>-</sup> mycelium and 88 genes upregulated  
667 in a *mat*<sup>+</sup> mycelium. Among these genes, we search for the DE genes identified in this study.  
668 A total of 32 DE genes (46 %) were identified among the 69 genes which were found to be  
669 upregulated in the *mat*<sup>-</sup> mycelium (Table S8) and 45 DE genes (51%) are found among the  
670 *mat*<sup>+</sup> upregulated genes (Table S9). A statistical analysis indicates that the DE genes are over-  
671 represented in both categories, when compared with the 3,466 DE genes in the 10,507 *P.*  
672 *anserina*'s gene models. This overrepresentation was higher for the *mat*<sup>+</sup> upregulated genes  
673 (Chi<sup>2</sup>, p-value = 0.0003) than for the *mat*<sup>-</sup> upregulated genes (Chi<sup>2</sup>, p-value = 0.0183). This  
674 results is likely due to the use of a *mat*<sup>+</sup> maternal strain for the fruiting body development  
675 presented here and indicates that half of the genes upregulated in fertilization competent  
676 mycelium are also up-regulated in maternal tissues during perithecial development.

## 677 **Discussion**

### 678 **Mating-type gene expression**

679 In this study, we analyzed on a genome-wide scale the transcriptional changes associated with  
680 sexual development in *P. anserina* over a 10 time points course, from fertilization to ascospore  
681 maturation. We found that more than 32% of the *P. anserina*'s genes are differentially

682 expressed during at least one interval covered by this kinetic. Interestingly, more than 36% of  
683 the proteins encoded by these genes do not have a Pfam domain, suggesting that a significant  
684 proportion of them remain functionally uncharacterized. Overall, the data generated by our  
685 study are rich in novel players involved in fungal sexual reproduction and are likely to provide  
686 an exploratory framework to analyze the pathways that are required for the differentiation of  
687 the 15 different tissues identified in a perithecium (Bistis, Perkins, et Read 2003).

688         Among these 15 tissues, the ascogenous hyphae plays a central role in the sexual cycle,  
689 since it is where karyogamy, meiosis and ascospore formation take place. Its formation starts  
690 at T24 (Fig. 1) and requires the expression of the four mating-type genes: *FMRI*, *SMR1*, *SMR2*  
691 and *FPR1* (Zickler et al. 1995; Arnaise et al. 2001). The ratio of ascogenous tissue to the other  
692 tissues of the perithecium is low, potentially masking the zygotic expression profile. This  
693 detection limit of our experiments can be assessed by searching the transcripts of *FMRI*, *SMR1*  
694 and *SMR2*. These three mating-type genes are specific to the *mat*- nuclei, which come from the  
695 male organs and are present only in the ascogonial cell and in the ascogenous hyphae. These  
696 three mating-type genes are listed in the DE genes (Table S1), indicating unambiguously that  
697 transcripts specific to the ascogenous hyphae are detected in our experiments, in spite of its  
698 scarcity. Moreover, upregulations of *FMRI* and *SMR2* transcripts are detected as soon as T24,  
699 correlating with the initial development of ascogenous hyphae (Table S1). In addition, 11 genes  
700 upregulated in *mat*- fertilization competent mycelium are among the DE genes (Table S8),  
701 confirming the detection of transcript from *mat*- male nuclei in the female organ. However, no  
702 significant upregulation of the *mat*- mating-type transcripts is detected in the early steps of  
703 perithecial development, *i.e.* from T0 to the formation of the ascogonial cell at T18 (Fig. 1).  
704 Correspondingly, no wave I DE genes are identified in genes that accumulate transcripts in the  
705 *mat*- mycelium (Table S8). The earliest DE genes upregulated in *mat*- competent mycelium  
706 belong to wave II (n = 4). Wave IV, which corresponds to upregulation of *mat*- mating-type  
707 genes and their correlated genes (Fig. 6), includes 412 genes (Fig. 3), among which six belong  
708 to genes upregulated in *mat*- mycelium (Table S8). Overall, these data suggests that critical  
709 target genes required for the development from fertilization to the formation of ascogonial cells  
710 are not included in the DE genes, because their expression level is below the detection threshold  
711 of microarrays. In contrast, the expression of genes required for the subsequent developmental  
712 stages (differentiation of ascogenous hyphae, meiosis and ascospore formation) are indeed  
713 detected in our experiments.

714 Surprisingly, *FPR1/Pa\_1\_20590*, the *mat*<sup>+</sup> specific mating type gene, does not belong to the  
715 DE genes, although this gene is also required for the formation of ascogenous hyphae (Zickler

716 et al. 1995; Arnaise et al. 2001). It is likely that any changes in expression of this gene in zygotic  
717 tissues is hidden by its transcription in the maternal tissues, which are predominant in the  
718 fruiting body. We have however found 45 DE genes that are specifically upregulated in *mat+*  
719 mycelium competent for fertilization (Table S9). Among these DE genes, 32 are direct or  
720 indirect target genes of FPR1 (Bidard et al. 2011). Their detection proved that FPR1 target  
721 genes do not escape our method, although FPR1 itself is not present in DE genes. Moreover,  
722 detection starts in the early stages of perithecium development as suggested by FPR1 target  
723 genes belonging to wave II (n = 9). Later stages are represented by FPR1 target genes belonging  
724 to wave IV (n = 3) and V (n = 2). However, our experiments do not allow us to determine  
725 unambiguously whether the differential expression of a gene occurs in the maternal or in the  
726 zygotic tissues

### 727 **Secondary metabolite gene are actively regulated during sexual development.**

728 As in *N. crassa* (Wang et al. 2022), *C. globosum* (Wang et al. 2019) and *A. nidulans* (Liu et al.  
729 2021), our results further support the active role of SMs in fungal sexual reproduction. The  
730 evolutionary conservation of their expression profile, *i.e.* downregulation from fertilization to  
731 meiosis followed by a coordinated upregulation during ascospore formation and maturation  
732 suggests that SMs protect fungal progeny, *i.e.* ascospores, from animal predation. However, we  
733 identified several genes of unknown function that were overexpressed during early phases. It is  
734 therefore possible that these metabolites protect newly-formed fructification but also that they  
735 are recruited to mediate developmental processes. This is reminiscent of the *Pa\_5\_2090/pks-6*  
736 gene, which encodes the insecticidal polyketide neurosporin A produced at a similar  
737 developmental stage by *S. macrospora* (Nowrousian 2009) and *N. crassa* (Zhao et al. 2017) to  
738 protect fructifications from arthropod attack. Whether this gene product is also involved in the  
739 process of morphogenesis is still an open question (Wang et al. 2022).

740 Our data reveals a relationship between sterigmatocystin synthesis and mating-type  
741 genes by correlating expression of *PaAflR/Pa\_2\_7360*, *HMG5/Pa\_1\_13940* and  
742 *HMG8/Pa\_6\_4110* (Table 6 and Fig. 6). *PaAflR/Pa\_2\_7360* is a regulator of the  
743 sterigmatocystin synthesis cluster (Shen et al. 2019), while *HMG5/Pa\_1\_13940* and  
744 *HMG8/Pa\_6\_4110* are HMG-domain TFs acting as hubs of regulation for the mating-type  
745 genes and mating-type target genes (Ait Benkhali et al. 2013; Gautier et al. 2018). Our results  
746 suggest that *PaAflR/Pa\_2\_7360* controls positively the transcription of the two other genes.  
747 Over expression of *PaAflR/Pa\_2\_7360* results in sterility, as does the over-expression of  
748 *HMG5/Pa\_1\_13940* (Shen et al. 2019; Ait Benkhali et al. 2013). This result is in agreement

749 with the positive transcriptional control of *HMG5/Pa\_1\_13940* by *PaAflR/Pa\_2\_7360*. To date,  
750 *PaAflR/Pa\_2\_7360* has not been deleted in *P. anserina*, and although its orthologue in *A.*  
751 *nidulans* has been knocked out, the effect on the sexual cycle remains poorly understood (Wang  
752 et al. 2022).

### 753 **Het genes operates important but yet unknown roles in sexual reproduction**

754 This study unveiled an interesting regulation of the *het* genes, which are involved in  
755 allorecognition and thus control the viability of heterokaryons formed after hyphal fusion  
756 during vegetative growth (Daskalov 2023). However, heterokaryotic syncytia are not restricted  
757 to vegetative thalli, as fertilization also leads to the formation of such multinucleate cells, and  
758 accordingly, several *het* loci have been shown to control hybrid fertility in *P. anserina* (Ament-  
759 Velásquez et al. 2020). When we compared the *het* genes expression profiles during sexual  
760 reproduction with those assayed during IR (Bidard, Clavé, et Saupe 2013) and bacterial defense  
761 response (Lamacchia et al. 2016) (Table S6 G), we found that they fall into two categories and  
762 four classes.

763 The first category consists of *het* genes that were always down-regulated during sexual  
764 reproduction but were either up-regulated during IR and/or bacterial confrontation (class 1) or  
765 down-regulated during IR and up-regulated during bacterial confrontation (class 4). This  
766 category therefore contains *het* genes that play a role only in vegetative growth and/or defense  
767 against pathogens, but must otherwise be silenced.

768 The second category consisted of *het* genes that were either up-regulated during IR but  
769 also at some time points during sexual reproduction (class 2) or were not DE during IR and  
770 bacterial confrontation but during sexual reproduction (class 3). Class 2 genes may be recruited  
771 to control cell fusion during morphogenesis of fructifications and/or of early zygotic lineage for  
772 wave II genes, or of the developing asci and ascospores for waves IV and V genes. Class 3  
773 genes may have been recruited to carry out specific programmed cell death events, not related  
774 to hyphal fusion but necessary for sexual morphogenesis. In this context, *Pa\_3\_5550* is of  
775 particular interest since it shows a peak of expression during dikaryon formation. Hce2 proteins  
776 also belong to this category, suggesting a specific role during both the pre-zygotic (*Pa\_2\_4765*)  
777 and post-zygotic (*Pa\_1\_17065*) stages.

778 The *het-6* and *het-13* genes were found to be DE during sexual reproduction in the  
779 heterothallic fungus *N. crassa*, whereas no such observation was made in the homothallic  
780 fungus *C. globosum* (Wang et al. 2019). In *P. anserina*, both orthologs *Pa\_6\_6160/het-6* and  
781 *Pa\_0\_150/het-13* were also DE but showed opposite expression patterns compared to *N. crassa*.

782 This suggests that the roles of *het* genes, although central in the sexual developmental network  
783 of fungi endowed with two different mating types, are polymorphic.

#### 784 **Rewiring genes in the RNAi pathway?**

785 In *P. anserina* (Grognet et al. 2019), as in *T. reesei* (Li, Chen, et Wang 2018) and *A.*  
786 *immersus* (Malagnac et al. 1997), the *rid* (*RIP defective*) ortholog is required for proper sexual  
787 reproduction, which is not the case in *N. crassa*. *Pa\_1\_19440/PaRid* showed a short peak of  
788 expression at T12 (Fig. S4AI), which is consistent with its requirement for the individualization  
789 of the dikaryotic cells leading to meiocytes (Grognet et al. 2019). In contrast, *N. crassa rid* up-  
790 regulation started at dikaryon formation and increased continuously throughout the second half  
791 of sexual reproduction (Wang et al. 2014). Interestingly, the *N. crassa rid* expression pattern  
792 was similar to that of the *P. anserina* DNA methyltransferase encoding gene *PaDnmt5*, which  
793 is absent in *N. crassa*. These results were in agreement with our previous data (Grognet et al.  
794 2019) and strengthened the body of evidence pointing to a role for DNA methylation during  
795 sexual development.

796 Although there is no experimental proof that either somatic or meiotic RNAi pathways  
797 (Quelling or MSUD, respectively) works efficiently in *P. anserina*, some of the corresponding  
798 genes are found to be DE during sexual reproduction. Globally the expression patterns of *P.*  
799 *anserina* genes involved in the RNAi pathways are opposite to those of *N. crassa*, except for  
800 *Pa\_2\_9830/Sad-2* and *Pa\_5\_11380/Sad-5*. During MSUD, *Sad-2* is thought to interact with the  
801 RdRP *Sad-1* as a scaffold. Yet, *Sad-1* and *Sad-2* showed opposite patterns of expression during  
802 *N. crassa* sexual development, which likely discards this hypothesis (Wang et al. 2014).  
803 However, *Sad-5*, which role is yet unknown, has an expression profile that is similar to that of  
804 *Sad-2*. The evolutive conservation of *Pa\_2\_9830/Sad-2* and *Pa\_5\_11380/Sad-5* profiles of  
805 expression in both *N. crassa* and *P. anserina*, suggests that they have been recruited together to  
806 play a role in sexual reproduction. In the homothallic ascomycete *S. macrospora*, *Dcl1*, *Dcl2*,  
807 *Sms2* and *Qde2* are involved in meiocyte formation (Girard et al. 2021), which is not consistent  
808 with the expression pattern of the *Pa\_6\_6150/Dcl-1* but rather with that of *Pa\_5\_8910/Sms-2*.  
809 Altogether, these results suggest a complex role for the RNAi machinery during fungal sexual  
810 reproduction.

#### 811 **Conclusion**

812 Our work reveals 3,466 DE genes during the differentiation of fruiting bodies of *P.*  
813 *anserina*. Only 86 of these DE genes have been previously functionally analyzed (Table S10),

814 leaving a large number of genes for future investigations. Although our analysis does not allow  
815 us to determine the DE genes specifically expressed in each tissue of the fruiting body, our  
816 method is sensitive enough to detect DE genes in ascogenous hyphae, at a stage where this  
817 tissue is represented by few cells in the mass of the fruiting body. The sensitivity of our method  
818 is emphasized by the detection of a peak of transcription of *Pa\_1\_19440/PaRid* at a time of the  
819 early perithecial development corresponding to its expected function. Among the DE genes, we  
820 have identified at least two pathways that may control mating-type genes. The first one involves  
821 *Pa\_7\_7000*, possibly revealing a new regulatory pathway upstream of mating-type genes. The  
822 second one is the relationship between *PaAflR/Pa\_2\_7360*, *HMG5/Pa\_1\_13940* and  
823 *HMG8/Pa\_6\_4110*. The latter two TFs control mating-type genes, thus revealing an additional  
824 putative pathway regulating indirectly these genes. Our analysis reveals also the intriguing  
825 expression patterns of the genes from the MSUD and RNAi pathways, although MSUD has not  
826 been found in *P. anserina* and RNAi not investigated. Further research should be focused in *P.*  
827 *anserina* on the role of the genes involved in these pathways in other species.

## 828 **Acknowledgements**

829 We are grateful to S. Saupe for the exchange of data and ideas.

## 830 **Funding**

831 This work and the salary of F. Bidard were funded by the French National Research Agency  
832 (L'Agence Nationale de la Recherche, ANR) grant number ANR-05-BLAN-0385, project  
833 SexDevMycol, coordinator R. Debuchy.

## 834 **Conflict of interest disclosure**

835 The authors declare that they comply with the PCI rule of having no financial conflicts of  
836 interest in relation to the content of the article.

## 837 **Data, scripts, code, and supplementary information availability**

838 Data are available online: Go to  
839 <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE93094>. Enter token  
840 whcrwemexfkfhqr into the box.

841

842

843 **References**

- 844 Ait Benkhali, Jinane, Evelyne Coppin, Sylvain Brun, Leonardo Peraza-Reyes, Tom Martin,  
845 Christina Dixelius, Nouredine Lazar, Herman van Tilbeurgh, et Robert Debuchy.  
846 2013. « A Network of HMG-Box Transcription Factors Regulates Sexual Cycle in the  
847 Fungus *Podospira anserina* ». *PLoS Genetics* 9 (7): e1003642.  
848 <https://doi.org/10.1371/journal.pgen.1003642>.
- 849 Ament-Velásquez, S. Lorena, Hanna Johannesson, Tatiana Giraud, Robert Debuchy, Sven J.  
850 Saupe, Alfons J.M. Debets, Eric Bastiaans, et al. 2020. « The taxonomy of the model  
851 filamentous fungus *Podospira anserina* ». *MycoKeys* 75 (novembre):51-69.  
852 <https://doi.org/10.3897/mycokeys.75.55968>.
- 853 Arnaise, S., R. Debuchy, et M. Picard. 1997. « What Is a Bona Fide Mating-Type Gene?  
854 Internuclear Complementation of Mat Mutants in *Podospira anserina* ». *Molecular &*  
855 *General Genetics: MGG* 256 (2): 169-78.
- 856 Arnaise, S., D. Zickler, S. Le Bilot, C. Poisier, et R. Debuchy. 2001. « Mutations in Mating-  
857 Type Genes of the Heterothallic Fungus *Podospira anserina* Lead to Self-Fertility ».  
858 *Genetics* 159 (2): 545-56.
- 859 Arnaise, Sylvie, Evelyne Coppin, Robert Debuchy, Denise Zickler, et Marguerite Picard.  
860 1995. « Models for mating type gene functions in *Podospira anserina* ». *Fungal*  
861 *Genetics Newsletter*, 42:79.
- 862 Ashburner, Michael, Catherine A. Ball, Judith A. Blake, David Botstein, Heather Butler, J.  
863 Michael Cherry, Allan P. Davis, et al. 2000. « Gene Ontology: Tool for the  
864 Unification of Biology ». *Nature Genetics* 25 (1): 25-29.  
865 <https://doi.org/10.1038/75556>.
- 866 Bailey, Lori A, et Daniel J Ebbole. 1998. « The fluffy Gene of *Neurospora crassa* Encodes a  
867 Gal4p-Type C6 Zinc Cluster Protein Required for Conidial Development ». *Genetics*  
868 148 (4): 1813-20. <https://doi.org/10.1093/genetics/148.4.1813>.
- 869 Basenko, Evelina Y., Jane A. Pulman, Achchuthan Shanmugasundram, Omar S. Harb,  
870 Kathryn Crouch, David Starns, Susanne Warrenfeltz, et al. 2018. « FungiDB: An  
871 Integrated Bioinformatic Resource for Fungi and Oomycetes ». *Journal of Fungi* 4 (1):  
872 39. <https://doi.org/10.3390/jof4010039>.
- 873 Berteaux-Lecellier, V., D. Zickler, R. Debuchy, A. Panvier-Adoutte, C. Thompson-Coffe, et  
874 M. Picard. 1998. « A Homologue of the Yeast SHE4 Gene Is Essential for the  
875 Transition between the Syncytial and Cellular Stages during Sexual Reproduction of  
876 the Fungus *Podospira anserina* ». *The EMBO Journal* 17 (5): 1248-58.  
877 <https://doi.org/10.1093/emboj/17.5.1248>.
- 878 Bhetariya, Preetida J., Taruna Madan, Seemi Farhat Basir, Anupam Varma, et Sarma P. Usha.  
879 2011. « Allergens/Antigens, Toxins and Polyketides of Important *Aspergillus*  
880 Species ». *Indian Journal of Clinical Biochemistry* 26 (2): 104-19.  
881 <https://doi.org/10.1007/s12291-011-0131-5>.
- 882 Bian, K, Sap Lenz, Q Tang, F Chen, R Qi, M Jost, Cl Drennan, Jm Essigmann, Sd Wetmore,  
883 et D Li. 2019. « DNA Repair Enzymes ALKBH2, ALKBH3, and AlkB Oxidize 5-  
884 Methylcytosine to 5-Hydroxymethylcytosine, 5-Formylcytosine and 5-  
885 Carboxylcytosine in Vitro ». *Nucleic Acids Research* 47 (11).  
886 <https://doi.org/10.1093/nar/gkz395>.
- 887 Bidard, F, S Imbeaud, N Reymond, O Lespinet, P Silar, C Clave, H Delacroix, V Berteaux-  
888 Lecellier, et R Debuchy. 2010. « A general framework for optimization of probes for  
889 gene expression microarray and its application to the fungus *Podospira anserina* » 3  
890 (1): 171.



- 891 Bidard, Frédérique, Jinane Aït Benkhali, Evelyne Coppin, Sandrine Imbeaud, Pierre Grognet,  
892 Hervé Delacroix, et Robert Debuchy. 2011. « Genome-wide gene expression profiling  
893 of fertilization competent mycelium in opposite mating types in the heterothallic  
894 fungus *Podospora anserina* ». *PLoS One* 6 (6): e21476.  
895 <https://doi.org/10.1371/journal.pone.0021476>.
- 896 Bidard, Frédérique, Corinne Clavé, et Sven J. Saupe. 2013. « The Transcriptional Response to  
897 Nonsens in the Fungus *Podospora anserina* ». *G3 (Bethesda, Md.)* 3 (6): 1015-30.  
898 <https://doi.org/10.1534/g3.113.006262>.
- 899 Bistis, George, David Perkins, et Nick Read. 2003. « Different cell types in *Neurospora*  
900 *crassa* ». *Fungal Genetics Reports* 50 (1): 17-19. [https://doi.org/10.4148/1941-  
901 4765.1154](https://doi.org/10.4148/1941-4765.1154).
- 902 Blank-Landeshammer, b, I Teichert, R Märker, M Nowrousian, U Kück, et A Sickmann.  
903 2019. « Combination of Proteogenomics with Peptide De Novo Sequencing Identifies  
904 New Genes and Hidden Posttranscriptional Modifications ». *mBio* 10 (5).  
905 <https://doi.org/10.1128/mBio.02367-19>.
- 906 Boisnard, Stéphanie, Eric Espagne, Denise Zickler, Anne Bourdais, Anne-Laure Riquet, et  
907 Véronique Berteaux-Lecellier. 2009. « Peroxisomal ABC Transporters and Beta-  
908 Oxidation during the Life Cycle of the Filamentous Fungus *Podospora anserina* ». *Fungal Genetics and Biology: FG & B* 46 (1): 55-66.  
909 <https://doi.org/10.1016/j.fgb.2008.10.006>.
- 910 Bonnet, Crystel, Eric Espagne, Denise Zickler, Stéphanie Boisnard, Anne Bourdais, et  
911 Véronique Berteaux-Lecellier. 2006. « The Peroxisomal Import Proteins PEX2, PEX5  
912 and PEX7 Are Differently Involved in *Podospora anserina* Sexual Cycle ». *Molecular*  
913 *Microbiology* 62 (1): 157-69. <https://doi.org/10.1111/j.1365-2958.2006.05353.x>.
- 914 Borkovich, Katherine A., Lisa A. Alex, Oded Yarden, Michael Freitag, Gloria E. Turner,  
915 Nick D. Read, Stephan Seiler, et al. 2004. « Lessons from the Genome Sequence of  
916 *Neurospora crassa*: Tracing the Path from Genomic Blueprint to Multicellular  
917 Organism ». *Microbiology and Molecular Biology Reviews* 68 (1): 1-108.  
918 <https://doi.org/10.1128/mmbr.68.1.1-108.2004>.
- 919 Boucher, Charlie, Tinh-Suong Nguyen, et Philippe Silar. 2017. « Species Delimitation in the  
920 *Podospora anserina*/P. *paucisetata*/P. *comata* Species Complex (Sordariales) ». *Cryptogamie, Mycologie* 38 (4): 485-506.
- 921 Bouhouche, K, D Zickler, R Debuchy, et S Arnaise. 2004. « Altering a gene involved in  
922 nuclear distribution increases the repeat-induced point mutation process in the fungus  
923 *Podospora anserina* ». *Genetics* 167 (1): 151-59.
- 924 Braus, Gerhard H, Stefan Irniger, et Özgür Bayram. 2010. « Fungal development and the  
925 COP9 signalosome ». *Current Opinion in Microbiology, Growth and development: eukaryotes/prokaryotes*, 13 (6): 672-76. <https://doi.org/10.1016/j.mib.2010.09.011>.
- 926 Breyer, Eva, et Federico Baltar. 2023. « The Largely Neglected Ecological Role of Oceanic  
927 Pelagic Fungi ». *Trends in Ecology & Evolution* 38 (9): 870-88.  
928 <https://doi.org/10.1016/j.tree.2023.05.002>.
- 929 Carlier, F., M. Li, L. Maroc, R. Debuchy, C. Souaid, D. Noordermeer, P. Grognet, et F.  
930 Malagnac. 2021. « Loss of EZH2-like or SU(VAR)3-9-like Proteins Causes  
931 Simultaneous Perturbations in H3K27 and H3K9 Tri-Methylation and Associated  
932 Developmental Defects in the Fungus *Podospora anserina* ». *Epigenetics & Chromatin*  
933 14 (1): 22. <https://doi.org/10.1186/s13072-021-00395-7>.
- 934 Cheng, Steven Yan, et J. Michael Bishop. 2002. « Suppressor of Fused represses Gli-  
935 mediated transcription by recruiting the SAP18-mSin3 corepressor complex ». *Proceedings of the National Academy of Sciences* 99 (8): 5442-47.  
936 <https://doi.org/10.1073/pnas.082096999>.

- 941 Chevanne, Damien, Sven J Saupe, Corinne Clavé, et Mathieu Paoletti. 2010. « WD-repeat  
942 instability and diversification of the *Podospora anserina* hnwD non-self recognition  
943 gene family ». *BMC Evolutionary Biology* 10 (mai):134. [https://doi.org/10.1186/1471-](https://doi.org/10.1186/1471-2148-10-134)  
944 2148-10-134.
- 945 Chou, Jui-Yu, Po-Chen Hsu, et Jun-Yi Leu. 2022. « Enforcement of Postzygotic Species  
946 Boundaries in the Fungal Kingdom ». *Microbiology and Molecular Biology Reviews*:  
947 *MMBR* 86 (4): e0009822. <https://doi.org/10.1128/membr.00098-22>.
- 948 Cogoni, C, et G Macino. 1999. « Gene silencing in *Neurospora crassa* requires a protein  
949 homologous to RNA-dependent RNA polymerase ». *Nature* 399 (6732): 166-69.  
950 <https://doi.org/10.1038/20215>.
- 951 Coppin, Evelyne, Véronique Berteaux-Lecellier, Frédérique Bidard, Sylvain Brun, Gwenaél  
952 Ruprich-Robert, Eric Espagne, Jinane Aït-Benkhalil, et al. 2012. « Systematic Deletion  
953 of Homeobox Genes in *Podospora anserina* Uncovers Their Roles in Shaping the  
954 Fruiting Body ». *PloS One* 7 (5): e37488.  
955 <https://doi.org/10.1371/journal.pone.0037488>.
- 956 Cottingham, F. R., L. Gheber, D. L. Miller, et M. A. Hoyt. 1999. « Novel Roles for  
957 *Saccharomyces cerevisiae* Mitotic Spindle Motors ». *The Journal of Cell Biology* 147  
958 (2): 335-50. <https://doi.org/10.1083/jcb.147.2.335>.
- 959 Curto, M.-Ángeles, Mohammad Reza Sharifmoghdam, Eduardo Calpena, Nagore De León,  
960 Marta Hoya, Cristina Doncel, Janet Leatherwood, et M.-Henar Valdivieso. 2014.  
961 « Membrane Organization and Cell Fusion during Mating in Fission Yeast Requires  
962 Multipass Membrane Protein Prm1 ». *Genetics* 196 (4): 1059-76.  
963 <https://doi.org/10.1534/genetics.113.159558>.
- 964 Daskalov, Asen. 2023. « Emergence of the fungal immune system ». *iScience* 26 (6): 106793.  
965 <https://doi.org/10.1016/j.isci.2023.106793>.
- 966 Dyer, Paul S., et Ulrich Kück. 2017. « Sex and the Imperfect Fungi ». *Microbiology Spectrum*  
967 5 (3): 10.1128/microbiolspec.funk-0043-2017.  
968 <https://doi.org/10.1128/microbiolspec.funk-0043-2017>.
- 969 Emms, David M., et Steven Kelly. 2019. « OrthoFinder: Phylogenetic Orthology Inference for  
970 Comparative Genomics ». *Genome Biology* 20 (1): 238.  
971 <https://doi.org/10.1186/s13059-019-1832-y>.
- 972 Erdman, S., L. Lin, M. Malczynski, et M. Snyder. 1998. « Pheromone-Regulated Genes  
973 Required for Yeast Mating Differentiation ». *The Journal of Cell Biology* 140 (3):  
974 461-83. <https://doi.org/10.1083/jcb.140.3.461>.
- 975 Espagne, Eric, Pascale Balhadère, Marie-Louise Penin, Christian Barreau, et Béatrice Turcq.  
976 2002. « HET-E and HET-D belong to a new subfamily of WD40 proteins involved in  
977 vegetative incompatibility specificity in the fungus *Podospora anserina*. » *Genetics*  
978 161 (1): 71-81.
- 979 Espagne, Eric, Olivier Lespinet, Fabienne Malagnac, Corinne Da Silva, Olivier Jaillon, Betina  
980 M Porcel, Arnaud Couloux, et al. 2008. « The genome sequence of the model  
981 ascomycete fungus *Podospora anserina* ». *Genome Biology* 9 (5): R77.  
982 <https://doi.org/10.1186/gb-2008-9-5-r77>.
- 983 Foster, Benjamin M., Paul Stolz, Christopher B. Mulholland, Alex Montoya, Holger Kramer,  
984 Sebastian Bultmann, et Till Bartke. 2018. « Critical Role of the UBL Domain in  
985 Stimulating the E3 Ubiquitin Ligase Activity of UHRF1 toward Chromatin ». *Molecular Cell* 72 (4): 739-752.e9. <https://doi.org/10.1016/j.molcel.2018.09.028>.
- 987 Froehlich, Allan C., Yi Liu, Jennifer J. Loros, et Jay C. Dunlap. 2002. « White Collar-1, a  
988 Circadian Blue Light Photoreceptor, Binding to the Frequency Promoter ». *Science*  
989 (*New York, N.Y.*) 297 (5582): 815-19. <https://doi.org/10.1126/science.1073681>.

- 990 Galagan, James E, Sarah E Calvo, Katherine A Borkovich, Eric U Selker, Nick D Read,  
991 David Jaffe, William FitzHugh, et al. 2003. « The genome sequence of the filamentous  
992 fungus *Neurospora crassa* ». *Nature* 422 (6934): 859-68.  
993 <https://doi.org/10.1038/nature01554>.
- 994 Gautier, Valérie, Laetitia Chan Ho Tong, Tinh-Suong Nguyen, Robert Debuchy, et Philippe  
995 Silar. 2018. « PaPro1 and IDC4, Two Genes Controlling Stationary Phase, Sexual  
996 Development and Cell Degeneration in *Podospora anserina* ». *Journal of Fungi (Basel,*  
997 *Switzerland)* 4 (3): 85. <https://doi.org/10.3390/jof4030085>.
- 998 Girard, Chloe, Karine Budin, Stéphanie Boisnard, Liangran Zhang, Robert Debuchy, Denise  
999 Zickler, et Eric Espagne. 2021. « RNAi-Related Dicer and Argonaute Proteins Play  
1000 Critical Roles for Meicyte Formation, Chromosome-Axes Lengths and Crossover  
1001 Patterning in the Fungus *Sordaria macrospora* ». *Frontiers in Cell and Developmental*  
1002 *Biology* 9 (juin). <https://doi.org/10.3389/fcell.2021.684108>.
- 1003 Gladyshev, Eugene. 2017. « Repeat-Induced Point Mutation and Other Genome Defense  
1004 Mechanisms in Fungi ». *Microbiology Spectrum* 5 (4).  
1005 <https://doi.org/10.1128/microbiolspec.FUNK-0042-2017>.
- 1006 Grognet, Pierre, Hélène Timpano, Florian Carlier, Jinane Aït-Benkhalil, Véronique Berteaux-  
1007 Lecellier, Robert Debuchy, Frédérique Bidard, et Fabienne Malagnac. 2019. « A RID-  
1008 like Putative Cytosine Methyltransferase Homologue Controls Sexual Development in  
1009 the Fungus *Podospora anserina* ». *PLoS Genetics* 15 (8): e1008086.  
1010 <https://doi.org/10.1371/journal.pgen.1008086>.
- 1011 He, Qiyang, Ping Cheng, Yuhong Yang, Lixing Wang, Kevin H. Gardner, et Yi Liu. 2002.  
1012 « White Collar-1, a DNA Binding Transcription Factor and a Light Sensor ». *Science*  
1013 *(New York, N.Y.)* 297 (5582): 840-43. <https://doi.org/10.1126/science.1072795>.
- 1014 Hutchison, Elizabeth A, et N Louise Glass. 2010. « Meiotic Regulators Ndt80 and Ime2 Have  
1015 Different Roles in *Saccharomyces* and *Neurospora* ». *Genetics* 185 (4): 1271-82.  
1016 <https://doi.org/10.1534/genetics.110.117184>.
- 1017 Iliiev, Iliyan D., Gordon D. Brown, Petra Bacher, Sarah L. Gaffen, Joseph Heitman, Bruce S.  
1018 Klein, et Michail S. Lionakis. 2024. « Focus on Fungi ». *Cell* 187 (19): 5121-27.  
1019 <https://doi.org/10.1016/j.cell.2024.08.016>.
- 1020 Julius, D., A. Brake, L. Blair, R. Kunisawa, et J. Thorner. 1984. « Isolation of the Putative  
1021 Structural Gene for the Lysine-Arginine-Cleaving Endopeptidase Required for  
1022 Processing of Yeast Prepro-Alpha-Factor ». *Cell* 37 (3): 1075-89.  
1023 [https://doi.org/10.1016/0092-8674\(84\)90442-2](https://doi.org/10.1016/0092-8674(84)90442-2).
- 1024 Kaneko, Isao, Karine Dementhon, Qijun Xiang, et N. Louise Glass. 2006. « Nonallelic  
1025 Interactions between Het-c and a Polymorphic Locus, Pin-c, Are Essential for Nonsell  
1026 Recognition and Programmed Cell Death in *Neurospora crassa* ». *Genetics* 172 (3):  
1027 1545-55. <https://doi.org/10.1534/genetics.105.051490>.
- 1028 Kim, Wonyong, Zheng Wang, Hyeonjae Kim, Kasey Pham, Yujia Tu, Jeffrey P. Townsend,  
1029 et Frances Trail. 2022. « Transcriptional Divergence Underpinning Sexual  
1030 Development in the Fungal Class Sordariomycetes ». *mBio*, mai, e0110022.  
1031 <https://doi.org/10.1128/mbio.01100-22>.
- 1032 Kurahashi, Hiroshi, Yoshiyuki Imai, et Masayuki Yamamoto. 2002. « Tropomyosin Is  
1033 Required for the Cell Fusion Process during Conjugation in Fission Yeast ». *Genes to*  
1034 *Cells: Devoted to Molecular & Cellular Mechanisms* 7 (4): 375-84.  
1035 <https://doi.org/10.1046/j.1365-2443.2002.00526.x>.
- 1036 Lamacchia, Marina, Witold Dyrka, Annick Breton, Sven J. Saupe, et Mathieu Paoletti. 2016.  
1037 « Overlapping *Podospora anserina* Transcriptional Responses to Bacterial and Fungal  
1038 Non Self Indicate a Multilayered Innate Immune Response ». *Frontiers in*  
1039 *Microbiology* 7:471. <https://doi.org/10.3389/fmicb.2016.00471>.

- 1040 Leipe, Detlef D., Eugene V. Koonin, et L. Aravind. 2004. « STAND, a Class of P-Loop  
1041 NTPases Including Animal and Plant Regulators of Programmed Cell Death: Multiple,  
1042 Complex Domain Architectures, Unusual Phyletic Patterns, and Evolution by  
1043 Horizontal Gene Transfer ». *Journal of Molecular Biology* 343 (1): 1-28.  
1044 <https://doi.org/10.1016/j.jmb.2004.08.023>.
- 1045 Lelandais, Gaëlle, Damien Remy, Fabienne Malagnac, et Pierre Grognet. 2022. « New  
1046 Insights into Genome Annotation in *Podospira anserina* through Re-Exploiting  
1047 Multiple RNA-Seq Data ». *BMC Genomics* 23 (1): 1-19.  
1048 <https://doi.org/10.1186/s12864-022-09085-4>.
- 1049 Li, Wan-Chen, Chia-Ling Chen, et Ting-Fang Wang. 2018. « Repeat-Induced Point (RIP)  
1050 Mutation in the Industrial Workhorse Fungus *Trichoderma reesei* ». *Applied*  
1051 *Microbiology and Biotechnology*, janvier. <https://doi.org/10.1007/s00253-017-8731-5>.
- 1052 Liu, Li, Christoph Sasse, Benedict Dirnberger, Oliver Valerius, Enikő Fekete-Szücs, Rebekka  
1053 Harting, Daniela E Nordzieke, et al. 2021. « Secondary metabolites of *Hülle* cells  
1054 mediate protection of fungal reproductive and overwintering structures against  
1055 fungivorous animals ». Édité par Antonis Rokas, Detlef Weigel, Milton Drott, et Jae-  
1056 Hyuk Yu. *eLife* 10 (octobre):e68058. <https://doi.org/10.7554/eLife.68058>.
- 1057 Lütkenhaus, Ramona, Stefanie Traeger, Jan Breuer, Laia Carreté, Alan Kuo, Anna Lipzen,  
1058 Jasmyn Pangilinan, et al. 2019. « Comparative Genomics and Transcriptomics to  
1059 Analyze Fruiting Body Development in Filamentous Ascomycetes ». *Genetics*,  
1060 octobre. <https://doi.org/10.1534/genetics.119.302749>.
- 1061 MacPherson, Sarah, Marc Laroche, et Bernard Turcotte. 2006. « A Fungal Family of  
1062 Transcriptional Regulators: the Zinc Cluster Proteins ». *Microbiology and Molecular*  
1063 *Biology Reviews* 70 (3): 583-604. <https://doi.org/10.1128/mubr.00015-06>.
- 1064 Malagnac, F, B Wendel, C Goyon, G Faugeron, D Zickler, J L Rossignol, M Noyer-Weidner,  
1065 P Vollmayr, T A Trautner, et J Walter. 1997. « A gene essential for de novo  
1066 methylation and development in *Ascobolus* reveals a novel type of eukaryotic DNA  
1067 methyltransferase structure ». *Cell* 91 (2): 281-90.
- 1068 Mistry, Jaina, Sara Chuguransky, Lowri Williams, Matloob Qureshi, Gustavo A. Salazar, Erik  
1069 L. L. Sonnhammer, Silvio C. E. Tosatto, et al. 2021. « Pfam: The Protein Families  
1070 Database in 2021 ». *Nucleic Acids Research* 49 (D1): D412-19.  
1071 <https://doi.org/10.1093/nar/gkaa913>.
- 1072 Nelson, M. A., S. T. Merino, et R. L. Metzberg. 1997. « A Putative Rhamnogalacturonase  
1073 Required for Sexual Development of *Neurospora crassa* ». *Genetics* 146 (2): 531-40.  
1074 <https://doi.org/10.1093/genetics/146.2.531>.
- 1075 Nowrousian, Minou. 2009. « A Novel Polyketide Biosynthesis Gene Cluster Is Involved in  
1076 Fruiting Body Morphogenesis in the Filamentous Fungi *Sordaria Macrospora* and  
1077 *Neurospora crassa* ». *Current Genetics* 55 (2): 185-98. <https://doi.org/10.1007/s00294-009-0236-z>.
- 1078 ———. 2022. « The Role of Chromatin and Transcriptional Control in the Formation of  
1079 Sexual Fruiting Bodies in Fungi ». *Microbiology and Molecular Biology Reviews*:  
1080 *MMBR* 86 (4): e0010422. <https://doi.org/10.1128/mubr.00104-22>.
- 1081 Oh, Yong-Seok, Pu Gao, Ko-Woon Lee, Ilaria Ceglia, Ji-Seon Seo, Xiaozhu Zhang, Jung-  
1082 Hyuck Ahn, et al. 2013. « SMARCA3, a Chromatin-Remodeling Factor, Is Required  
1083 for P11-Dependent Antidepressant Action ». *Cell* 152 (4): 831-43.  
1084 <https://doi.org/10.1016/j.cell.2013.01.014>.
- 1085 Ottmann, Christian, Borries Lubracki, Isabell Küfner, Wolfgang Koch, Frédéric Brunner,  
1086 Michael Weyand, Laura Mattinen, et al. 2009. « A common toxin fold mediates  
1087 microbial attack and plant defense ». *Proceedings of the National Academy of*  
1088 *Sciences* 106 (25): 10359-64. <https://doi.org/10.1073/pnas.0902362106>.
- 1089

- 1090 Otto, Sarah P. 2009. « The Evolutionary Enigma of Sex ». *The American Naturalist* 174  
1091 Suppl 1 (juillet):S1-14. <https://doi.org/10.1086/599084>.
- 1092 Pearl, LH. 2000. « Structure and Function in the Uracil-DNA Glycosylase Superfamily ». *Mutation Research* 460 (3-4). [https://doi.org/10.1016/s0921-8777\(00\)00025-2](https://doi.org/10.1016/s0921-8777(00)00025-2).
- 1093 Peraza-Reyes, Leonardo, Sylvie Arnaise, Denise Zickler, Evelyne Coppin, Robert Debuchy,  
1094 et Véronique Berteaux-Lecellier. 2011. « The Importomer Peroxins Are Differentially  
1095 Required for Peroxisome Assembly and Meiotic Development in *Podospora anserina*:  
1096 Insights into a New Peroxisome Import Pathway ». *Molecular Microbiology* 82 (2):  
1097 365-77. <https://doi.org/10.1111/j.1365-2958.2011.07816.x>.
- 1098 Peraza-Reyes, Leonardo, et Fabienne Malagnac. 2016. *Sexual Development in Fungi*. The  
1099 Mycota I, Growth, Differentiation and Sexuality. Springer.
- 1100 Romano, Nicoletta, et Giuseppe Macino. 1992. « Quelling: Transient Inactivation of Gene  
1101 Expression in *Neurospora crassa* by Transformation with Homologous Sequences ». *Molecular Microbiology* 6 (22): 3343-53. <https://doi.org/10.1111/j.1365-2958.1992.tb02202.x>.
- 1102  
1103  
1104
- 1105 Rose, M. D., L. M. Misra, et J. P. Vogel. 1989. « KAR2, a Karyogamy Gene, Is the Yeast  
1106 Homolog of the Mammalian BiP/GRP78 Gene ». *Cell* 57 (7): 1211-21.  
1107 [https://doi.org/10.1016/0092-8674\(89\)90058-5](https://doi.org/10.1016/0092-8674(89)90058-5).
- 1108 Shen, Ling, François-Hugues Porée, Thomas Gaslonde, Hervé Lalucque, Florence Chapeland-  
1109 Leclerc, et Gwenaël Ruprich-Robert. 2019. « Functional Characterization of the  
1110 Sterigmatocystin Secondary Metabolite Gene Cluster in the Filamentous Fungus  
1111 *Podospora anserina*: Involvement in Oxidative Stress Response, Sexual Development,  
1112 Pigmentation and Interspecific Competitions ». *Environmental Microbiology* 21 (8):  
1113 3011-26. <https://doi.org/10.1111/1462-2920.14698>.
- 1114 Shiu, P K, N B Raju, D Zickler, et R L Metzenberg. 2001. « Meiotic silencing by unpaired  
1115 DNA ». *Cell* 107 (7): 905-16.
- 1116 Slot, Jason C., et Antonis Rokas. 2011. « Horizontal Transfer of a Large and Highly Toxic  
1117 Secondary Metabolic Gene Cluster between Fungi ». *Current Biology: CB* 21 (2):  
1118 134-39. <https://doi.org/10.1016/j.cub.2010.12.020>.
- 1119 Stajich, Jason E., Mary L. Berbee, Meredith Blackwell, David S. Hibbett, Timothy Y. James,  
1120 Joseph W. Spatafora, et John W. Taylor. 2009. « The Fungi ». *Current Biology* 19  
1121 (18): R840-45. <https://doi.org/10.1016/j.cub.2009.07.004>.
- 1122 Steffens, Eva Katharina, Kordula Becker, Sabine Krevet, Ines Teichert, et Ulrich Kück. 2016.  
1123 « Transcription Factor PRO1 Targets Genes Encoding Conserved Components of  
1124 Fungal Developmental Signaling Pathways ». *Molecular Microbiology* 102 (5):  
1125 792-809. <https://doi.org/10.1111/mmi.13491>.
- 1126 Stergiopoulos, Ioannis, Jérôme Collemare, Rahim Mehrabi, et Pierre J. G. M. De Wit. 2013.  
1127 « Phytotoxic Secondary Metabolites and Peptides Produced by Plant Pathogenic  
1128 Dothideomycete Fungi ». *FEMS Microbiology Reviews* 37 (1): 67-93.  
1129 <https://doi.org/10.1111/j.1574-6976.2012.00349.x>.
- 1130 Sun, Xianyun, Luning Yu, Nan Lan, Shiping Wei, Yufei Yu, Hanxing Zhang, Xinyu Zhang,  
1131 et Shaojie Li. 2012. « Analysis of the role of transcription factor VAD-5 in conidiation  
1132 of *Neurospora crassa* ». *Fungal Genetics and Biology* 49 (5): 379-87.  
1133 <https://doi.org/10.1016/j.fgb.2012.03.003>.
- 1134 Takano-Rojas, Harumi, Denise Zickler, et Leonardo Peraza-Reyes. 2016. « Peroxisome  
1135 Dynamics during Development of the Fungus *Podospora anserina* ». *Mycologia* 108  
1136 (3): 590-602. <https://doi.org/10.3852/15-112>.
- 1137 Teichert, Ines, Tim A. Dahlmann, Ulrich Kück, et Minou Nowrousian. 2017. « RNA Editing  
1138 During Sexual Development Occurs in Distantly Related Filamentous Ascomycetes ». *Genome Biology and Evolution* 9 (4): 855-68. <https://doi.org/10.1093/gbe/evx052>.
- 1139

- 1140 Teichert, Ines, Stefanie Pöggeler, et Minou Nowrousian. 2020. « Sordaria Macrospora:  
1141 25 Years as a Model Organism for Studying the Molecular Mechanisms of Fruiting  
1142 Body Development ». *Applied Microbiology and Biotechnology* 104 (9): 3691-3704.  
1143 <https://doi.org/10.1007/s00253-020-10504-3>.
- 1144 Teichert, Ines, Gabriele Wolff, Ulrich Kück, et Minou Nowrousian. 2012. « Combining laser  
1145 microdissection and RNA-seq to chart the transcriptional landscape of fungal  
1146 development ». *BMC Genomics* 13 (1): 511. <https://doi.org/10.1186/1471-2164-13-511>.
- 1148 The Gene Ontology Consortium, Suzi A Aleksander, James Balhoff, Seth Carbon, J Michael  
1149 Cherry, Harold J Drabkin, Dustin Ebert, et al. 2023. « The Gene Ontology  
1150 knowledgebase in 2023 ». *Genetics* 224 (1): iyad031.  
1151 <https://doi.org/10.1093/genetics/iyad031>.
- 1152 Thompson-Coffe, C., G. Borioli, D. Zickler, et A. L. Rosa. 1999. « Pyruvate Decarboxylase  
1153 Filaments Are Associated with the Cortical Cytoskeleton of Asci and Spores over the  
1154 Sexual Cycle of Filamentous Ascomycetes ». *Fungal Genetics and Biology: FG & B*  
1155 26 (1): 71-80. <https://doi.org/10.1006/fgbi.1998.1106>.
- 1156 Tong, Kevin, Thomas Keller, Charles S. Hoffman, et Anthony T. Annunziato. 2012.  
1157 « Schizosaccharomyces Pombe Hat1 (Kat1) Is Associated with Mis16 and Is Required  
1158 for Telomeric Silencing ». *Eukaryotic Cell* 11 (9): 1095-1103.  
1159 <https://doi.org/10.1128/EC.00123-12>.
- 1160 Turgeon, B Gillian, et R. Debuchy. 2007. « Cochliobolus and Podospora: mechanism of sex  
1161 determination and the evolution of reproductive lifestyle ». In *Sex in Fungi, Molecular  
1162 Determination and Evolutionary Implications*.
- 1163 Wang, Amy, Siavash K. Kurdistani, et Michael Grunstein. 2002. « Requirement of Hos2  
1164 Histone Deacetylase for Gene Activity in Yeast ». *Science (New York, N.Y.)* 298  
1165 (5597): 1412-14. <https://doi.org/10.1126/science.1077790>.
- 1166 Wang, Chengshu, et Raymond J. St Leger. 2007. « The MAD1 Adhesin of Metarhizium  
1167 Anisopliae Links Adhesion with Blastospore Production and Virulence to Insects, and  
1168 the MAD2 Adhesin Enables Attachment to Plants ». *Eukaryotic Cell* 6 (5): 808-16.  
1169 <https://doi.org/10.1128/EC.00409-06>.
- 1170 Wang, Peng, Jia Xu, Perng-Kuang Chang, Zhemin Liu, et Qing Kong. 2022. « New Insights  
1171 of Transcriptional Regulator AflR in Aspergillus flavus Physiology ». *Microbiology  
1172 Spectrum* 10 (1): e00791-21. <https://doi.org/10.1128/spectrum.00791-21>.
- 1173 Wang, Zheng, Aditya Gudibanda, Ugochukwu Ugwuowo, Frances Trail, et Jeffrey P.  
1174 Townsend. 2018. « Using evolutionary genomics, transcriptomics, and systems  
1175 biology to reveal gene networks underlying fungal development ». *Fungal Biology  
1176 Reviews, Complex multicellularity in Fungi*, 32 (4): 249-64.  
1177 <https://doi.org/10.1016/j.fbr.2018.02.001>.
- 1178 Wang, Zheng, Francesc Lopez-Giraldez, Nina Lehr, Marta Farré, Ralph Common, Frances  
1179 Trail, et Jeffrey P Townsend. 2014. « Global Gene Expression and Focused Knockout  
1180 Analysis Reveals Genes Associated with Fungal Fruiting Body Development in  
1181 Neurospora crassa ». *Eukaryotic Cell* 13 (1): 154-69.  
1182 <https://doi.org/10.1128/EC.00248-13>.
- 1183 Wang, Zheng, Francesc Lopez-Giraldez, Jason Slot, Oded Yarden, Frances Trail, et Jeffrey P.  
1184 Townsend. 2022. « Secondary Metabolism Gene Clusters Exhibit Increasingly  
1185 Dynamic and Differential Expression during Asexual Growth, Conidiation, and Sexual  
1186 Development in Neurospora crassa ». *mSystems* 7 (3): e00232-22.  
1187 <https://doi.org/10.1128/msystems.00232-22>.
- 1188 Wang, Zheng, Francesc López-Giráldez, Junrui Wang, Frances Trail, et Jeffrey P. Townsend.  
1189 2019. « Integrative Activity of Mating Loci, Environmentally Responsive Genes, and

- 1190 Secondary Metabolism Pathways during Sexual Development of *Chaetomium*  
1191 *globosum* ». *mBio* 10 (6): 10.1128/mbio.02119-19.  
1192 <https://doi.org/10.1128/mbio.02119-19>.
- 1193 White, B. T., et C. Yanofsky. 1993. « Structural Characterization and Expression Analysis of  
1194 the *Neurospora* Conidiation Gene *Con-6* ». *Developmental Biology* 160 (1): 254-64.  
1195 <https://doi.org/10.1006/dbio.1993.1303>.
- 1196 Wilson, Andi M., P. Markus Wilken, Magriet A. van der Nest, Michael J. Wingfield, et  
1197 Brenda D. Wingfield. 2019. « It's All in the Genes: The Regulatory Pathways of  
1198 Sexual Reproduction in Filamentous Ascomycetes ». *Genes* 10 (5): 330.  
1199 <https://doi.org/10.3390/genes10050330>.
- 1200 Yang, Qiuying, Qiaohong Anne Ye, et Yi Liu. 2015. « Mechanism of siRNA Production from  
1201 Repetitive DNA ». *Genes & Development* 29 (5): 526-37.  
1202 <https://doi.org/10.1101/gad.255828.114>.
- 1203 Young, B. P., R. A. Craven, P. J. Reid, M. Willer, et C. J. Stirling. 2001. « Sec63p and Kar2p  
1204 Are Required for the Translocation of SRP-Dependent Precursors into the Yeast  
1205 Endoplasmic Reticulum in Vivo ». *The EMBO Journal* 20 (1-2): 262-71.  
1206 <https://doi.org/10.1093/emboj/20.1.262>.
- 1207 Zhao, Yanxia, Jianing Ding, Weihua Yuan, Jinjin Huang, Wenxiu Huang, Yan Wang, et  
1208 Weifa Zheng. 2017. « Production of a fungal furocoumarin by a polyketide synthase  
1209 gene cluster confers the chemo-resistance of *Neurospora crassa* to the predation by  
1210 fungivorous arthropods ». *Environmental Microbiology* 19 (10): 3920-29.  
1211 <https://doi.org/10.1111/1462-2920.13791>.
- 1212 Zheng, Qian, Rui Hou, null Juanyu, null Zhang, Jiwen Ma, Zhongshou Wu, Guanghui Wang,  
1213 Chenfang Wang, et Jin-Rong Xu. 2013. « The MAT Locus Genes Play Different  
1214 Roles in Sexual Reproduction and Pathogenesis in *Fusarium graminearum* ». *PloS*  
1215 *One* 8 (6): e66980. <https://doi.org/10.1371/journal.pone.0066980>.
- 1216 Zickler, D., S. Arnais, E. Coppin, R. Debuchy, et M. Picard. 1995. « Altered Mating-Type  
1217 Identity in the Fungus *Podospira anserina* Leads to Selfish Nuclei, Uniparental  
1218 Progeny, and Haploid Meiosis ». *Genetics* 140 (2): 493-503.  
1219