



Overexpression of POLRMT increases mitochondrial transcription initiation in mouse

Maria Miranda, Andrea Mesaros, Louise Pérard, Irina Kuznetsov, Martin Purrio, Ilian Atanassov, Aleksandra Filipovska, Arnaud Mourier, Inge Kuhl

► To cite this version:

Maria Miranda, Andrea Mesaros, Louise Pérard, Irina Kuznetsov, Martin Purrio, et al.. Overexpression of POLRMT increases mitochondrial transcription initiation in mouse. Gordon Research Conference 2024 - Mitochondria and Chloroplasts: Energy-Transducing Organelles: Fundamental Processes and Translation in Agriculture and Medicine, Jul 2024, Barcelone, Spain. . hal-04943900

HAL Id: hal-04943900

<https://hal.science/hal-04943900v1>

Submitted on 12 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OVEREXPRESSION OF POLRMT INCREASES MITOCHONDRIAL TRANSCRIPTION INITIATION IN MOUSE

Maria Miranda^{1,6}, Andrea Mesaros¹, Louise Pérard², Irina Kuznetsov³, Martin Purrio¹, Ilian Atanassov¹, Aleksandra Filipovska⁴, Arnaud Mourier⁵, Inge Kühl^{2*}

1 Max Planck Institute for Biology of Ageing, 50931 Cologne, Germany; 2 Institute of Integrative Biology of the Cell, UMR9198, CEA, CNRS, Université Paris-Saclay, 91190 Gif-sur-Yvette, France; 3 Harry Perkins Institute of Medical Research and Centre for Medical Research, The University of Western Australia, Nedlands 6009, Australia and School of Molecular Sciences, The University of Western Australia, 6009 Crawley, Australia; 4 Telethon Kids Institute, Northern Entrance, Perth Children's Hospital, 15 Hospital Avenue, Nedlands, Western Australia, 6009, Australia; 5 Université de Bordeaux and the Centre National de la Recherche Scientifique, Institut de Biochimie et Génétique Cellulaires UMR 5095, 33077 Bordeaux, France; 6 Current address: Massachusetts General Hospital, Boston, MA 02114, USA, Harvard Medical School, Boston, MA 02115, USA, Broad Institute of MIT and Harvard, Cambridge, MA, 02142.

*inge.kuhl@i2bc.paris-saclay.fr

POLRMT is the sole RNA polymerase in human mitochondria where it generates primers for mitochondrial DNA (mtDNA) replication and transcribes the mitochondrial genome to express genes encoding essential components of the oxidative phosphorylation (OXPHOS) system. Elevated POLRMT levels are found in several cancers and in mouse models with severe mitochondrial dysfunction. Here, we generated and characterized mice over-expressing Polrmt to investigate the physiological and molecular consequences of elevated POLRMT levels. Increasing POLRMT did not result in any pathological phenotype but instead positively affected exercise capacity under stress conditions. POLRMT overexpression increased in organello transcription initiation, resulting in higher steady-state levels of the promoter-proximal L-strand transcript 7S RNA and higher mtDNA levels. Surprisingly, the abundance of mature mitochondrial RNAs was not affected by the elevated POLRMT levels. Furthermore, ubiquitous simultaneous overexpression of POLRMT and LRPPRC, which stabilizes mitochondrial messenger RNAs, did not increase steady-state levels of mitochondrial transcripts in the mouse. Our data show that POLRMT levels regulate transcription initiation, and that additional regulatory steps downstream of transcription initiation and transcript stability limit OXPHOS biogenesis.

THE MAMMALIAN MITOCHONDRIAL GENOME

