



HAL
open science

Active depinning of bacterial droplets: The collective surfing of *Bacillus subtilis*

Marc Hennes, Julien Tailleur, Gaëlle Charron, Adrian Daerr

► **To cite this version:**

Marc Hennes, Julien Tailleur, Gaëlle Charron, Adrian Daerr. Active depinning of bacterial droplets: The collective surfing of *Bacillus subtilis*. Congrès des 150 ans de la SFP, Jul 2023, Paris, France. . hal-04947043

HAL Id: hal-04947043

<https://hal.science/hal-04947043v1>

Submitted on 14 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License


Active depinning of bacterial droplets: the collective surfing of *Bacillus subtilis*

Context: Surface translocation of microbes



Swarming *B. subtilis*


- Rich phenomenology, mechanisms unclear
- Often sharp colony boundary: no leaving colony alone
- Movement strongly constrained by interfacial forces





- Role of wetting and contact line depinning ?

Wetting of bacterial droplets


- Wetting hysteresis pins even millimetric water droplets to vertical



- Bacteria depin their drop




- Bacteria achieve depinning on very shallow slopes ...




- ... initial droplet of bacterial suspension
- ... water

- ... but this colony surfing occurs only after several hours.


B. subtilis produces biosurfactant surfactin → reduces surface tension



Moving grid profilometry



Drop volume increases prior to depinning



- slow volume decrease for water
- the more bacteria ($OD \approx 1.5 > OD \approx 0.27$), the faster volume increases
- supernatant (SN = bacteria removed) behaves the same initially
- surfactant deficient strain 168 ⇒ no volume increase, as for water
- pure surfactant solution ⇒ volume increase

Bacterial droplets harness 3 distinct effects to unpin



drop depinning condition:

Weight > Contact line pinning force

$$\rho V g \sin \alpha > \text{cst.} \cdot V^{1/3} \gamma (\cos \theta_r - \cos \theta_a)$$

Volume ↑, Surface tension ↓ ⇒ $Bo = \rho V^{2/3} g \sin \alpha / \gamma$ increases,


Wetting hysteresis ↓ ⇒ critical Bond from 0.2 down to ~0.03



Control experiments

- How many bacteria are needed ?


Fraction of colony surfing events ($N = 5$) for different bacterial concentrations (OD) in the deposited drop.



- $OD \leq 0.02$ no surfing (out of five exp.)
- $OD \geq 0.27$ surfing always occurs

- Are bacteria really needed ?

- Centrifuge & filter, keep supernatant (SN)
- Put droplets of bacterial suspension and of SN side-by-side




⇒ Contamination triggers drop sliding

- Do propelling forces contribute to depinning ?

Pressure from active swimmers could help depin contact line

⇒ Compare with surfing probability for non-flagellated mutant



⇒ Colony surfing independent of individual motility