



HAL
open science

A new species of *Androctonus* Ehrenberg, 1828 from the Tassili n'Ajjer, Algeria (Scorpiones: Buthidae).

Eric Ythier, Salah Eddine Sadine, Youcef Alioua, Wilson R. Lourenço

► To cite this version:

Eric Ythier, Salah Eddine Sadine, Youcef Alioua, Wilson R. Lourenço. A new species of *Androctonus* Ehrenberg, 1828 from the Tassili n'Ajjer, Algeria (Scorpiones: Buthidae).. *Faunitaxys*, 2025, 13 (8), pp.1-8. 10.57800/faunitaxys-13(08) . hal-04950545

HAL Id: hal-04950545

<https://hal.science/hal-04950545v1>

Submitted on 16 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*



Volume 13
Numéro 08

Février 2025

ISSN (Print) : 2269 - 6016
ISSN (Online) : 2970 - 4960
Dépôt légal : Février 2025

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*

ZooBank : <https://zoobank.org/79A36B2E-F645-4F9A-AE2B-ED32CE6771CC>

Directeur de la publication, rédacteur, conception graphique et PAO:

Lionel Delaunay

Cette revue ne peut pas être vendue

Elle est distribuée par échange aux institutions (version papier)

et sur simple demande aux particuliers (format PDF)

à l'adresse suivante:

AFCFF (Association française de Cartographie de la Faune et de la Flore)

28, rue Voltaire, F- 42100 Saint Etienne

E-mail: faunitaxys.journal@gmail.com

Elle est disponible librement au téléchargement à partir du site:

<https://faunitaxys.fr/>

La parution de *Faunitaxys* est apériodique

Faunitaxys est indexé dans / *Faunitaxys* is indexed in:

- **Zoological Record**

Articles and nomenclatural novelties are referenced by:

- **ZooBank** (<https://zoobank.org>)

Online Archives:

- **HAL** (<https://hal.archives-ouvertes.fr>)

- **Internet Archive** (<https://archive.org/details/faunitaxys>)

Imprimée sur les presses de SPEED COPIE, 6, rue Tréfilerie, F- 42100 Saint-Etienne

Imprimé le 15 02 2025

A new species of *Androctonus* Ehrenberg, 1828 from the Tassili n'Ajjer, Algeria (Scorpiones: Buthidae)

ERIC YTHIER (1, *), SALAH EDDINE SADINE (2, 3), YUCEF ALIOUA (2, 4) & WILSON R. LOURENÇO (5)

(1) BYG Taxa, 382 rue des Guillates, 71570 Romanèche-Thorins, France.

– E-mail: contact@bygtaxa.com – ZooBank: <https://zoobank.org/06FD0852-A88E-49E5-B8E6-E1494B86C4E1> – Orcid: <https://orcid.org/0000-0002-3194-5184>

(2) Laboratoire Valorisation et Conservation des Ecosystèmes Arides (LVCEA), Faculté des Sciences de la Nature et de la Vie et Sciences de la terre, Université de Ghardaïa, 47000 Ghardaïa, Algeria.

(3) – E-mail: sse.scorpion@yahoo.fr – ZooBank: <https://zoobank.org/45564AC0-B7ED-4C1E-95DD-9EAB3CED9336> – Orcid: <https://orcid.org/0000-0002-0026-8280>

(4) – E-mail: yucef900@yahoo.fr – ZooBank: <https://zoobank.org/597CDBA8-7FAE-49B9-B8B7-516091847F96> – Orcid: <https://orcid.org/0000-0003-2548-0795>

(5) Muséum national d'Histoire naturelle, Sorbonne Universités, Institut de Systématique, Evolution, Biodiversité (ISYEB), UMR7205-CNRS, MNHN, UPMC, EPHE, CP 53, 57 rue Cuvier, 75005 Paris, France.

– E-mail: wilson.lourenco@mnhn.fr – ZooBank: <https://zoobank.org/58448BD6-79D7-46CE-AFDD-91EFF2B7D4EF> – Orcid: <https://orcid.org/0000-0002-2386-363X>

* Correspondence.

Keywords:

Scorpion;
Buthidae;
Androctonus;
ajjer;
taxonomy;
new species;
description;
morphology;
Saharan massifs;
Tassili n'Ajjer;
Algeria.

Abstract. – A new species of *Androctonus* Ehrenberg, 1828 is described on the basis of one male specimen collected in Eferi, at the foothills of the Tassili n'Ajjer massif, Wilaya of Djanet, Algeria. This new scorpion taxon represents the 42nd known species of the genus *Androctonus* and the 7th reported from Algeria. A geographical distribution map of the *Androctonus* species occurring in the region of the Saharan Mountain ranges of Algeria and Niger is presented.

Ythier E., Sadine S. E., Alioua Y. & Lourenço W. R., 2025. – A new species of *Androctonus* Ehrenberg, 1828 from the Tassili n'Ajjer, Algeria (Scorpiones: Buthidae). *Faunitaxys*, 13(08): 1 – 8.

DOI: [https://doi.org/10.57800/faunitaxys-13\(08\)](https://doi.org/10.57800/faunitaxys-13(08))

ZooBank: <https://zoobank.org/5F076343-9D35-4918-87C6-3F44E948B588>

Received: 05/02/2025 – Revised: 10/02/2025 – Accepted: 11/02/2025

Introduction

Since the monographic work of Vachon (1952) on North African scorpions, the scorpion fauna of Algeria has been the subject of several studies, leading to an increasing number of new species as well as a new genus (Abidi *et al.*, 2021; Bouisset & Larrouy, 1962; Khammassi *et al.*, 2023; Lourenço, 2001, 2002a,b, 2006, 2010, 2013a,b, 2017; Lourenço & Leguin, 2011, 2014; Lourenço & Rossi, 2015, 2016; Lourenço & Sadine, 2014, 2015, 2016, 2017, 2021; Lourenço *et al.*, 2016, 2017, 2018a,b, 2020; Sadine *et al.*, 2016, 2024; Ythier, 2022; Ythier *et al.*, 2021, 2024). Although the core Saharan region, defined by Vachon (1952) as the 'central compartment', has been the subject of many studies, other regions in the peri-Saharan zone (surrounding most of the central compartment) also called the attention of scorpologists, such as the Saharan massifs of Tassili n'Ahaggar (Hoggar) and Tassili n'Ajjer in the South of Algeria. The patterns of distribution of North African scorpions including the role of refugia represented by the Saharan massifs for scorpion species less well adapted to xeric environments have already been discussed in several recent papers (*e.g.* Lourenço, 2001, 2002, 2010, 2013; Lourenço & Leguin, 2014; Lourenço & Rossi, 2015, 2016; Ythier, 2021; Ythier *et al.*, 2021, 2024) so it will not be further commented here.

In his contributions to the study of North African scorpions, Pallary (1928, 1929) described two *Androctonus* species from

the Saharan massifs of southern Algeria. The first one, *Androctonus eburneus* (Pallary, 1928) (= *Prionurus eburneus* Pallary, 1928), was described on the basis of two specimens collected in Djanet, in the Tassili n'Ajjer massif. Vachon (1952) raised doubts about the validity of this small yellowish species but without taking any decision on its status since he was not able to study the type material. He then placed this species in the synonymy of *A. amoreuxi* (Audouin, 1826) based of new material collected in Djanet (Vachon, 1955). Lourenço (2008) was finally able to locate the type material in the collections of the Muséum national d'Histoire naturelle of Paris, France (MNHN) and the analysis of such material revealed that this species was different from *A. amoreuxi*, so it was revalidated. It should be noted that Mousaid & Tang (2024) recently stated that “a specimen (MNHN-RS-RS2026) under the name “*Androctonus amoreuxi* Audouin & Savigny, 1825” was allegedly considered the holotype of *A. amoreuxi* from Djanet, Algeria, by the National Museum of Natural History, Paris, France”. It is obviously a mistake on the website of the MNHN, due to the fact that *A. eburneus* was previously synonymized with *A. amoreuxi* (type originally from Egypt, now lost) and this mistake will soon be corrected (E.-A. Leguin in litt.).

Androctonus hoggarensis (Pallary, 1929) (= *Prionurus hoggarensis* Pallary, 1928) was the second species described, based on specimens (number not indicated and types not designated) collected in the Hoggar massif. This large dark brown to blackish species was later

Reviewer: Gérard Dupré (France).



This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Copyright 2025. The Authors. *Faunitaxys* published by Lionel Delaunay on behalf of the AFCFF (Association française de Cartographie de la Faune et de la Flore).

also collected in the Tassili n'Ajjer massif as well as in the Air massif in northern Niger. However, Vachon (1952) called the attention to the population distributed in the Air massif, differing by a smaller size and a different coloration. New material collected in the Air massif and its comparison with specimens of *A. hoggarensis* from Hoggar and Tassili n'Ajjer massifs confirmed the existence of distinct characters allowing Lourenço (2015) to describe this population as a separate species, *Androctonus santi* Lourenço, 2015.

The study of an *Androctonus* specimen recently collected in Eferi, at the foothills of the Tassili n'Ajjer massif, Wilaya of Djanet, and its comparative analysis with other *Androctonus* species occurring in the region, has led to the description of another new species, *Androctonus ajjer* sp. n., confirming again that the Saharan Massifs are biodiversity hot spots with a very high degree of endemism within the Sahara Desert. This new species seems to show affinities with *A. amoreuxi*, distributed in Algeria in the core region of the Sahara, extending up to the Algerian Saharan massifs to the South (Fig. 13). Both species can however be easily distinguished by several morphological features. As previously mentioned, in its note synonymizing *A. eburneus* with *A. amoreuxi*, Vachon (1955) studied material of *A. amoreuxi* from Djanet. He suggested that morphological variations in *A. amoreuxi* could lead to its division in several forms (as already suggested in Vachon, 1952 and later confirmed with the description of several new species from the peri-Saharan zone, previously associated with *A. amoreuxi*). For the population of Djanet, he indicated several different morphological characters which correspond to the new species described here, notably the length of lateral inframedian carinae of metasomal segment II, shape of the scalloping of the proximal dentate margin of fixed finger in male, male pectinal tooth count, and especially a less slender metasoma with bigger spinoid granules posteriorly on segments III-IV (especially well visible on segment IV in Fig. 1 of Vachon, 1955). It therefore appears that the *Androctonus* population of Djanet examined by Vachon (1955) most certainly corresponds to the new species described here.

Material and methods

Illustrations and measurements were produced with the aid of a Motic DM143 digital stereomicroscope with an ocular micrometre. Habitus photographs were made with a Canon EOS RP and Adobe Photoshop software. Map was made using maps-for-free.com and Adobe Photoshop software. Measurements follow Stahnke (1970) and are given in mm. Trichobothrial notations follow Vachon (1974) and morphological terminology mostly follows Vachon (1952) and Hjelle (1990). Material studied herein is deposited in the collections of the MNHN (Muséum national d'Histoire naturelle), Paris, France.

Taxonomic treatment

Family **Buthidae** C. L. Koch, 1837

Genus *Androctonus* Ehrenberg, 1828

Composition of the genus *Androctonus* in Algeria (in order of description)

- *A. australis* (Linnaeus, 1758)
- *A. amoreuxi* (Audouin, 1826)
- *A. aeneas* C. L. Koch, 1839
- *A. liouvillei* (Pallary, 1924)
- *A. eburneus* (Pallary, 1928)
- *A. hoggarensis* (Pallary, 1929)
- *A. ajjer* sp. n.

Androctonus ajjer sp. n.

(Fig. 1-10)

ZooBank: <https://zoobank.org/D740E1BF-147C-4F1F-BCD3-7072EC5225A7>

Holotype, ♂, Algeria, Djanet Wilaya, Eferi, foothills of the Tassili n'Ajjer, 1010 m a.s.l., 42°29'N - 009°29'E, 09/XI/2022 (Y. Alioua). Type material deposited in the collections of the Muséum national d'Histoire naturelle, Paris, France.

Etymology. – The specific name is placed in apposition to the generic name and refers to Tassili n'Ajjer (Tassili of the Kel Ajjer Tuareg confederation), where the new species was collected.

Diagnosis. – Scorpion of large size for the genus, with a total length of 87.26 mm for the male holotype. General coloration yellowish to brownish yellow; metasomal segments IV-V and telson with dark pigmentation; legs pale yellow. Metasomal segments I-IV of approximately the same width, with a moderately to strongly marked dorsal depression; dorsal carinae strongly marked with granules becoming spinoid posteriorly; segments III-IV with one bigger posterior spinoid granule. Anal arc composed of 14 ventral teeth and four lateral lobes. Fixed and movable fingers with 12 rows of granules. Leg retrolateral basitarsal spurs with basal tooth simple. Pectines with 31-33 teeth in male holotype; fulcra with 6-7 macrosetae externally and one microseta internally.

Description based on male holotype (morphometric measurements after the description)

Coloration (in alcohol). – Basically, yellowish to brownish yellow. Prosoma: carapace yellowish orange; eyes marked with dark pigment. Mesosoma yellowish orange without spots; carinae not pigmented. Metasomal segments I-III yellowish to yellowish orange, IV-V yellowish orange with brownish (IV) to dark brown (V) pigmentation; carinae pigmented from II-V, especially ventrally; telson vesicle yellowish with dark brown pigmentation, aculeus yellowish orange at its base and blackish at its extremity. Venter yellowish; genital operculum and pectines paler than the other zones. Chelicerae yellowish without any spots; fingers yellowish with dark red teeth. Pedipalps yellowish orange; fingers with the oblique rows of granules dark red. Legs pale yellow without spots.

Morphology. – Carapace moderately granular; anterior margin almost straight, slightly concave. Carinae moderately to strongly marked; anterior median, central median and posterior median carinae strongly granular, other carinae moderately granular. All furrows weak. Median ocular tubercle located slightly anterior to the centre of the carapace; eyes separated by about three ocular diameters; five pairs of lateral eyes, the first three of moderate size, the last two reduced. Sternum triangular and narrow, longer than wide. Mesosoma: tergites moderately granular. Three longitudinal carinae moderately crenulate on tergites I to VI; tergite VII pentacarinata with carinae moderately to strongly crenulate. Venter: genital operculum divided longitudinally, each plate with an oval shape. Pectines: pectinal tooth count 31-33 in male holotype; middle basal lamella of the pectines not dilated; fulcra with 6-7 macrosetae externally, one microseta internally. Sternites III-VI with elongated spiracles, almost smooth with only some minute granules laterally; sternites III-V acarinate, VI with four vestigial carinae; sternite VII with four moderate carinae and minute granulation on whole surface except between central carinae. Metasomal segments I-IV of approximately the same width (not enlarged posteriorly), with a smooth and moderately to strongly marked dorsal depression; segment I with ten complete carinae, crenulated; segments II to IV with eight complete carinae, strongly crenulated; lateral inframedian carinae represented by 10-12 granules on distal two thirds of segment II, by 2-3 granules on III, absent on IV; dorsal carinae strongly marked with granules becoming spinoid posteriorly, segments III-IV with one bigger posterior spinoid granule; segment V with five carinae; the latero-ventral carinae strongly crenulate with several lobate denticles; ventral median carinae not divided posteriorly; anal arc composed of 14 ventral teeth and with four lateral lobes; all metasomal segments with intercarinal spaces slightly granular to smooth and a weak setation. Telson with some moderate granulations ventrally; aculeus moderately curved and of about same size as the vesicle, without a subaculear tubercle. Cheliceral dentition as defined by Vachon (1963) for the family Buthidae; external distal and internal distal teeth approximately the same length; ventral aspect of both fingers and manus covered with long dense setae. Pedipalps with a weak setation; femur



Fig. 1-2. *Androctonus ajjer* sp. n., ♂ holotype, habitus. 1. Dorsal aspect. 2. Ventral aspect.

pentacarinat; patella with 7 carinae but only dorso-internal and internal are well marked; other carinae are weak to vestigial; chela with vestigial carinae; all faces very weakly granular to smooth; fixed and movable fingers with 12 rows of granules, internal and external accessory granules present, strong; two accessory granules on the distal end of the movable finger next to the terminal denticle; scalloping of the proximal dentate margin of fixed finger in male. Legs: tarsus with numerous thin setae ventrally, arranged in more or less two rows; tibial spur moderate on legs III and IV; basitarsal spurs moderate to strong on legs I to IV; retrolateral basitarsal spur with basal tooth simple (not bifid). Trichobothriotaxy: trichobothrial pattern of Type A, orthobothriotaxic as defined by Vachon (1974). Dorsal trichobothria of femur arranged in β (beta) configuration (Vachon, 1975).

Morphometric measurements (mm) of the male holotype.

– **Total length:** 87.26.

– Carapace

length 9.50;
anterior width 5.63;
posterior width 10.63.

– **Mesosoma:** length 23.0.

– Metasomal segments

I: length 6.75; width 6.75;
II: length 7.88; width 6.88;
III: length 8.50; width 7.13;
IV: length 10.13; width 6.88;
V: length 11.0; width 6.25; depth 5.25.

– **Telson:** length 10.5.

– **Vesicle:** width 3.94; depth 3.19.

– Pedipalp

femur length 7.63; width 2.88;
patella length 8.88; width 4.38;
chela length 16.63; width 4.13; depth 4.88;
movable finger length 10.63.

Relationships. – By its general morphology, size and coloration, *Androctonus ajjer* sp. n. shows similarities with *A. amoreuxi*, described from Egypt and distributed in Algeria in the core region of the Sahara, extending up to the Algerian Saharan massifs to the south (Fig. 13).

The new species can however be readily distinguished by the following main features:

- (i) metasomal segments I-IV with a moderately to strongly marked dorsal depression (reduced in *A. amoreuxi*);
- (ii) dorsal carinae strongly marked with granules becoming spinoid posteriorly, segments III-IV with one bigger posterior spinoid granule (small rounded granules in *A. amoreuxi* and no pronounced posterior spinoid granules);
- (iii) movable finger with 12 rows of granules (13-14 in *A. amoreuxi*);
- (iv) distinct type of habitat (*A. amoreuxi* is distributed in the deserts of the core Saharan region, generally under 1000 m a.s.l.). The *Androctonus* population of Djanet examined by Vachon (1955) most certainly corresponds to the new species described here (see Introduction).

Androctonus ajjer sp. n. can also be easily distinguished from the other *Androctonus* species distributed in the Saharan massifs, notably by the following main features:

– *A. eburneus* described from Djanet:

- (i) much smaller overall size with 47.6 mm in adult male lectotype (87.26 in the new species);
- (ii) lighter general coloration, without darker pigmentation on metasoma and telson;
- (iii) higher pectinal tooth count with 38-37 teeth in male lectotype (31-33 in the new species);
- (iv) movable finger with 13 rows of granules (12 in the new species).

– *A. hoggarensis*, described from the Hoggar Massif (Tassili n'Ajger) and also distributed in the Tassili n'Ajger:

- (i) general coloration dark brown to blackish (yellowish to brownish yellow in the new species);
- (ii) fixed and movable fingers with 13 and 14-15 rows of granules, respectively (12 in the new species);
- (iii) leg retrolateral basitarsal spurs with basal tooth bifid (simple in the new species);
- (iv) distinct type of habitat (*A. hoggarensis* is a mountain species that lives at high altitude, generally between 1200-2000 m a.s.l.).




Fig. 3-9. *Androctonus ajjer* sp. n., male holotype (Scale bars: 2 mm).

3. Metasomal segments III-V and telson, lateral aspect. 4. Cutting edge of chela movable finger. 5-9. Trichobothrial pattern. 5-6. Chela, external (5) and ventral (6) aspects. 7. Femur, dorsal aspect. 8-9. Patella, dorsal (8) and external (9) aspects.



Fig. 10. *Androctonus ajjer* sp. n., male holotype alive in its natural habitat.

- *A. santi*, described from the Aïr Massif:

- (i) smaller overall size with 65-70 mm in adult males (87.26 in the new species);
- (ii) general coloration dark brown to blackish (yellowish to brownish yellow in the new species);
- (iii) distinct type of habitat (*A. santi* is a mountain species that lives at high altitude (recorded between 1200-1600 m a.s.l.) in the Aïr Massif, separated from the Hoggar and Tassili n'Ajjer massifs by arid plains) (Fig. 13).

Finally, *Androctonus ajjer* sp. n. can be distinguished from *A. australis*, distributed in Algeria between the high plateaus to the north, extending up to Ghardaïa and Ouargla to the south, notably by the following main features:

- (i) metasomal segments strongly enlarged posteriorly (approximately the same width in the new species);
- (ii) movable finger with 14 rows of granules (12 in the new species);
- (iii) leg retrolateral basitarsal spurs with basal tooth bifid (simple in the new species);
- (iv) anal arc with three lateral lobes (four in the new species);
- (v) pectines with fulcra bearing one macroseta internally (one macroseta in the new species).

Ecological characteristics of the type locality. – The region of Djanet is located on a geological fault at the far southeast of Algeria, forming a sandstone cliff that rises along the western edge of Oued Edjériou. These sandstones have preserved the traces of significant geological and climatic events that the region has experienced (orogeny, glaciation, volcanism, marine transgressions and regressions, sedimentation, fluvial and aeolian erosion) (Dubief, 1999). The region is now dominated by dune fields or ergs that crown and envelop the Tassili n'Agger foothills, which are signs of increasingly growing aridity (Beddiaf, 2020). The factor exacerbating this aridity is the rainfall, which is low (around 20mm/year) and irregular, with the months receiving the most rain being September (3.5 mm) and March (2.7 mm) (Abdoun, 2002). The temperature is relatively stable, with an average temperature of over 31.1°C, a minimum temperature of 1.1°C recorded in January, and a maximum temperature of 32.6°C recorded in August (Beddiaf, 2020). Eferi,

the type locality of *Androctonus ajjer* sp. n., is an urban area surrounded by sandblasted cliffs, interspersed with date palm orchards and other crops, contributing to the improvement of the relative humidity in this desert region (Fig. 11-12).

References

- Abidi H., Sadine S. E., Houhamdi M., Madoui A. & Lourenço W. R., 2021. – The genus *Buthus* Leach, 1815 in Algeria (Scorpiones: Buthidae) and a possible new case of vicariant species. *Revista Ibérica de Aracnología*, 38: 81-86.
- Abdoun F., 2002. – Etude de la dynamique spatio-temporelle des populations de *Cupressus dupersiana* au Tassili N'Ajjer. Thèse de Doctorat, Université de droit, d'Economie et des Sciences d'Aix- Marseille III, Faculté des Sciences et Techniques de Saint- Jérôme, Marseille, 171 p.
- Beddiaf R., 2020. – Importance de trois bio-agresseurs (*Ceratitis capitata*, *Virachola livia* et *Epilachna chrysomelina*) sur quelques cultures dans l'extrême Sud- est algérien (Djanet et Tamarrasset). Thèse de Doctorat en Sciences Agronomique. Université de Ouargla, Algérie, 150 p.
- Bouisset L. & Larrouy G., 1962. – Une nouvelle sous-espèce de *Scorpio maurus* du Nord-Ouest Oranais. *Bulletin de la Société d'Histoire Naturelle de Toulouse*, 97(3): 316-322.
- Dubief J., 1999. – L'Ajjer Sahara central, Ed. Karthala, Paris, 709 pp.
- Hjelle J. T., 1990. – Anatomy and morphology. Pp. 9-63. In: G. A. Polis (ed.), *The Biology of Scorpions*. Stanford Univ. Press, 587 p.
- Khammassi M., Harris D. J., Sadine S. E., El Bouhissi M. & Nouria S., 2023. – Description of a new species of *Scorpio* (Scorpiones: Scorpionidae) from Northwestern Algeria using morphological and molecular data. *Biologia*, 78(1): 1-12.
- Lourenço W. R., 2001. – Un nouveau genre et une nouvelle espèce de scorpion d'Algérie, avec des considérations taxonomiques sur le genre *Lissothus* Vachon, 1948 (Scorpiones, Buthidae). *Zoosystema*, 23(1): 51-57.



Fig. 11-12. Habitat of *Androctonus ajjer* sp. n. in Eferi, Djanet Wilaya, Algeria.

Lourenço W. R., 2002a. – Nouvelles considérations sur la systématique et la biogéographie du genre *Butheoloides* Hirst (Scorpiones, Buthidae) avec description d'un nouveau sous-genre et de deux nouvelles espèces. *Revue suisse de Zoologie*, 109(4): 725 - 733.

Lourenço W. R., 2002b. – Considérations sur les modèles de distribution et différenciation du genre *Buthus* Leach, 1815, avec la description d'une nouvelle espèce des montagnes du Tassili des Ajjer, Algérie (Scorpiones, Buthidae). *Biogeographica*, 78(3): 109 - 127.

Lourenço W. R., 2005. – Nouvelles considérations taxonomiques sur les espèces du genre *Androctonus* Ehrenberg, 1828 et description de deux nouvelles espèces (Scorpiones, Buthidae). *Revue suisse de Zoologie*, 112(1): 145 - 171.

Lourenço W. R., 2006. – Further considerations on the genus *Buthacus* Birula, 1908 (Scorpiones, Buthidae) with a description of one new species and two new species. *Boletín de la Sociedad Entomológica Aragonesa*, 38: 59 - 70.

Lourenço W. R., 2008. – A new species of *Androctonus* Ehrenberg, 1828 from Togo (Scorpiones, Buthidae). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 15(179), 37-44.

Lourenço W. R., 2010. – The *Compsobuthus* species from 'Tassili des Ajjer', Algeria (Scorpiones, Buthidae) and description of a new species. *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 15(182): 147 - 155.

Lourenço W. R., 2013a. – The *Buthacus* Birula, 1908 populations from Tassili n'Ajjer, Algeria (Scorpiones, Buthidae) and description of a new species Pallary (1928, 1929). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 16(190): 89 - 99.

Lourenço W. R., 2013b. – A new species of *Buthus* Leach, 1815 from Algeria (Scorpiones, Buthidae). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 16(189): 63 - 68.

Lourenço W. R., 2015. – A new species of *Androctonus* Ehrenberg, 1828 from the Aïr Massif in Niger (Scorpiones: Buthidae). *Serket*, 14(4): 167 - 175.

Lourenço W. R., 2017. – Encore une nouvelle espèce de *Buthus* Leach, 1815 (Scorpiones: Buthidae) pour le nord-est du Maroc. *Revista Ibérica de Aracnología*, 31: 59 - 63.

Lourenço W. R., Bissati S. & Sadine S. E., 2016. – One more new species of *Buthacus* Birula, 1908 from the region of Ghardaïa, Algeria (Scorpiones: Buthidae). *Arachnida - Rivista Aracnologica Italiana*, 8: 2 - 11.

Lourenço W. R., Chichi S. & Sadine S. E., 2018a. – A new species of *Buthus* Leach, 1815 from the region of Bou Sâada-M'sila, Algeria. A possible case of vicariance for the genus (Scorpiones: Buthidae). *Revista Ibérica de Aracnologia*, 32: 15-20.

Lourenço W. R., El Bouhissi M. & Sadine S. E., 2020. – Further considerations on the *Buthus* Leach, 1815 species present in Algeria with description of a new species (Scorpiones : Buthidae). *Revista Ibérica de Aracnologia*, 36: 103-108.

Lourenço W. R., Kourim M. L. & Sadine S. E., 2017. – Scorpions from the region of Tamanrasset, Algeria. Part I. A new species of *Buthacus* Birula, 1908 (Scorpiones: Buthidae). *Arachnida - Rivista Aracnologica Italiana*, 13: 31-41.

Lourenço W. R., Kourim M. L. & Sadine S. E., 2018b. – Scorpions from the region of Tamanrasset, Algeria, Part II. A new African species of the genus *Leiurus* Ehrenberg, 1828 (Scorpiones: Buthidae). *Arachnida - Rivista Aracnologica Italiana*, 16: 3-14.

Lourenço W. R. & Leguin E. A., 2011. – Further considerations on the species of the genus *Orthochirus* Karsch, 1891 from Africa, with

description of three new species (Scorpiones: Buthidae). *Euscorpium*, 123: 1-18.

Lourenço W. R. & Leguin E. A., 2014. – Une nouvelle espèce d'*Hottentotta* Birula, 1908 pour le Massif du Hoggar en Algérie (Scorpiones, Buthidae); conséquences biogéographiques sur la répartition du genre. *Revista Ibérica de Aracnologia*, 24: 15-18.

Lourenço W. R. & Rossi A., 2015. – Two new species of *Cicileus* Vachon, 1948 from Hoggar Mountains in Algeria (Scorpiones: Buthidae). *Arachnida - Rivista Aracnologica Italiana*, 4: 2-12.

Lourenço W. R. & Rossi A., 2016. – Confirmation of a new species of *Scorpio* Linnaeus, 1758 in Tassili n'Ajjer Mountains, South Algeria (Scorpiones: Scorpionidae). *Onychium*, 12: 11-18.

Lourenço W. R. & Sadine S. E., 2014. – A new species of the rare buthid scorpion genus *Lissothus* Vachon, 1948 from Central Algeria (Scorpiones, Buthidae). *Comptes Rendus Biologie*, 337: 416-422.

Lourenço W. R. & Sadine S. E., 2015. – A new species of *Buthacus* Birula, 1908 from the region of Ghardaïa, Algeria (Scorpiones, Buthidae). *Revista Ibérica de Aracnologia*, 27: 55-59.




Fig. 13. Topographic map of the three Saharan Mountain ranges of Tassili n'Ajjer, Hoggar and Aïr showing the approximate distribution of the *Androctonus* species known from this region, based on material examined and literature.

A. amoreuxi (black and white dashed area), *A. eburneus* (blue square), *A. hoggarensis* (pink circle), *A. santi* (green triangle) and *A. ajjer* sp. n. (yellow star).

- Lourenço W. R. & Sadine S. E., 2016. – One more new species of *Buthus* Leach, 1815 from Algeria (Scorpiones, Buthidae). *Revista Ibérica de Aracnologia*, 28: 13-17.
- Lourenço W. R. & Sadine S. E., 2021. – The genus *Orthochirus* Karsch, 1891, in Algeria with description of a new species (Scorpiones, Buthidae). *Bulletin de la Société Entomologique de France*, 126 (2): 175-181.
- Lourenço W. R., Sadine S. E., Bissati S. & Houtia A., 2017. – The genus *Buthacus* Birula, 1908 in Northern and Central Algeria; description of a new species and comments on possible micro-endemic populations (Scorpiones: Buthidae). *Arachnida - Rivista Aracnologica Italiana*, 12: 18-30.
- MNHN-RS-RS2026. – *Androctonus amoreuxi* Audouin & Savigny, 1826. Holotype. Muséum National d'Histoire Naturelle, Paris, France (accessed 04. II. 2025). <https://science.mnhn.fr/institution/mnhn/collection/rs/item/rs2026?listIndex=5&listCount=60>.
- Mousaid M. & Tang V., 2024. – Review of the distribution of *Androctonus amoreuxi* (Audouin, 1825) (Scorpiones: Buthidae) in Morocco. *Serket*, 20(2): 78-92.
- Pallary P., 1928. – Description de quatre scorpions nouveaux de la Berbérie. *Bulletin du Muséum National d'Histoire Naturelle, Paris*, 5: 346-351.
- Pallary P., 1929. – Les scorpions du Sahara central. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 20(6/7): 133-141.
- Sadine S. E., Bissati T. S. & Lourenço W. R., 2016. – The first true deserticolous species of *Buthus* Leach, 1815 from Algeria (Scorpiones: Buthidae); Ecological and biogeographic considerations. *Comptes Rendus Biologies*, 339(1): 44-49.
- Sadine S. E., Souilem Z., Chedad A., Chebihi B., Zebra R., Houhamdi M., Lourenço W. R. & Ythier E., 2024. – A new species of *Buthacus* Birula, 1908 from the Algerian Saharan Desert (Scorpiones: Buthidae). *Faunitaxys*, 2024, 12(9): 1-9. [https://doi.org/10.57800/faunitaxys-12\(9\)](https://doi.org/10.57800/faunitaxys-12(9))
- Stahnke H. L., 1970. – Scorpion nomenclature and mensuration. *Entomological News*, 81: 297-316.
- Vachon M., 1952. – Etudes sur les scorpions. Publications de l'Institut Pasteur d'Algérie, Alger: 482 pp.
- Vachon M., 1955. – Le Scorpion jaune des pays Ajjjer: *Androctonus amoreuxi* (Aud. et Sav., 1812 et 1826) (= *Prionurus eburneus* Pallary, 1928). *Archives de l'Institut Pasteur d'Algérie*, 33: 54-58.
- Vachon M., 1963. – De l'utilité, en systématique, d'une nomenclature des dents des chélicères chez les Scorpions. *Bulletin du Muséum national d'Histoire naturelle, Paris*, 2e sér., 35 (2): 161-166.
- Vachon M., 1974. – Etude des caractères utilisés pour classer les familles et les genres de Scorpions (Arachnides). 1. La trichobothriotaxie en arachnologie. Sigles trichobothriaux et types de trichobothriotaxie chez les Scorpions. *Bulletin du Muséum national d'Histoire naturelle, Paris*, 3e sér., n° 140, Zool. 104: 857-958.
- Vachon M., 1975. – Sur l'utilisation de la trichobothriotaxie du bras des pédipalpes des Scorpions (Arachnides) dans le classement des genres de la famille des Buthidae Simon. *Comptes Rendus des Séances de l'Académie de Sciences*, 281 (D): 1597-1599.
- Ythier E., 2021. – A new species of *Androctonus* Ehrenberg, 1828 from the Sahelian wooded steppes of Burkina Faso (Scorpiones: Buthidae). *Faunitaxys*, 9(31): 1-7. [https://doi.org/10.57800/faunitaxys-9\(31\)](https://doi.org/10.57800/faunitaxys-9(31))
- Ythier E., 2022. – A new species of *Buthacus* Birula, 1908 from Western Algeria (Scorpiones: Buthidae). *Faunitaxys*, 10(28): 1-6. [https://doi.org/10.57800/faunitaxys-10\(28\)](https://doi.org/10.57800/faunitaxys-10(28))
- Ythier E., Sadine S. E., Bengaid Y. & Lourenço W. R., 2024. – A new species of *Scorpio* Linnaeus, 1758 from Algeria (Scorpiones: Scorpionidae) and a new case of vicariance. *Arachnides*, 113: 1-10.
- Ythier E., Sadine S. E., Haddadi M. L. & Lourenço W. R., 2021. – A new species of *Buthus* Leach, 1815 from Algeria (Scorpiones: Buthidae) and an interesting new case of vicariance. *Faunitaxys*, 9(21): 1-9. [https://doi.org/10.57800/faunitaxys-9\(21\)](https://doi.org/10.57800/faunitaxys-9(21))

Authors contribution ¹

Publisher correspondence. – EY

Writing the article. – EY, SS, YA, WL

Description. – EY, SS, YA, WL

Article proofreading. – EY, SS, YA, WL

Bibliographic work. – EY, SS, YA, WL

Material study. – EY, SS, YA, WL

Author of the figures. – EY (Fig. 1-9, 13), SS (Fig. 10), YA (Fig. 11-12).

¹ EY = Ythier E. – SS = Sadine S. E. – YA = Alioua Y. – WL = Lourenço W. R.

Résumé

Ythier E., Sadine S. E., Alioua Y. & Lourenço W. R., 2025. – Une nouvelle espèce d'*Androctonus* Ehrenberg, 1828 du Tassili n'Ajjer en Algérie (Scorpiones: Buthidae). *Faunitaxys*, 13(08): 1-8.

Une nouvelle espèce appartenant au genre *Androctonus* Ehrenberg, 1828 est décrite sur la base d'un spécimen mâle collecté à Eferi, au pied du massif du Tassili n'Ajjer, Wilaya de Djanet, Algérie. Ce nouveau taxon représente la 42^{ème} espèce d'*Androctonus* décrite à ce jour et la septième connue pour l'Algérie. Une carte de la distribution des espèces d'*Androctonus* présentes dans la région des massifs montagneux du Sahara d'Algérie et du Niger est présentée.

Mots-clés. – Scorpion, Buthidae, *Androctonus*, *ajjer*, taxonomie, nouvelle espèce, description, morphologie, massifs Sahariens, Tassili n'Ajjer, Algérie.

Derniers articles publiés

- Hui Wen, Xin-Yuan Zhang, Shu-Tong Zhang & Hao Xu, 2024. – Distribution of the flower beetle *Pseudodicerus nigrocyaneus* (Bourgoin, 1917) (Coleoptera: Scarabaeidae: Cetoniinae) in Asia with biological notes. *Faunitaxys*, 12(57): 1 – 15. [https://doi.org/10.57800/faunitaxys-12\(57\)](https://doi.org/10.57800/faunitaxys-12(57))
- Cumming R. T., Engel M. S., Lian Z. & Ulitzka M. R., 2024. – Small, intricate, and beautiful; a new species of lophioneurid from the Cretaceous (Insecta: Thripida: †Lophioneurida). *Faunitaxys*, 12(58): 1 – 10. [https://doi.org/10.57800/faunitaxys-12\(58\)](https://doi.org/10.57800/faunitaxys-12(58))
- Ythier E. & Lourenço W. R., 2024. – A new species of *Hadrurochactas* Pocock, 1893 (Scorpiones: Chactidae) from the Mitaraka Massif in French Guiana. *Faunitaxys*, 12(59): 1 – 9. [https://doi.org/10.57800/faunitaxys-12\(59\)](https://doi.org/10.57800/faunitaxys-12(59))
- Háva J., 2024. – *Mycetophagus* (s. str.) *lenkae* sp. nov. from India: Arunachal Pradesh (Coleoptera: Mycetophagidae), along with new records for *Mycetophagus* (*Mycetophagoides*) *alni* Champion in two states in India. *Faunitaxys*, 12(60): 1 – 3. [https://doi.org/10.57800/faunitaxys-12\(60\)](https://doi.org/10.57800/faunitaxys-12(60))
- Herbin D., Giusti A. & Mielke C. G. C., 2024. – Taxonomic revision of the genera *Tamphana* Schaus and *Tuyvae* Orlandin, Piovesan & Carneiro with description of eleven new species (Lepidoptera, Apatelodidae). *Faunitaxys*, 12(61): 1 – 31. [https://doi.org/10.57800/faunitaxys-12\(61\)](https://doi.org/10.57800/faunitaxys-12(61))
- Niisato T. & Tichý T., 2024. – A new species of the genus *Microdebilissa* Pic, 1925 from West Kalimantan (Coleoptera, Cerambycidae, Cerambycinae, Stenopterini). *Faunitaxys*, 12(62): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(62\)](https://doi.org/10.57800/faunitaxys-12(62))
- Moret P., 2024. – Description of new Platynini from the montane cloud forest of Ecuador, with a redefinition of the genera *Glyptolenus* Bates and *Glyptolenoides* Perrault (Coleoptera, Carabidae). *Faunitaxys*, 12(63): 1 – 23. [https://doi.org/10.57800/faunitaxys-12\(63\)](https://doi.org/10.57800/faunitaxys-12(63))
- Cerda J.-A., 2024. – Corrigenda de *Butlerius* et *Guerinius* (Lepidoptera, Noctuoidea, Erebidae, Arctiinae, Arctiini, Ctenuchina). Quinzième note. *Faunitaxys*, 12(64): 1. [https://doi.org/10.57800/faunitaxys-12\(64\)](https://doi.org/10.57800/faunitaxys-12(64))
- Vives E., 2024. – Un nuevo género de Strogylurini de las islas Filipinas. New or interesting Cerambycidae from Philippines (Col.: Cerambycidae, Cerambycinae) (Pars 31). *Faunitaxys*, 12(65): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(65\)](https://doi.org/10.57800/faunitaxys-12(65))
- Vlasak J., Santos-Silva A. & Bezark L. G., 2024. – South American Acanthocini (Coleoptera, Cerambycidae, Lamiinae): description of a new species; transfer of a species from *Urgleptes* to *Lepturges* (*Lepturges*); designation of a lectotype; and corrections. *Faunitaxys*, 12(66): 1 – 11. [https://doi.org/10.57800/faunitaxys-12\(66\)](https://doi.org/10.57800/faunitaxys-12(66))
- Bollino M. & Racheli T., 2025. – A new taxon of the *Pterourus menatius* complex from West Ecuador (Lepidoptera: Papilionidae). *Faunitaxys*, 13(01): 1 – 6. [https://doi.org/10.57800/faunitaxys-13\(01\)](https://doi.org/10.57800/faunitaxys-13(01))
- Lourenço W. R. & Velten J., 2025. – One more new species for the genus *Cretaceoushormiops* Lourenço, 2018 from Cretaceous Burmite (Scorpiones: Protoischnuridae). *Faunitaxys*, 13(02): 1 – 6. [https://doi.org/10.57800/faunitaxys-13\(02\)](https://doi.org/10.57800/faunitaxys-13(02))
- Lourenço W. R. & Velten J., 2025. – A new species for the genus *Archaeoananteroides* Lourenço, 2016, fossil scorpion from Burmite (Scorpiones: Buthoidea: Buthidae). *Faunitaxys*, 13(03): 1 – 6. [https://doi.org/10.57800/faunitaxys-13\(03\)](https://doi.org/10.57800/faunitaxys-13(03))
- Lourenço W. R. & Ythier E., 2025. – A new cave species of *Isometrus* Ehrenberg 1828 from Myanmar (Scorpiones: Buthidae). *Faunitaxys*, 13(04): 1 – 6. [https://doi.org/10.57800/faunitaxys-13\(04\)](https://doi.org/10.57800/faunitaxys-13(04))
- Platia G. & Pedroni G., 2025. – Five new species and new records of click beetles from Saudi Arabia with an updated catalogue (Coleoptera, Elateridae). *Faunitaxys*, 13(05): 1 – 12. [https://doi.org/10.57800/faunitaxys-13\(05\)](https://doi.org/10.57800/faunitaxys-13(05))
- Háva J. & Hodeček J., 2025. – Two additional new species of *Cryptorhopalum* (Coleoptera: Dermestidae: Megatominae) from the Dominican Republic. *Faunitaxys*, 13(06): 1 – 5. [https://doi.org/10.57800/faunitaxys-13\(06\)](https://doi.org/10.57800/faunitaxys-13(06))
- Vitali F., 2025. – The genus *Epicasta* Thomson, 1864 and the limits of the morphological systematics in the study of Lamiinae (Coleoptera Cerambycidae). *Faunitaxys*, 13(07): 1 – 5. [https://doi.org/10.57800/faunitaxys-13\(07\)](https://doi.org/10.57800/faunitaxys-13(07))

Faunitaxys est échangée avec les revues suivantes (« print versions ») :

- Annali del Museo Civico di Storia Naturale G. Doria (Italie)
- Boletín de la Asociación española de Entomología (Espagne)
- Boletín de la Sociedad Andaluza de Entomología (Espagne)
- Bollettino del Museo di Storia Naturale di Venezia (Italie)
- Bulletin de la Société linnéenne de Lyon (France)
- Bulletin of Insectology (Italie)
- Heteropterus Rev. Entomol. (Espagne)
- Integrative Systematics (Allemagne)
- Israel Journal of Entomology (Israël)
- Klapalekiana (République Tchèque)
- Koleopterologische Rundschau (Allemagne)
- Memorie del Museo Civico di Storia Naturale di Verona (Italie)
- Nova Supplementa Entomologica (Allemagne)
- Proceedings of the Entomological Society of Washington (USA)
- Revue suisse de Zoologie (Suisse)
- Spixiana (Allemagne)
- Zoosystematica Rossica (Russie)

Faunitaxys

Volume 13, Numéro 08, Février 2025

SOMMAIRE

Une nouvelle espèce d'*Androctonus* Ehrenberg, 1828 du Tassili n'Ajjer en Algérie (Scorpiones: Buthidae).

Eric Ythier, Salah Eddine Sadine, Youcef Alioua & Wilson R. Lourenço 1 – 8

CONTENTS

A new species of *Androctonus* Ehrenberg, 1828 from the Tassili n'Ajjer, Algeria (Scorpiones: Buthidae).

Eric Ythier, Salah Eddine Sadine, Youcef Alioua & Wilson R. Lourenço 1 – 8



Illustration de la couverture :

Tassili n'Ajjer landscape (photo: Djaren-Sefar).

Crédits:

Eric Ythier : Fig. 1-9 & 13.

Salah Eddine Sadine : Fig. 10.

Youcef Alioua : Fig. 11-12.

Djaren-Sefar / Pixabay : couverture.