

**Table 1. Characteristics of bariatric surgery patients with ESRD at baseline (before BS) and during follow-up**

	<b>Before BS n=22</b>	<b>1 year n=22</b>	<b>5 years n=13</b>
Sex (% female)	12 (54.5%)		
Age (years)	47.1 (10.1)		
Type 2 diabetes	12 (54.5%)		
Hypertension	18 (81.8%)		
Treated obstructive sleep apnea	13 (59.1%)		
Weight (kg)	111.9 (14.3)	87.7 (13.3)	89.4 (14.7)
BMI (kg/m <sup>2</sup> )	40.5 (4.6)	31.7 (4.8)	31.7 (5)
Fat mass (kg)	48.8 (10.2)	35.2 (8.6)	33.9 (7.8)
Fat mass (%)	43.3 (7.3)	39.4 (8)	39.7 (9.3)
Lean mass (kg)	60.4 (11.6)	52.2 (10.9)	54 (12.1)
Lean mass (%)	53.9 (7.3)	58.3 (8.1)	58.8 (9.7)
Appendicular lean mass/ total weight (%)			
Male	26.4 (4.8)	28.2 (5.9)	27.5 (3.8)
Female	21.7 (2.6)	23.2 (4.3)	23.7 (3.4)
Sarcopenic obesity (n, %)*	8 (36.3)	4 (18)	2 (15.3)
Weight-loss (%)**		21.4 (8.5)	23 (12)
Lean mass-loss (%)***		13.5 (5.9)	15.3 (9)
Lean mass-loss/weight-loss (%)		36.7 (19.7)	47 (25.3)

Continuous variables are expressed as mean (SD), categorical variables are expressed as n (%); BMI: Body Mass Index

\* according to 2022 EASO-ESPEN consensus, based solely on DXA measurements

\*\*Weight at 1 or 5 years – preoperative weight

\*\*\*Lean mass at 1 or 5 years – preoperative weight