



HAL
open science

Macroecological rules predict how biomass scales with species richness in nature

Alex Pigot, Laura Dee, Anthony Richardson, Declan Cooper, Nico Eisenhauer, Richard Gregory, Simon Lewis, Callum Macgregor, Dario Massimino, Daniel Maynard, et al.

► To cite this version:

Alex Pigot, Laura Dee, Anthony Richardson, Declan Cooper, Nico Eisenhauer, et al.. Macroecological rules predict how biomass scales with species richness in nature. *Science*, 2025, 387 (6740), pp.1272-1276. <10.1126/science.adq3278>. <hal-05029605>

HAL Id: hal-05029605

<https://hal.science/hal-05029605v1>

Submitted on 10 Apr 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Macroecological rules predict how biomass scales with species richness in nature

Alex L. Pigot^{1*}, Laura Dee², Anthony J. Richardson^{3,4}, Declan L. M. Cooper^{1,5}, Nico Eisenhauer^{6,7}, Richard D. Gregory^{1,8}, Simon L. Lewis^{5,9}, Callum J. Macgregor¹⁰, Dario Massimino¹¹, Daniel Maynard¹, Helen R. P. Phillips^{12, 13}, Marina Rillo¹⁴, Michel Loreau^{15,16}, Bart Haegeman¹⁷.

¹Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, Gower Street, London, WC1E 6BT, UK

²Laura E. Dee, Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, USA

³School of Environment, University of Queensland, St Lucia, 4072, Queensland, Australia

⁴CSIRO Environment, Queensland Biosciences Precinct, St Lucia, 4067, Queensland, Australia

⁵Department of Geography, University College London, Gower Street, London, WC1E 6BT, UK

⁶German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁷Leipzig University, Institute of Biology, Leipzig, Germany

⁸RSPB Centre for Conservation Science, Sandy, Bedfordshire SG19 2DL, UK

⁹School of Geography, University of Leeds, Leeds, UK

¹⁰British Trust for Ornithology (BTO) Cymru, Thoday Building, Deiniol Road, Bangor, Gwynedd, LL57 2UW, UK

¹¹British Trust for Ornithology, Thetford, UK

¹²Netherlands Institute of Ecology (NIOO-KNAW), Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, 6708 PB Wageningen, The Netherlands

¹³Saint Mary's University, Department of Environmental Sciences, Halifax, Nova Scotia, Canada

¹⁴Institute for Chemistry and Biology of the Marine Environment (ICBM), University of Oldenburg, Wilhelmshaven, Germany

¹⁵Theoretical and Experimental Ecology Station, CNRS, 2 route du CNRS, 09200 Moulis, France

¹⁶Institute of Ecology, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

¹⁷CNRS/Sorbonne Université, Laboratory of Microbial Oceanography, Av. Pierre Fabre, 66650 Banyuls-sur-Mer, France

*Corresponding author. Email: a.pigot@ucl.ac.uk

Despite advances in theory and experiments, how biodiversity influences the structure and functioning of natural ecosystems remains debated. By applying new theory to data on 84,695 plant, animal and protist assemblages we show that the general positive effect of species richness on stocks of biomass, as well as substantial variation in the strength and sign of this effect, is predicted by a fundamental macroecological quantity: the scaling of species abundance with body mass. Standing biomass increases with richness when large-bodied species are numerically rare but is independent of richness when species size and abundance are uncoupled. These results suggest a new fundamental law in the structure of ecological communities and show that the impacts of changes in species richness on biomass are predictable.

Introduction

Human activities are rapidly eroding global biodiversity, creating a pressing need to understand the consequences of these losses for the structure and function of Earth's ecosystems (1-3). Hundreds of experimental studies based on artificial species assemblages have provided critical evidence that biodiversity supports higher standing stocks of biomass and fluxes of energy through ecosystems (4), and theory identifies several underlying mechanisms for these effects (5-7). But understanding the relationship between biodiversity and the structure and functioning of assemblages in nature, with their greater scale and complexity, remains a major challenge (8). On the one hand, many empirical studies of animal and plant assemblages have reported positive associations between species richness and biomass that are consistent with, or stronger than, effects estimated from experiments (9-11). However, in other cases, the biomass supported by an ecosystem appears to be decoupled from changes in species richness, or even declines at higher levels of biodiversity (11-13). This heterogeneity has led to the suggestion that the effect of species richness on ecosystem function in nature is likely to be idiosyncratic and context dependent, eluding a general unifying theory (14). The problem is therefore to explain not only the widespread empirical trend for biomass stocks to increase with biodiversity, but also the substantial variation in the strength and sign of this relationship (15). In contrast to artificial species assemblages, which typically consist of random mixtures of species, natural assemblages exhibit a number of general regularities, or 'macroecological rules', that describe how the abundance of individuals and their traits—such as body size—are distributed across species and space (16, 17). One potentially promising approach for understanding the effect of biodiversity on ecosystem functioning is therefore to consider the patterns that arise from the macroecological rules governing how assemblages in nature are structured (18).

Here we develop and test a macroecological model to predict the effect of species richness on standing biomass. We acknowledge that biomass represents a simplistic proxy that does not capture the full multi-dimensional nature of ecosystem functioning (19), but we use it here as it provides a universal and widely measured property of ecosystems linked to a variety of critical functions and services (13), including carbon storage (20), primary production (21) and trophic energy flows (22). Our model is designed to make minimal assumptions while capturing the key macroecological rules that describe how assemblages in nature are structured and sampled. These rules should be applicable to any kind of organism or environment (23) and thus we assess the predictions of this model using data on the richness and standing biomass of a diverse compilation of aquatic and terrestrial assemblages spanning plants, animals and protists (Table S1). Our analysis reveals how the general positive effect of species richness on stocks of biomass

observed across taxa arises from a well-known but previously unlinked macroecological rule: the trend for large bodied species to be numerically rare.

A macroecological model of biomass scaling with species richness

To understand the link between biodiversity effects, species abundance and body size, consider a model consisting of a regional pool of species that vary in their abundance (A) and body mass (M) and where local assemblages represent random samples of individuals drawn from the environment (Fig. 1) (24). We assume that variation in species body masses and abundances both follow the widely supported lognormal distribution (17, 25, 26) and that, as is common in nature, A and M are negatively related (27, 28) and follow a power-law relationship with exponent λ . That is, when plotted on a log-log scale this relationship is linear with slope λ .

$$\log(A) = \text{constant} + \lambda \times \log(M)$$

Such a negative effect of M on A is often attributed to energetic or spatial constraints on population carrying capacity, as larger organisms that consume more resources and occur at higher trophic levels can be supported at lower densities (28-30). Differences in richness and biomass across assemblages arise in our model due to stochastic variation in the sampled abundance of each species. Because we are interested in predicting the relative change in biomass given a relative change in species richness, we focus on predicting the slope (θ) of the least squares regression of log-log transformed biomass (B) on richness (R).

$$\log(B) = \text{constant} + \theta \times \log(R)$$

Our use of power laws to estimate λ and θ should not be taken to imply we are assuming a particular mechanistic basis to these relationships, or that these are necessarily the best fitting models. Rather we follow previous broad-scale analyses (e.g., (31)) in using these descriptors to allow comparison across organisms groups where body mass is measured on very different scales.

A positive relationship between species richness and biomass can arise through two distinct routes with potentially different implications for ecosystem functioning (32): species rich assemblages could (i) contain a greater number of individuals and/or (ii) contain individuals which are on average larger. Because the effect of the number of individuals is relatively straightforward, we focus on the average size effect, by examining the relationship between species richness and biomass across samples of a fixed total abundance. We later relax this assumption to explore the joint effect of the number of individuals and average size.

Our simple theoretical model reveals that a positive effect of species richness on standing biomass ($\theta > 0$) arises due to the general macroecological rule for abundance to decline with body mass ($\lambda < 0$) (Fig. 1). First, note that when a constant number of individuals is repeatedly sampled at random from the species pool, variation in richness and biomass is mathematically constrained. Assemblages that, by chance, contain only a single species can have anything from a very low to very high total biomass, depending on whether the individuals come from the smallest- or largest-bodied species respectively. In contrast, a species-rich assemblage must contain both small- and large-bodied species (because there are a limited number of small-bodied and large-bodied species, respectively, to sample from) and is thus constrained to have an intermediate biomass. Second, note that while this mathematical constraint defines the space of

possible richness and biomass values, assemblages are unlikely to occupy this space evenly. This is because when individuals are randomly sampled from the environment, small-bodied species that occur at a higher abundance are more likely to be included in the sample. Thus, for an assemblage where, by chance, all individuals that were sampled happened to belong to a single species, it is much more likely to be a common small-bodied species than a rare large-bodied species. This dominance of small individuals will result in this species-poor assemblage having a relatively low standing biomass. Equally, the rarity of large-bodied species makes them less likely to be sampled, and these rare species will tend to only be represented in those assemblages that by chance have a higher species richness. The presence of these rare but large-bodied species ensures a relatively high standing biomass in species-rich assemblages compared to assemblages comprising fewer species. Thus, when large-bodied species are numerically rare ($\lambda < 0$), the random sampling of individuals alone can generate a positive relationship between species richness and biomass ($\theta > 0$).




Fig. 1. How size-dependent constraints on species abundance determine the effect of richness on biomass stocks. (A) For simplicity here we show a species pool consisting of 5 hypothetical species represented by circles of different colors, with circle size denoting mean species body mass. (B) Species abundance varies inversely with body mass (here, $\lambda = -1$). (C) A fixed number of individuals (here $n = 5$) are repeatedly drawn at random from the species pool to form local assemblages, which exhibit stochastic variation in both richness and biomass. While not evident in this simplified example with only five individuals, local assemblages exhibit species body mass-abundance relationships mirroring the regional pool (fig. S1). (D) The two-dimensional space of all possible richness and biomass states from our model are constrained (dashed black lines) by the body masses available in the species pool. Cells indicate the probability (dark shading indicates higher probability) of each richness-biomass combination under a random draw of individuals. Solid line shows the expected effect of species richness on

biomass (θ) estimated via least squares regression. For each level of species richness, assemblages with the highest and lowest possible biomass are shown. Numerals correspond to the assemblages shown in (C). For simplicity in this example, intraspecific size variation is ignored, but the dependency of θ on λ is robust to intraspecific size variability (24).

While it has been argued that the species body mass-abundance relationship has a canonical form of $\lambda = -3/4$ (26), empirical studies show that λ varies across taxa and contexts from strongly negative to absent or even weakly positive (27). We therefore next examined the effects of variation in λ on the expected slope of the richness biomass relationship. Throughout, we use λ estimated from the abundance of species recorded across local assemblages matching the data typically available in empirical studies, but note that similar results are obtained when using the body mass-abundance relationship of the regional pool (fig. S1). Our macroecological model shows that variation in λ is expected to give rise to predictable variation in the sign and strength of θ (Fig. 2A). Specifically, when body mass is negatively correlated with abundance ($\lambda < 0$), then as illustrated in Fig. 1, the expected effect of richness on biomass is positive ($\theta > 0$). In contrast, when variation in species body mass and abundance are independent ($\lambda = 0$), biomass is expected to vary independently of species richness ($\theta = 0$). And, when the typical relationship between species body mass and abundance is reversed—so that numerically abundant species have a larger body mass ($\lambda > 0$)—the expected effect of richness on biomass is also reversed, with biomass declining with increasing richness ($\theta < 0$).

While λ acts as a master variable controlling the sign of θ (Figs. 2A), our simulations show that it explains only limited variation in the magnitude of θ , as this also depends on the variance in log-transformed M ($\sigma_{\log(M)}^2$) and A ($\sigma_{\log(A)}^2$) (Fig. 2A). When there is relatively little variation in body mass among species, then regardless of how these species are mixed within assemblages, total biomass will vary little. When there is relatively high variation in abundance among species, most assemblages will comprise the same highly abundant species, again resulting in little variation in biomass. Thus, for a given λ , the expected magnitude of θ can vary substantially (Fig. 2A). Rescaling λ by the relative variance in log-transformed M ($\sigma_{\log(M)}^2$) and A ($\sigma_{\log(A)}^2$) provides a metric, λ_S , that accounts for these effects.

$$\lambda_S = \lambda \times \frac{\sigma_{\log(M)}^2}{\sigma_{\log(A)}^2}$$

Having done this rescaling, the unexplained variation in θ largely collapses, with λ_S reliably predicting both the expected sign and steepness of θ (Fig. 2B). An exact analytical link between θ and λ_S is lacking and the exact form of the relationship depends on the precise parameters of the model, such as the number of species in the regional pool (figs. S2-S3). However, sensitivity analyses show that under a range of realistic scenarios, λ_S could serve as a heuristic metric to predict how changes in species richness influence the biomass of ecological assemblages (24). For example, the strong theoretical dependency of θ on λ_S is robust to relaxing key assumptions of our model, including the shape of the species abundance distribution (fig. S3), the presence of intra-specific size variation (fig. S4) and non-random sampling arising due to the spatial aggregation of species populations (figs. S5-S6). This theoretical dependency is also not an artefact of sampling assemblages consisting of a fixed number of individuals, because we

observed the same effect when allowing assemblages to contain different numbers of individuals (Fig. 2c). In this case, the expected richness-biomass relationship is always positive because assemblages containing more individuals will tend to sample more species and contain a higher total biomass (fig. S7), with this effect becoming stronger with increasing variation in the number of individuals across assemblages (Fig. 2c). Taken together, these results confirm the strong theoretical dependency of the richness-biomass relationship on the number of individuals, while revealing λ_S an additional macroecological rule, whereby differences in $\sigma_{log(M)}$ modulate the richness-biomass relationship by altering the average size of the species in an assemblage.




Fig. 2. Theoretical expectation of how differences in size-dependent constraints on species abundance drive heterogeneity in the sign and steepness of the species richness-biomass relationship (θ). (A) The slope λ of the relationship between species body-mass (M) abundance (A) determines the expected sign of θ . For a given λ , the steepness of θ varies widely depending on the relative variance in species body mass ($\sigma_{log(M)}^2$) and species abundance ($\sigma_{log(A)}^2$), indicated by different colors. (B) When accounting for the variance in body mass and abundance, the rescaled species body-mass abundance relationship λ_S strongly predicts both the sign and steepness of θ . Each point shows the mean θ across 500 replicate simulations. (C) the dependency of θ on λ_S when the total number of individuals varies across assemblages, showing scenarios from a 2-fold to 20-fold variation in total abundance. In (A-C) λ and λ_S are estimated from species abundance across local assemblages to match the empirical data typically available.

Biomass scaling with species richness across taxa in nature

To test this theoretical prediction, we collated a database of assemblages comprising records of species abundance across multiple survey sites and for which we could obtain estimates of individual or species average mass (24). In total, our database includes 84,695 assemblages, consisting of ~60 million individual organisms and 4,092 species. Those species come from 11 major taxonomic groups with vastly different ecologies, from earthworms to elephants and from unicellular nanoplankton weighing one trillionth of a gram to trees weighing 200 tonnes (Table S1). The empirical relationship between richness and biomass could be sensitive to difference in the area or volume (i.e. grain size) of survey sites, as larger grains will encompass more habitats and individuals (33). While grain size necessarily varies substantially across organism groups,

we selected datasets such that within each group assemblages were generally sampled using a standardised grain size. Nevertheless, the number of sampled individuals often varied substantially across assemblages driving consistently strong positive richness-biomass relationships as predicted by our theoretical model (fig. S8, table S2). Variation in the number of individuals could reflect a causal effect of biodiversity on organism density (32). However, it could also reflect the stochastic nature of sampling (24) or environmental (e.g., energetic (34)) constraints on carrying capacity, both of which could drive the appearance of a positive effect of richness on biomass. To isolate the effect of richness on biomass independent of total assemblage abundance, we therefore rarefy assemblages to a constant number of individuals before estimating biomass, richness and θ (24).

Across most taxa, stocks of biomass increase with species richness, supporting the widely reported positive effect of biodiversity on biomass (11). However, θ varies substantially across taxa (Fig. 3A): it is significantly positive for six of the eleven taxonomic groups, is not significantly different from zero for three (mammals, earthworms and coccolithophores) and for two others is significantly negative (moths and birds) (Table S3). This is consistent with the heterogeneity in the effects of biodiversity on biomass previously reported for natural assemblages of animals and plants (11-13). However, in accordance with our theoretical model, much of the variation in the sign and steepness of θ across taxa follows predictably from variation in the sign and magnitude of λ_S estimated from local assemblage data (Fig. 3C). For example, fish exhibit the steepest negative λ_S of -0.29 and accordingly also exhibit the steepest increase in biomass with species richness. In contrast, λ_S is weakly positive in our dataset of UK breeding birds ($\lambda_S = 0.04$), and in this group biomass declines with species richness. Other taxa have intermediate values of λ_S between these extremes, with the effect of species richness on biomass varying as predicted i.e. becoming more strongly positive as the effect of body size on abundance becomes more strongly negative (Table S3).

Overall, a linear model including λ_S alone is able to explain 60% of the variation in θ across taxa (intercept = -0.02, slope = -2.6, $p = 0.003$ (24)). This strong dependency and predictive power is robust to uncertainty in estimates of λ_S and θ , including resulting from how assemblages are subsampled (fig. S9-10) and whether they are standardised to a constant abundance or level of sampling completeness (fig. S11). Variation among taxa in θ is also predicted by the unscaled slope λ (intercept = 0.08, slope = -0.88, $p = 0.03$, $r^2 = 36\%$), but in accordance with our theoretical model, this dependency is weaker than the dependency on λ_S (Fig. 3B). Examples from particular taxa highlight the reason for this. For instance, species abundance in Dinoflagellates declines strongly with size ($\lambda = -0.55$), but because there is relatively little variation in body mass among species, θ is weaker than expected from λ alone. In contrast, fish exhibit relatively large variation in body mass and this, combined with the strong negative scaling of abundance with size ($\lambda = -0.47$), results in a particularly steep increase in biomass with species richness. Richness-biomass relationships are thus most steeply positive in those taxa that exhibit both strong size-dependent constraints on abundance and substantial variation in body mass across species.




Fig. 3. Size-dependent constraints on species abundance predict heterogeneity in the slope of the species richness-biomass relationship (θ) across taxa. (A) The observed relationship for each taxonomic group ($n = 11$) between log-log transformed species richness and biomass and log-log transformed species body mass and abundance. Lines are estimated regression slopes (Table S3). Taxa are ordered according to the rescaled species body-mass abundance relationship λ_S , from steeply negative (left) to positive (right). (B, C) Covariation between θ and (B) λ and (C) λ_S across taxa. Colors indicate the taxonomic group in (A). Bars show 95% confidence intervals (not visible for some taxa). In (B, C) the fitted line is the ordinary least square regression slope (\pm S.E) between λ_S the rescaled (or λ , the unscaled) slope of the relationship between species body-mass and abundance and θ the slope of the species richness-biomass relationship. See (24) for sources of taxon images.

Conclusions

Together, our theoretical and empirical findings provide evidence that the general positive effect of species richness on the standing biomass of natural assemblages arises as a mirror image of the general negative effect of body mass on species abundance. Furthermore, far from being idiosyncratic, we show that differences among taxa and environments in the sensitivity of biomass to changes in species richness arise predictably from differences in the scaling of abundance with body size. For example, biomass increases more strongly with species richness in aquatic compared to terrestrial groups (31), as expected from the stronger negative scaling of abundance with body size in aquatic ecosystems (35). While in this work we have focussed on

comparing organism groups with very different size structures, human activities are driving profound changes in the distribution of species size and abundance (36, 37). Our model could provide a basis for predicting the consequences of these changes in size-structure for how biodiversity influences ecosystem function.

Our results should not be taken to suggest a simple bivariate relationship between richness and biomass in nature, and its dependency on the species body mass-abundance slope. As we show, uncovering these relationships requires first controlling for differences in the relative extent of variation in body mass across species (Fig. 2B) and for differences across assemblages in the number of individuals (Fig. 2C). An important next step will be to integrate our findings into more general theoretical frameworks that aim to predict these additional state variables that are critical for a comprehensive understanding of how biodiversity and ecosystem function are connected (18, 34). Further work is also required to understand how different regional pool structures, environmental contexts, non-random patterns of sampling and community assembly processes—including interactions between species—alter our theoretical predictions and ability to explain the empirical patterns.

The processes shaping ecosystem functioning are undoubtedly more complex than those considered in our theoretical model. For example, the key ecological mechanisms of niche complementarity and facilitation that experiments have shown promote higher biomass in more species rich systems (6) are not explicitly considered. This should not be taken to imply that such mechanisms are unimportant in explaining patterns of ecosystem functioning in nature. Rather, our results suggest that knowledge of how these mechanisms operate may not be required to predict the impact of changes in species richness on biomass (38). For example, a key constraint identified in our theoretical model that drives increases in biomass with species richness, is for numerically rare (large bodied) species to preferentially occur in more species rich assemblages (Fig. 1). In our model this arises due to random sampling, but the same effect could be driven by other niche-based mechanisms that cause numerically rare species to congregate in the richest assemblages (e.g. lower extinction rates) (34). Our analysis is unable to resolve what ultimately drives the distribution of rarity across species assemblages and body size classes (27), but the quest to identify the causes of these macroecological rules (39) emerges from our analysis as a priority for understanding the role of biodiversity in the functioning of ecosystems.

Here we identify a previously unappreciated fundamental law in the scaling of biomass with species richness that appears to hold across contrasting physical environments and kingdoms of life. Because metabolic rates and fluxes of energy and matter consistently scale with body mass (40), and because abundance covaries with a number of other species traits (41), this law could encompass similar generalities in the relationship between other facets of biodiversity (e.g. functional trait diversity) and metrics of ecosystem function (e.g. productivity or stability). Such a prospect points towards a unification of macroecology and ecosystem science, in which changes in the structure and functioning of ecosystems can be predicted from the natural rules governing the abundance and distribution of individual species.

References

1. B. J. Cardinale *et al.*, Biodiversity loss and its impact on humanity. *Nature* **486**, 59-67 (2012).
2. IPBES, "Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services," (Bonn, Germany, 2019).
3. F. Isbell *et al.*, Linking the influence and dependence of people on biodiversity across scales. *Nature* **546**, 65-72 (2017).
4. A. Hector *et al.*, Plant diversity and productivity experiments in European grasslands. *Science* **286**, 1123-1127 (1999).
5. M. Loreau, Biodiversity and ecosystem functioning: A mechanistic model. *P Natl Acad Sci USA* **95**, 5632-5636 (1998).
6. M. Loreau, A. Hector, Partitioning selection and complementarity in biodiversity experiments (vol 412, pg 72, 2001). *Nature* **413**, 548-548 (2001).
7. D. Tilman, C. L. Lehman, K. T. Thomson, Plant diversity and ecosystem productivity: Theoretical considerations. *P Natl Acad Sci USA* **94**, 1857-1861 (1997).
8. A. Gonzalez *et al.*, Scaling-up biodiversity-ecosystem functioning research. *Ecol Lett* **23**, 757-776 (2020).
9. R. Danovaro *et al.*, Exponential decline of deep-sea ecosystem functioning linked to benthic biodiversity loss. *Curr Biol* **18**, 1-8 (2008).
10. C. Mora *et al.*, Global Human Footprint on the Linkage between Biodiversity and Ecosystem Functioning in Reef Fishes. *Plos Biol* **9**, (2011).
11. J. E. Duffy, C. M. Godwin, B. J. Cardinale, Biodiversity effects in the wild are common and as strong as key drivers of productivity. *Nature* **549**, 261-+ (2017).
12. L. E. Dee *et al.*, Clarifying the effect of biodiversity on productivity in natural ecosystems with longitudinal data and methods for causal inference (vol 14, 2607, 2023). *Nat Commun* **14**, (2023).
13. F. van der Plas, Biodiversity and ecosystem functioning in naturally assembled communities. *Biol Rev* **94**, 1220-1245 (2019).
14. N. Eisenhauer *et al.*, The heterogeneity-diversity-system performance nexus. *Natl Sci Rev* **10**, (2023).
15. N. Eisenhauer *et al.*, A multitrophic perspective on biodiversity-ecosystem functioning research. *Adv Ecol Res* **61**, 1-54 (2019).
16. J. H. Brown, B. A. Maurer, Macroecology - the Division of Food and Space among Species on Continents. *Science* **243**, 1145-1150 (1989).
17. K. J. Gaston, T. M. Blackburn, *Pattern and process in macroecology*. (Blackwell Science, Oxford, OX ; Malden, MA, USA, 2000), pp. xii, 377 p.
18. J. Harte, M. Brush, E. A. Newman, K. Umemura, An equation of state unifies diversity, productivity, abundance and biomass. *Commun Biol* **5**, (2022).
19. A. Hector, R. Bagchi, Biodiversity and ecosystem multifunctionality. *Nature* **448**, 188-190 (2007).
20. S. Fauset *et al.*, Hyperdominance in Amazonian forest carbon cycling. *Nat Commun* **6**, (2015).
21. D. G. Boyce, M. R. Lewis, B. Worm, Global phytoplankton decline over the past century. *Nature* **466**, 591-596 (2010).
22. Y. Malhi *et al.*, Logged tropical forests have amplified and diverse ecosystem energetics. *Nature* **612**, (2022).

23. A. Shade *et al.*, Macroecology to Unite All Life, Large and Small. *Trends Ecol Evol* **33**, 731-744 (2018).
24. S. M. a. Methods.
25. B. J. McGill *et al.*, Species abundance distributions: moving beyond single prediction theories to integration within an ecological framework. *Ecol Lett* **10**, 995-1015 (2007).
26. F. W. Preston, The Commonness, and Rarity, of Species. *Ecology* **29**, 254-283 (1948).
27. T. M. Blackburn *et al.*, The Relationship between Abundance and Body-Size in Natural Animal Assemblages. *J Anim Ecol* **62**, 519-528 (1993).
28. E. P. White, S. K. M. Ernest, A. J. Kerkhoff, B. J. Enquist, Relationships between body size and abundance in ecology. *Trends Ecol Evol* **22**, 323-330 (2007).
29. T. M. Blackburn, K. J. Gaston, Linking patterns in macroecology. *J Anim Ecol* **70**, 338-352 (2001).
30. J. Damuth, Population-Density and Body Size in Mammals. *Nature* **290**, 699-700 (1981).
31. M. I. O'Connor *et al.*, A general biodiversity-function relationship is mediated by trophic level. *Oikos* **126**, 18-31 (2017).
32. S. Tatsumi, M. Loreau, Partitioning the biodiversity effects on productivity into density and size components. *Ecol Lett*, (2023).
33. R. J. Whittaker, Meta-analyses and mega-mistakes: calling time on meta-analysis of the species richness-productivity relationship. *Ecology* **91**, 2522-2533 (2010).
34. D. Storch, E. Bohdalková, J. Okie, The more-individuals hypothesis revisited: the role of community abundance in species richness regulation and the productivity-diversity relationship. *Ecol Lett* **21**, 920-937 (2018).
35. R. Trebilco, J. K. Baum, A. K. Salomon, N. K. Dulvy, Ecosystem ecology: size-based constraints on the pyramids of life. *Trends Ecol Evol* **28**, 423-431 (2013).
36. B. J. Enquist, A. J. Abraham, M. B. J. Harfoot, Y. Malhi, C. E. Doughty, The megabiota are disproportionately important for biosphere functioning. *Nat Commun* **11**, (2020).
37. J. Sheridan, D. Bickford, Shrinking body size as an ecological response to climate change. *Nature Climate Change* **1**, 401-406 (2011).
38. B. J. McGill, J. C. Nekola, Mechanisms in macroecology: AWOL or purloined letter? Towards a pragmatic view of mechanism. *Oikos* **119**, 591-603 (2010).
39. B. J. McGill, Towards a unification of unified theories of biodiversity. *Ecol Lett* **13**, 627-642 (2010).
40. G. B. West, B. J. H., B. Enquist, A General Model for the Origin of Allometric Scaling Laws in Biology. *Science* **276**, 122-126 (1997).
41. D. Mouillot *et al.*, Rare Species Support Vulnerable Functions in High-Diversity Ecosystems. *Plos Biol* **11**, (2013)69. J. Chamagne *et al.*, Forest diversity promotes individual tree growth in central European forest stands. *J Appl Ecol* **54**, 71-79 (2017).