



HAL
open science

Can legume crop residues contribute to sustainable intensification of rainfed rice production in Madagascar?

Lalaina Ranaivoson, Gatien N. Falconnier, Patrice Autfray, Bertrand Muller, Louise Leroux, François Affholder, Aude Ripoche

► **To cite this version:**

Lalaina Ranaivoson, Gatien N. Falconnier, Patrice Autfray, Bertrand Muller, Louise Leroux, et al.. Can legume crop residues contribute to sustainable intensification of rainfed rice production in Madagascar?. International Crop Modelling Symposium (iCROP 2020), CIRAD; INRAE; INRIA, Feb 2020, Montpellier, France. 2 p. <hal-05174820>

HAL Id: hal-05174820

<https://hal.science/hal-05174820v1>

Submitted on 21 Jul 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Copyright - All rights reserved

Can legume crop residues contribute to sustainable intensification of rainfed rice production in Madagascar?

Ranaivoson Lalaina^{*1}, *Falconnier Gatien*², *Autfray Patrice*², *Muller Bertrand*^{3,4}, *Leroux Louise*², *Affholder François*², *Ripoche Aude*^{2,3}

¹Centre National de Recherche Appliquée au Développement Rural (FOFIFA), BP 1690, Antananarivo, Madagascar, lalaina.ranaivoson@fofifa.mg

²Centre de coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UPR AIDA, F-34398 Montpellier France AIDA, Univ Montpellier, CIRAD, Montpellier, France

³CRR FOFIFA, BP 230, Antsirabe, Madagascar

⁴Centre de coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UMR AGAP, F-34398 Montpellier France AGAP, Univ Montpellier, CIRAD, Montpellier, France

Rainfed agriculture on smallholder farms across the tropics is crucial for food security and livelihoods when availability of irrigated land is limited. Coarse-textured soils with low organic carbon and poor inherent fertility prevail in these systems (Tifton and Giller, 2013). In the absence of sufficiently remunerative market, smallholders cannot afford mineral fertilizers so that nutrient inputs are generally limited to manure. Integrated soil fertility management, *e.g.* integration of legumes combined with additional supply of mineral fertilizers, is required to sustainably increase rainfed agriculture productivity (Vanlauwe et al. 2014). In Madagascar, most smallholders do not use mineral fertilizer under rainfed agriculture. Incorporation of Nitrogen (N) rich crop residues could increase nutrient supply and improve N use efficiency (Aggarwal et al., 1997). Nutrient supply from residues is a complex process related to decomposition rates of residue, which is impacted by residue type and climate variability, *e.g.* increased temperatures can accelerate residue decomposition while intense rainfall can increase the loss of the precious mineralized N through leaching. The potential contribution of N rich crop residues to increase staple crop productivity is therefore complex to anticipate. Soil-crop models can account for such complexity. This study aims at (i) calibrating the STICS crop model for rice yield modeling in the smallholder context of cool humid uplands in Madagascar and (ii) use the model to explore the effect of incorporation of rice and legume residues for low (25 kg N ha⁻¹) and high (160 kg N ha⁻¹) fertilizer inputs with a variable climate.

The soil-crop model *STICS* (Brisson et al., 2003) was chosen for its capacity to account for soil water and nutrient dynamics during crop cycle for various climates and crop management. Rice experiments (cultivar NERICA 4) carried out in 2016-2017 and 2017-2018 cropping season in Ivory (19°33'S, 46°24'E, 950 m a.s.l) on Ferralsols were used for model calibration and testing. Two previous crops were compared: (i) rice and (ii) *Mucuna cochinchinensis* - *Crotalaria spectabilis* intercropping, combined with two levels of nutrient input, *i.e.* manure only (25 kg N ha⁻¹) or manure and mineral fertilizer (160 kg N ha⁻¹). Residues were incorporated at plowing. Measured data include rice phenology, above ground biomass and plant N, grain yield and in-season soil moisture and soil N.

The calibrated STICS model reproduced adequately rainfed rice emergence, flowering and maturity, with on average an error of less than four days. Simulated in-season soil N and rice biomass agreed with the observations with a relative Root Mean Square Error from 33% to 36% (see Figure 1 for rice residues with high N input treatment). The model could reproduce the additional mineral N supply and plant uptake associated with the incorporation of N rich legume residue for the two fertilizer treatments.

Calibration for grain yield and scenario analysis using historical climate (1980-2010) are on-going and will allow to determine the profitability and risk associated with the different simulated options. This study will contribute to gain new insights on the relevancy of legume residue for sustainable intensification of cropping systems in a tropical smallholder context.




Figure 1: Comparison of observed and STICS simulated rice biomass (a) and soil N (b) dynamics during rice growing cycle for rice residues with manure and mineral fertilizer (160 kg N ha⁻¹)

Keywords: upland rice, tropical climate, green manure, soil nitrogen, low-input

References

- Aggarwal RK, Kumar P, Power JF (1997) Use of crop residue and manure to conserve water and enhance nutrient availability and pearl millet yields in an arid tropical region. *Soil Tillage Res* 41:43–51. doi: 10.1016/S0167-1987(96)01082-3
- Brisson N, Gary C, Justes E, et al (2003) An overview of the crop model STICS. *Eur J Agron* 18:309–332
- Tittonell P, Giller KE (2013) When yield gaps are poverty traps: The paradigm of ecological intensification in African smallholder agriculture. *F Crop Res* 143:76–90. doi: 10.1016/j.fcr.2012.10.007
- Vanlauwe B, Coyne D, Gockowski J, et al (2014) Sustainable intensification and the African smallholder farmer. *Curr Opin Environ Sustain* 8:15–22. doi: 10.1016/j.cosust.2014.06.001