



HAL
open science

GNSS spoofing in conflict zones disrupts wildlife tracking and hampers research and conservation efforts

Frédéric Jiguet, Asaf Mayrose, Markus Piha, Ron Efrat, Tuomas Seimola, Nir Sapir, Sami Timonen, Yohay Wasserlauf, Aurélien Besnard, Pierrick Bocher, et al.

► **To cite this version:**

Frédéric Jiguet, Asaf Mayrose, Markus Piha, Ron Efrat, Tuomas Seimola, et al.. GNSS spoofing in conflict zones disrupts wildlife tracking and hampers research and conservation efforts. *Nature Communications*, 2025, 16, pp.1199. <10.1038/s41467-025-56630-2>. <hal-05323095>

HAL Id: hal-05323095

<https://hal.science/hal-05323095v1>

Submitted on 21 Oct 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

GNSS spoofing in conflict zones disrupts wildlife tracking and hampers research and conservation efforts

Frédéric Jiguet, Asaf Mayrose, Markus Piha, Ron Efrat, Tuomas Seimola, Nir Sapir, Sami Timonen, Yohay Wasserlauf, Aurélien Besnard, Pierrick Bocher, Andrea Kölzsch, Aleksi Lehikoinen, Anne-Christine Monnet, Petr Procházka & Yoav Perlman

 Check for updates

In war and conflict zones, the jamming of Global Navigation Satellite System (GNSS) signals by military forces disrupts the tracking of tagged animals, and has increased in frequency following the recent escalation of conflicts in Eastern Europe and the Middle East. Such disruption to data collection strongly hampers research into the protection and conservation of endangered animals.

For decades, scientists have been uncovering the secrets of animal movements using various technological solutions¹. Many of these technologies rely on the GNSS (including the Global Positioning System—GPS) to geolocate the tagged animals². Although originally developed for military purposes³, GNSS has gained enormous popularity in a wide range of civilian applications, including those related to research on animal movement and conservation biology. By receiving signals from multiple satellites, a GNSS tracking device can gain a high-accuracy location of the carrying animal almost everywhere on the globe within milliseconds. Yet due to the common use of GNSS-enabled applications in military operations, the disruption of GNSS signals is becoming frequent, especially in conflict areas. This jamming or spoofing of GNSS signals is intended to disrupt navigation systems of enemy forces. However, the non-specific nature of this electronic warfare affects all GNSS based devices, including multiple civilian applications such as civilian aircraft⁴ and research involving animal tracking devices⁵.

GNSS spoofing

We report here the strong effects of GNSS spoofing on bird tracking following the recent escalation of conflicts in Eastern Europe and the Middle East. By remotely monitoring the daily movements of birds across these regions during autumn 2023 to summer 2024, we repeatedly recorded erroneous positioning for many individuals across multiple species. This resulted in a significant loss of invaluable data, which in turn may have a severe impact on the ability to monitor and interpret animal movements and draw relevant conclusions for understanding their biology and developing conservation strategies. We observed such cases for multiple species (including eagles, falcons, shorebirds, and bustards), involving both resident and migratory birds. The erroneously recorded positions occurred over multiple countries and were often—but not exclusively—translocated to

international airports, for example in Russia, Ukraine, Lebanon, Jordan, Syria, and Egypt. We illustrate the issue with the tracks of black-tailed godwits (*Limosa limosa*) migrating from Finland to Romania while flying in or near Russia, Belarus, and Ukraine, and of Bonelli's eagles (*Aquila fasciata*) dispersing from their natal sites in Israel.

Black-tailed godwits

Fifteen black-tailed godwits were tagged in Finland in May 2024, as part of the Habitrack EU-funded research program (<https://habitrack.eu>). Breeding near Oulu or in Karelia, the godwits migrated southwards later in June or July. Most godwits followed a migratory pathway over Russia, Belarus and Ukraine leading to the Danube River delta in Romania. Eight displayed spoofed geolocations during their migrations, many birds being localized at the exact same place despite not migrating simultaneously (Fig. 1). We identified three obvious locations: west of St Petersburg (4 individuals), in Smolensk oblast (for 7 individuals migrating over Belarus), and in Crimea (5 individuals). For the latter, the tags mostly pointed to Simferopol airport in Crimea (45.037°N, 33.966°E) when the birds actually reached the Odessa region of Ukraine. As examples, one female godwit (ring ST340089, in pink) had her location moved to Simferopol airport on 22 June 2024 while she was flying just north of Mykolaiv. The total distance added to this migration track was estimated at 550 km, though covered in 16 min only. For another female godwit (ringed ST320391, in black), we recorded 9 locations at Simferopol airport, while the interleaved positions recorded by the tag indicated that the bird was in the Danube River delta in Romania, close to the Ukrainian border: 4 on 28 June, 1 on 29 June, 2 on 4 July, 1 on 14 July and 1 on 21 July. The distance between the actual locations and the airport ranged between 380 and 420 km, so that the nine erroneously-recorded round trips represented a total distance of ~7200 km. This however represents less than 0.1% of all locations (9 out of 9 053) obtained while the bird lingered in the Danube River delta between 26 June and 10 August 2024. Seven other godwits took more western migratory routes and their tracking devices did not record spoofed geolocations.

Bonelli's eagles

Forty-eight Bonelli's eagles were tracked with GPS tags across Israel between October 2023 and September 2024, as part of a national conservation program led by the Israel Nature and Parks Authority. Electrocutation has been identified as the primary threat to this population⁶. Most of the tagged eagles typically exhibit local dispersal movements within Israel and adjacent Jordan, Egypt, Lebanon, and Syria. Coinciding with escalating regional conflict from October 2023 onwards, an increase in GPS interference was observed. The majority




Fig. 1 | Tracking of Finnish black-tailed godwits migrating over Russia, Belarus and Ukraine to the Danube River delta in Romania. Details of GPS spoofing are illustrated in the region of (upper insert) Saint Petersburg, (middle insert) Smolensk (near the Belarus-Russian border), and (lower insert) Odessa (Ukraine) and Tulcea (Romania). Tracks of 8 individuals are shown, each bird with a different color, for tracking periods between mid-June and the end of September, depending on the individual. Black circles enclose spoofed locations. Godwits have been captured under licenses VARELY/968/2024 and POPELY/1475/2024, while tagging was

authorized under ESAVI/12867/2024. Black-tailed godwits were equipped with 5.5 g solar-powered GPS-GSM transmitters (Ornitema OT-6). The incubating adults were captured at nest with a clapnet at the late incubation stage. The tags were fitted by a leg-loop silicone harness (1.6 mm-diameter tube). Sampling frequency was set to every 10–60 min, depending on battery level. The total mass of the tag, including harness, was on average 2.5% of the bird's body mass. Data reported here is archived on the movebank.org platform, under the study identity reference 4595061729.




Fig. 2 | Tracks of 10 tagged Bonelli's eagles from 1 April to 1 September 2024, each individual represented with a different color. Black circles enclose spoofed locations. The inserted graph reports the daily proportion (in %) of spoofed locations for 18 randomly selected eagles out of the 48 tagged. Tracks of only ten individuals are shown for clarity. Bonelli's eagle were equipped with 30 g solar-powered GPS-GSM transmitters (Ornitema OT-30). The tagging process was performed in the nest when the nestlings were 45–60 days old. The tags were fitted by

a Teflon and Silicon pelvic harness (4 mm wide Teflon tube and 2 mm thick silicone string). Sampling frequency was set to every 15–60 min, depending on battery level. The total mass of the tag, including the harness, was below 3% of the bird's body mass. The eagle study is licensed by the Israeli Nature and Parks Authority (License 2495/2018). Data reported here is archived on the movebank.org platform, under the study identity reference 108979007.

of spoofed locations were diverted to international airports in the region, while some outliers appeared in a specific area in the Mediterranean Sea (Fig. 2). By early April 2024, interference levels reached 100% for some individuals, with the entire monitored population experiencing 20–50% spoofed locations (Fig. 2 insert). This disruption hampers efforts to identify risk factors such as hazardous power pylons, poisoning events, and direct persecution, thereby significantly weakening long-term conservation and mitigation efforts.

Handling spoofed positions

For scientists studying animal movements, these virtual translocations can be detected if they are truly outlying, or repeated. For example, an animal commuting back and forth daily to an international airport, or several birds that receive the same geolocation despite their routes diverge so they are clearly not migrating in the same flock. However, some cases might be less obvious and may require a refined knowledge of the species' typical movement patterns in order to be detected.

Whether obvious or not, researchers must consider the risk of location errors caused by GNSS spoofing when analyzing movement trajectories or habitat use. As we show with these data samples, such errors are widespread, and might appear in many new places in the future. GNSS spoofing in conflict zones poses a significant challenge to wildlife tracking and conservation efforts. This phenomenon compromises not only the accuracy of migration studies but also critical conservation activities such as mortality detection and epidemic monitoring. The implications extend beyond scientific research, potentially affecting endangered species management and human-wildlife conflict mitigation^{7–9}. In response to these challenges, researchers may start exploring potential solutions to mitigate the effects of GNSS spoofing. While advanced anti-spoofing algorithms and encrypted signals are being developed in other fields like civil aviation and military applications, such technologies have not yet been widely applied to wildlife tracking due to cost and complexity. Given the gravity of environmental crises worldwide and the ubiquity with which wildlife research relies on GNSS technologies, such solutions are no less imperative and should be developed and shared among practitioners.

Frédéric Jiguet¹✉, Asaf Mayrose^{2,3}, Markus Piha⁴, Ron Efrat⁵, Tuomas Seimola⁴, Nir Sapir², Sami Timonen⁶, Yohay Wasserlauf², Aurélien Besnard⁷, Pierrick Bocher⁸, Andrea Kölzsch^{5,9}, Aleksii Lehikoinen¹⁰, Anne-Christine Monnet¹, Petr Procházka¹¹ & Yoav Perlman¹²

¹Muséum National d'Histoire Naturelle, UMR 7204 CESCO MNHN CNRS Sorbonne Université, Paris, France. ²Department of Evolutionary and Environmental Biology and Institute of Evolution, University of Haifa, Haifa, Israel. ³Israel Nature and Parks Authority, Science Division, Jerusalem, Israel. ⁴Natural Resources Institute Finland, Helsinki, Finland. ⁵Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany. ⁶Center for Economic Development, Transport and the Environment, Oulu, Finland. ⁷CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France. ⁸UMR LIENSs, CNRS-La Rochelle University, La Rochelle, France. ⁹Department of Ecology, Radboud Institute for Biological and Environmental Sciences (RIBES), Radboud University, Nijmegen, The Netherlands. ¹⁰The Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland. ¹¹Institute of Vertebrate Biology, Czech Academy of Sciences, Brno, Czechia. ¹²BirdLife Israel, Society for the Protection of Nature in Israel, Tel Aviv, Israel. ✉e-mail: frederic.jiguet@mnhn.fr

Received: 8 October 2024; Accepted: 23 January 2025;
Published online: 30 January 2025

References

1. Kays, R., Crofoot, M. C., Jetz, W. & Wikelski, M. Terrestrial animal tracking as an eye on life and planet. *Science* **348**, 6240 (2015).
2. Jetz, W. et al. Biological Earth observation with animal sensors. *Trends Ecol. Evol.* **37**, 293–298 (2022).

3. Bonnor, N. A brief history of global navigation satellite systems. *J. Navig.* **65**, 1–14 (2012).
4. International Air Transport Association. EASA Partners with IATA to Counter Safety Threat from GNSS Spoofing & Jamming. Press Release No: 4, 26 January 2024. <https://www.iata.org/en/pressroom/2024-releases/2024-01-26-01/> (2024).
5. Nathan, R. et al. Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science* **375**, eabg1780 (2022).
6. Mayrose, A. et al. Bonelli's Eagle electrocution risk in Israel can be reduced by 80% by insulating only 4% of the pylons. *Ornith. Appl.* **126**, duae004 (2024).
7. Barton, P. S. et al. Guidelines for using movement science to inform biodiversity policy. *Environ. Manag.* **56**, 791–801 (2015).
8. Doherty, T. S. & Driscoll, D. A. Coupling movement and landscape ecology for animal conservation in production landscapes. *Proc. R. Soc.* **285**, 20172272 (2018).
9. Pekarsky, S., Schiffner, I., Markin, Y. & Nathan, R. Using movement ecology to evaluate the effectiveness of multiple human-wildlife conflict management practices. *Biol. Conserv.* **262**, 109306 (2021).

Acknowledgements

We thank helpers at godwit catching in Finland, namely Jimmy Allain, Esko Pasanen, Jaana Rintala, Laura Oinas, Karoliina Hämäläinen, Mika Jokikokko, Marie Donnez, and Carole Doucerain. Research on godwits was conducted by the Habitrack research consortium and is funded by the European Union; views and opinions are however those of the authors only and do not necessarily reflect those of the EU, which cannot be held responsible for them. The research on Bonelli's eagles is funded and performed by the Israel Nature and Parks Authority (INPA), as part of the Endangered Raptor Conservation project, which is a collaboration between INPA, Israel Electric Company and the Society for the Protection of Nature in Israel (SPNI). We would like to thank the steering committee of the project: Ohad Hatzofe, Uri Naveh, Noam Leader and Roni Elias. Accessing the nests and tagging juvenile eagles were performed by: Walter Nesor, Yotam Orchan, Betachia Rokovsky, Noam Maizliesh, and Dudu Yifrach. Surveys and nest monitoring were done by many birdwatchers, INPA rangers, SPNI team, and ecologists.

Author contributions

Conceptualization: F.J., Y.P., A.M., Writing—original draft: F.J., A.M., M.P., R.E., T.S., N.S., S.T., Y.W., A.B., P.B., A.K., A.L., A.C.M., P.P., Y.P., Writing—review & editing: F.J., A.M., Funding acquisition: F.J., A.M., M.P., R.E., T.S., N.S., S.T., Y.W., A.B., P.B., A.K., A.L., P.P., Y.P.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Frédéric Jiguet.

Peer review information *Nature Communications* thanks Daniel Ruthrauff and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2025