



**HAL**  
open science

## Autistic Spectrum disorder and Anaesthesia

Maria I Dalamagka

► **To cite this version:**

Maria I Dalamagka. Autistic Spectrum disorder and Anaesthesia. World Journal of Advanced Research and Reviews, 2024, 21, pp.1656 - 1658. <10.30574/wjarr.2024.21.3.0918>. <hal-05432487>

**HAL Id: hal-05432487**

**<https://hal.science/hal-05432487v1>**

Submitted on 26 Dec 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



(CASE REPORT)



## Autistic Spectrum disorder and Anaesthesia

Maria I Dalamagka \*

*Anaesthesia Department, General Hospital of Larisa, Greece*

World Journal of Advanced Research and Reviews, 2024, 21(03), 1656–1658

Publication history: Received on 12 February 2024; revised on 19 March 2024; accepted on 21 March 2024

Article DOI: <https://doi.org/10.30574/wjarr.2024.21.3.0918>

### Abstract

One of the groups of patients which may require the greatest flexibility of approach is those with autistic spectrum disorder (ASD). ASD is a lifelong developmental disability, affecting four times as many males than females, which affects how a person communicates with, and relates to, other people and the world around them. Challenging behaviors may be triggered by the unfamiliar perioperative environment and may make anaesthesia related procedures difficult or impossible. Children with more severe ASD may exhibit verbal or physical aggression, antisocial or disruptive behavior, temper tantrums, screaming, panic attacks, and self-injurious behavior, and may have aberrant responses to sensory stimuli. Although anaesthesia and sedation do not present a problem for most children with ASD, unpredictable regression in skills and behaviour is noted in a small number of patients after general anaesthesia. A 10-year-old child, 53 kg, with autism under treatment, came for a dental abscess and related dental work under general anaesthesia. The pediatric neurological assessment indicated hyperactivity and lack of cooperation. Post-operatively, during resuscitation, the child showed agitation and an attempt to withdraw oxygen, with the physical presence of his family environment.

**Keywords:** Autism; Anaesthesia; ASD; Disability; Epilepsy; Behavior

### 1. Introduction

Autism spectrum disorder (ASD) is characterized by deficits in social communication and interaction; restrictive, repetitive patterns of behavior and interests that may include inflexible adherence to nonfunctional routines or rituals; and hyper- or hyposensitivity to sensory input. The prevalence of Autistic spectrum disorders (ASD) is estimated to be 1% worldwide. The incidence of autism is about 0.2%. One of the groups of patients which may require the greatest flexibility of approach is those with autistic spectrum disorder (ASD). ASD is a lifelong developmental disability, affecting four times as many males than females, which affects how a person communicates with, and relates to, other people and the world around them. This complex mental disability is about four times more prevalent in males but is more severe in females. It manifests during the first 3 years of life. It is important to recognize that for almost all cases, genetic testing is unlikely to establish a diagnosis of autism in the absence of careful clinical evaluation, since Fragile X Mental Retardation-1 gene, maternally inherited 15q11-q13 duplications, and other syndromes greatly increase the risk for autism, but do not lead to autism spectrum disorders in all cases. Epilepsy is an enduring predisposition for generating seizures which is active in approximately 0.4%–0.8% of the population. The majority of studies that examined sex differences in people with ASD found an increased risk in males, which is consistent with higher male prevalence for ASD in the general population. These results support the strong connection between intellectual disability, epilepsy and ASD [1-8].

\* Corresponding author: Maria I. Dalamagka

## 2. Case Study

A 10-year-old child, 53 kg, with autism under treatment, came for a dental abscess and related dental work. The pediatric neurological assessment indicated hyperactivity and lack of cooperation. Without cerebellar semiology, pyramidal and extrapyramidal involvement, as well as without neurodermal spots. His gait was normal, with no scoliosis present. Medication included Sir. Risperdal 0.75 ml, Melatonin spray 2 mg. Preanesthesia was done with 8 mg of Dormicum intranasally 20 minutes before induction into anesthesia. Anesthesia was induced with 0.1 mg Fentanyl, 180 mg Propofol and 50 mg Esmeron. An additional 0.1 mg of Fentanyl was given and anesthesia was maintained with 6% Desflurane in a mixture of 50% oxygen and 50% air. The child was mechanically ventilated on a Drager ventilator and volume control ventilation model. The duration of the surgery was 2.5 hours and the awakening, with 100% oxygen, took place in 15 minutes with the administration of Bridion 150 mg. Post-operatively, during resuscitation, the child showed agitation and an attempt to withdraw oxygen, with the physical presence of his family environment.

### 2.1. Management and outcome

Many children with special needs may attend hospital for investigations, medical management, or surgery. They may exhibit high levels of anxiety when faced with hospital treatment, and may have difficulty conforming to the usual pattern of care. In extreme cases, children may become so uncooperative that their procedure is postponed or abandoned, or can only be undertaken with the use of heavy sedative premedication or restraint. Anesthesia providers generally are aware of the prevalence of diagnosed ADHD and the various drugs these children might be on. Autistic children might also be on stimulant or antipsychotic drugs; therefore, they must consider specific, necessary anesthetic considerations. These drugs are combined with certain anesthetic drugs and an increase in central nervous system depression might result. Recommendations include good hydration, minimal fasting, care with Hartmann's solution (due to elevated blood lactate levels), maintenance of normal blood glucose, body temperature and acid–base balance, and avoidance of oxidative stress.

---

## 3. Discussion

Although there is a range of severity of ASD, most children with ASD are uncomfortable with or upset by unexpected or new situations. Challenging behaviors may be triggered by the unfamiliar perioperative environment and may make anesthesia related procedures (eg, application of a face mask for induction, intravenous catheter placement, application of monitors) difficult or impossible. Children with more severe ASD may exhibit verbal or physical aggression, antisocial or disruptive behavior, temper tantrums, screaming, panic attacks, and self-injurious behavior, and may have aberrant responses to sensory stimuli. A goal for perioperative care should be to avoid these difficulties, for patient comfort, and to avoid negative associations with the hospital environment.

---

## 4. Conclusion

Although anaesthesia and sedation do not present a problem for most children with ASD, unpredictable regression in skills and behaviour is noted in a small number of patients after general anaesthesia. It may be appropriate therefore to adopt an anaesthetic technique suitable for patients with mitochondrial disease.

---

## Compliance with ethical standards

### *Statement of informed consent*

Informed consent was obtained from all individual participants included in the study.

---

## References

- [1] Khan NZ, Gallo LA, Arghir A, et al. Autism and the grand challenges in global mental health. *Autism Res* 2012; 5: 156–59.
- [2] Bauman ML, Kemper TL. Neuroanatomic observations of the brain in autism: a review and future directions. *Int J Dev Neurosci* 2005; 23: 183–87.
- [3] American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*, 5th edn Washington, DC: American Psychiatric Association Publishing, 2013.

- [4] Lord C, Petkova E, Hus V, et al. A multisite study of the clinical diagnosis of different autism spectrum disorders. *Arch Gen Psychiatry* 2012; 69: 306–13.
- [5] Maenner MJ, Rice CE, Arneson CL, et al. Potential impact of DSM-5 criteria on autism spectrum disorder prevalence estimates. *JAMA Psychiatry* 2014; 71: 292–300.
- [6] Weitlauf AS, Gotham KO, Vehorn AC, Warren ZE. Brief report: DSM-5 “levels of support:” a comment on discrepant conceptualizations of severity in ASD. *J Autism Dev Disord* 2014; 44: 471–76.
- [7] Sicherman N, Loewenstein G, Tavassoli T, Buxbaum JD. Grandma knows best: Family structure and age of diagnosis of autism spectrum disorder. *Autism* 2018; 22: 368–76.
- [8] Weitlauf AS, McPheeters ML, Peters B, et al. Therapies for children with autism spectrum disorder: behavioral interventions update. Comparative Effectiveness Review No. 137. Rockville, MD: Agency for Healthcare Research and Quality, 2014