



HAL
open science

Molecular underpinning of heavy metal sequestration through advanced remediation strategies in higher plants

Himani Agarwal, Divya Chaudhary, Himanshi Aggarwal, Chhavi Karala, Niharika Purkait, Neha Sharma, Arti Mishra, Vaibhav Mishra, Ajay Kumar, Prashantkumar Singh, et al.

► To cite this version:

Himani Agarwal, Divya Chaudhary, Himanshi Aggarwal, Chhavi Karala, Niharika Purkait, et al.. Molecular underpinning of heavy metal sequestration through advanced remediation strategies in higher plants. *Plant stress*, 2025, 16, pp.100881. 10.1016/j.stress.2025.100881 . hal-05490812

HAL Id: hal-05490812

<https://hal.science/hal-05490812v1>


Submitted on 3 Feb 2026

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Review

Molecular underpinning of heavy metal sequestration through advanced remediation strategies in higher plants

Himani Agarwal^{a,1}, Divya Chaudhary^{a,1}, Himanshi Aggarwal^{a,1}, Chhavi Karala^{a,1}, Niharika Purkait^{a,1}, Neha Sharma^b, Arti Mishra^{c,d}, Vaibhav Mishra^a, Ajay Kumar^e, PrashantKumar Singh^g, Laurent Dufossé^{f,*}, , NaveenChandra Joshi^{a,**}

^a Amity Institute of Microbial Technology, Amity University, Uttar Pradesh, India

^b Department of Life Science, Faculty of Allied Health Sciences, SGT University, Gurugram, Haryana, India

^c Department of Botany, Hansraj College, Delhi University, India

^d Umeå Plant Science Centre (UPSC), Department of Plant Physiology, Umeå University, Umeå 901 87, Sweden

^e Amity Institute of Biotechnology, Amity University, Uttar Pradesh, India

^f Laboratoire de Chimie et Biotechnologie des Produits Naturels, ESIROI Agroalimentaire, Université de La Réunion, CHEMBIOPRO, 15 Avenue René Cassin Ile de La Réunion, Saint-Denis F-97400, France

^g Department of Biotechnology, Mizoram Central University, Mizoram, India



ARTICLE INFO

Keywords:

Nano-bioremediation
Heavy metals remediation
HMs ATPase
ZIP family
Phytoremediation
Phytochelatin

ABSTRACT

Anthropogenic emissions, particularly from industrial and agriculture activities, have significantly elevated the concentrations of highly toxic Heavy Metals (HMs), such as lead (Pb), cadmium (Cd), and arsenic (As), in the soil, leading to their accumulation in plants. These HMs, when exceeding toxicity thresholds (e.g., Pb >10 mg/kg, Cd >0.5 mg/kg, As >1 mg/kg), disrupt the plant physiology and metabolism. To mitigate this toxicity, plants employ diverse detoxification and sequestration strategies, including mycorrhizal associations, root exudates, cellular compartmentalization, and the production of organic acids, phytochelatin, metallothioneins, proline, stress proteins, and plant hormones. This review aims to critically examine the molecular mechanisms by which key crop plants, such as rice, wheat, maize, and other higher plants, sequester these primary heavy metal contaminants. Additionally, it highlights the role of nanotechnology in enhancing plant resistance and facilitating nano-bioremediation under HMs stress conditions. This review provides valuable insights into innovative clean-up strategies for agriculturally important crops by exploring nanoparticle-mediated remediation mechanisms.

1. Introduction

The geosphere is made up of rocks, minerals, and heavy metals (HMs). It starts with soil surface and extends to sand beaches, mountain summits, and molten rock at the earth's center (Mukherjee et al., 2023). HMs are found in soil as natural components or by spontaneous occurrence such as volcanic eruptions. Moreover, it also occurred due to anthropogenic activities like mining or construction. These HMs enter the ecosystem and take part in biogeochemical processes, thereby increasing their spread in the earth's ecosystems (Zhang et al., 2010).

According to reported literatures the most frequent contaminants that are toxic to human and animal health, even at low concentrations, are heavy metals like zinc (Zn), nickel (Ni), chromium (Cr), copper (Cu), mercury (Hg), arsenic (As), (metalloid), etc. (Lawal et al., 2021). Plants need some HMs like Zn, Cu, Ni, etc. as micronutrients to support normal growth and developmental activities. However, the quantity and concentration of both essential and non-essential nutrients must be carefully balanced to ensure their availability does not hinder plant development under natural circumstances (Hall, 2002) (Table 1).

Higher angiosperm and gymnosperm includes a diverse array of

Top of Form

* Corresponding author.

** Co-corresponding authors.

E-mail addresses: laurent.dufosse@univ-reunion.fr (L. Dufossé), naveen.joshi.jnu@gmail.com (N. Joshi).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.stress.2025.100881>

Received 6 December 2024; Received in revised form 5 May 2025; Accepted 6 May 2025

Available online 7 May 2025

2667-064X/© 2025 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Table 1
Essential heavy metals (Play a Metabolic Role).

Sl. No	Metal	Metabolic Role in Plants	Refs.
1.	Zinc (Zn)	Enzyme cofactor, protein synthesis, auxin production	Mousavi et al. (2013)
2.	Copper (Cu)	Photosynthesis, respiration, oxidative stress protection	Droppa & Horváth (1990); Ravet & Pilon (2013)
3.	Iron (Fe)	Chlorophyll synthesis, electron transport, enzyme cofactor	Kroh & Pilon (2020)
4.	Manganese (Mn)	Photosynthesis (PSII cofactor), antioxidant defence	Ducic & Polle (2005); Takashima et al. (2012)
5.	Nickel (Ni)	Urease enzyme activation, nitrogen metabolism	Nim & Wong (2019); Lavres et al. (2016)
6.	Molybdenum (Mo)	Nitrate reduction, nitrogen fixation	Demtröder et al. (2019)
Non-essential Heavy metals (No metabolic Role)			
1.	Cadmium (Cd)	No known essential role; may induce stress response pathways.	Haider et al. (2021)
2.	Lead (Pb)	No beneficial role; can accumulate in root cell walls.	Zulfiqar et al. (2019)
3.	Mercury (Hg)	No beneficial role; highly toxic in ionic and methylated forms.	Clemens, (2013)
4.	Chromium (Cr)	Some plants show enhanced stress tolerance with Cr (III) at very low levels.	Cervantes et al. (2001)
5.	Aluminium (Al)	Some plants (e.g., tea, Camellia) accumulate Al to enhance defence mechanisms.	Poschenrieder et al. (2013); Panda et al. (2009)
6.	Arsenic (As)	Some plants can tolerate arsenic and use it in stress signalling.	Bali & Sidhu (2021)

plant species that serve as rich sources of staple foods, medicinal compounds, and fruits. Among them, angiosperms represent the most evolutionary advanced and ecologically dominant group of land plants. They play a central role in global food security by providing primary food crops such as rice, wheat, and maize, as well as widely consumed fruits including apples, oranges and bananas. However, these plants are increasingly threatened by the accumulation of toxic heavy metals, notably Cd, Pb, Hg, and As. The bioaccumulations of these metals in the edible plant tissues not only poses severe health hazards to consumers but also disrupts ecological balance by introducing contaminants into the food chain. Therefore, exploring the mechanism of heavy metals sequestration in the higher plants and developing eco-friendly or cost-effective remediation strategies is crucial for promoting sustainable agriculture (Kafle et al., 2022). The visual signs of HMs toxicity brought on by its high concentrations occur at the molecular level. This toxicity is due to the bonding between metal and proteins, which impairs the functionality of biomolecules like pigments and enzymes, compromising the cell membrane's integrity and suppressing essential plant processes like photosynthesis and respiration, etc. (Rubino, 2015). Moreover, HMs contribute to the production of free radicals, reactive oxygen species (ROS) i.e., superoxide ions (O_2^-), hydroxyl ions (OH), and hydrogen peroxide (H_2O_2). In addition, toxic compounds like methylglyoxal (MG) tend to accumulate under several stress conditions, including HMs exposure and elevated ROS levels (Aslam et al. 2021). This accumulation of MG and ROS in plants interferes with the metabolic and physiological functions like photosynthesis, by deactivating the antioxidant defense system and disrupting internal homeostasis (Sytar et al., 2013).

Additionally, it is essential to explore innovative and eco-friendly remediation strategies that not only mitigate soil contamination but also aid in restoring soil health. The integration of nanotechnology into agriculture marks a major breakthrough in addressing the complex challenges of global food production (Zhang et al., 2022). The extensive application of nanomaterials in precision agriculture, nutrient management, pest control, and soil health offers unprecedented

opportunities for the adoption of sustainable and efficient farming practices (Yadav et al., 2023). In recent years, nanotechnology has gained significant recognition as an innovative and promising strategy for addressing the challenges associated with soil contamination in agricultural settings (Sable et al., 2024). The unique characteristics of nanomaterials, such as their large surface area, high reactivity, and ability to alter physicochemical properties, make them highly effective for soil remediation.

Nanotechnology has the potential to revolutionize traditional soil remediation techniques by offering more efficient, precise, and environmentally sustainable solutions for mitigating agricultural contamination (Dhanapal et al., 2024). This review aims to consolidate existing research on the application of nanotechnology in remediating contaminated agricultural soils. Through a detailed analysis of recent advancements, it explores various nanomaterials, nanocomposites, and nanotechnological methods used to remove, contain, and neutralize soil pollutants. Additionally, the review critically assesses the effectiveness, environmental implications, and possible risks associated with the use of nanotechnology in agricultural soil remediation.

To check the relevance of the topic we made a bibliometric analysis of the research papers published on the related topics and in the bibliometric data study, we retrieved a total of 24,597 articles between 2020 and 2024 from the Scopus database by searching the keywords "Heavy metals", AND "Bioremediation" or "bioaccumulation", or "biomagnification" or "Toxicity" Or "Cd" or "Hg" or "Pb" or "Bacteria" or "Mycoremediation" or "Exopolysaccharides" or "Oxidative stress", or "Water" Pollution and created the bibliometric analysis using VOS viewer processing software (v1.6.9) by selecting 10,000 articles. The different colours present in the network showed linked terms; however, the size of the labels showed the number of publications, and the distances present between two nodes represent the degree through which they are associated (Fig. 1). This study explores the diverse strategies employed by plants to mitigate the effects of HMs contamination, with a particular emphasis on the molecular mechanisms underlying HMs sequestration in higher plants.

2. HM uptake, translocation and distribution in plants

The process of absorbing HMs by plants begins in the soil through their respective roots and proceeds through the root hairs. The metal crosses the root tissues apoplastically and makes its way in the symplast of the endodermis cell which later passes into the cells of stele, facilitated by the plasmodesmata (Grennan, 2009). Metal ions are transported to the shoots through a process known as transpiration pull, which is the mass upward flow of water. Plants passively absorb nutrients in one of two ways: assisted diffusion or simple diffusion (Annamalainathan et al., 2013). Plants inhabiting metal-contaminated soil have established three basic defense strategies. The first one is metals excluders; these types of plants restrict HMs uptake and maintain low concentration in their respective shoots over many different soil levels. A plant is considered an excluder for Cd if its shoot concentration is below 10 mg/kg, for Zn if it is below 100 mg/kg, and for Ni if it is below 50 mg/kg, even when these metals are present in higher concentrations in the soil (Backer and Brooks, 1989; Cobbett, 2000). The second type of plants captures metals in their aerial tissues and act as a good metal contamination indicator. Consequently, a plant is considered a metal contamination indicator for Cd if its shoot concentration exceeds 1–10 mg/kg, for Zn if it exceeds 200–300 mg/kg, and for Ni if it exceeds 50–100 mg/kg, signaling contamination in the environment (Cobbett, 2000; Raskin et al., 1997; Backer and Brooks, 1989). The third category includes accumulators or hyperaccumulators. In this group plants begin to concentrate HMs in the above ground tissues at levels higher than those found in the surrounding soil (Verbruggen et al., 2009). Therefore, a plant is considered a hyperaccumulator if its shoot concentrations exceed 100 mg/kg for Cd, 1000 mg/kg for Zn and Ni (Backer and Brooks, 1989). Some examples of the common hyperaccumulators,

Table 2
Examples of Metal excluders, metal indicators and metal hyperaccumulators.

Sl. no.	Defense strategies	Plant examples	Metals	Refs.
		<i>Agrostis capillaris</i>	Cu, Zn, Cd, and Pb	Dahmani-Muller et al. (2000)
		<i>Bidens pilosa</i>	As excluder	Sun et al. (2009)
		<i>Commelina communis</i>	Cd-excluders	Wei et al. (2005)
		<i>Digitaria sanguinalis</i>	Cd-excluders	Wang et al. (2020)
		<i>Oenothera biennis</i>	Cd-excluders	Wei et al. (2005)
		<i>O. biennis</i>	Cu-excluder	Wei et al. (2005)
		<i>Silene vulgaris</i>	Cu, Zn, and Cd	Schat & Vooijs (1997)
		<i>Taraxacum mongolicum</i>	Zn-excluder	Wei et al. (2005)
		<i>Thlaspi arvense</i>	Cd-excluders	Martin et al. (2012)
		2.	Metal indicators	<i>Ageratum conyzoides</i>
<i>Phragmites australis</i>	Cd, Cu, Pb, and Zn			Bonanno & Giudice (2010); Phillips et al. (2015)
<i>Spartina maritima</i>				
<i>Typha capensis</i>				
<i>Arabidopsis halleri</i>	Cd			Zhao et al. (2000); Zhao et al. (2006)
3.	Metal hyperaccumulators	<i>Amaranthus gangeticus</i>	Cd	Zhou et al. (2013)
		<i>Alyssum</i> sp.	Ni	Mengoni et al. (2011); Broadhurst and Chaney (2016)
		<i>Brassica juncea</i>	Cd	Haag-Kerwer et al. (1999)
		<i>Berkheya codii</i>	Ni	Slatter (2013)
		<i>Eichhurnia crassipes</i>	Cd	Wang et al. (2012)
		<i>Helianthus annuus</i>	Cd	Júnior et al. (2015)
		<i>Microsorium pteropus</i>	Cd	Lan et al. (2018)
		<i>Noccaea</i> sp.	Cd, Zn, Ni	Sterckeman et al. (2017)
		<i>Phytolacca americana</i>	Cd	Peng et al. (2008)
		<i>Pteris vittata</i>	As	Kohda et al. (2021)
			Pb	Soongsombat et al. (2009)
		<i>Ptyrogramma calomelons</i>	Ar	Campos et al. (2018)
			Pb	Soongsombat et al. (2009)
		<i>Salix caprea</i>	Cd, Zn	Konlechner et al. (2013)
		<i>Sedum</i> sp.	Cd	Gao et al. (2013)

3.1. Extracellular speciation

Extracellular speciation refers to the interactions of HMs outside the cell, including in the rhizosphere, cell wall, and apoplastic space. Plants root release exudates with chelator properties into the rhizosphere which serves as the primary defense against the HMs uptake. In *Arabidopsis thaliana*, the uptake of Ni from soil was greatly reduced by the presence of Ni chelating histidine and citrate in root exudates (Nishida et al., 2011). However, the binding of HMs to plant cell wall components (pectin, cellulose and hemicellulose) is influenced by the availability of functional groups such as carboxyl (-COO⁻), hydroxyl (-OH), amino (-NH₂), and sulfhydryl (-SH), thereby facilitating the toxic HMs ions to be immobilized and prevents their entry into the cytosol. Pectin has the

highest metal binding capacity, particularly for Pb²⁺, Cr³⁺, and Cu²⁺, due to its abundant carboxyl (-COO⁻) groups. The cellulose and hemicellulose components bind to Cd²⁺, Zn²⁺, and Cr³⁺ through hydroxyl (-OH) groups, though with moderate efficiency (Krzesłowska, 2011). Whereas, bioavailability and speciation in plant-soil system are impacted by microbial activity in the soil matrix (Pierart et al., 2015). Some reports have shown an alteration in the phytoavailability of As in soil after microbial inoculation in soil, the microbes stimulate redox reactions between As(III) and As(V) (Gorny et al., 2015). Some bacterial species accountable for these interconversions are *Desulfotomaculum auripigmentum*, *Thermus thermophilus*, *Bacillus arsenicus*, *Geospirillum arsenophilus*, etc. The absorption sites of the cell wall are restricted, which accounts for its marginal impact on metal tolerance. Whereas, in *Arabidopsis halleri*, a negative correlation between Fe concentration and uranium uptake was studied, with Fe deficiency considerably increasing uranium uptake (Vaxevanidou et al., 2012; Yamamura and Amachi, 2014).

3.2. Intracellular speciation

In intracellular speciation, once HMs enter plant cells, they undergo intracellular detoxification and compartmentalization to minimize toxicity. HMs are promptly bounded by the intended cellular molecules upon arrival into the cytosol, eliminating the risk of toxicity brought on by metal ions freely available and enabling a number of synchronized metabolic activities including specialized integration in metalloproteins or vacuolar sequestration (Haydon and Cobbett, 2007). Nicotianamine (NA), a metal chelating agent, possess a high level of stability while binding transition metal ions along with some organic acids like citrates and is necessary for maintaining Fe, Zn and Cu homeostasis (Curie et al., 2009; Clemens et al., 2013). Furthermore, the formation of HM compounds in plants is aided by 1 mM GSH, a low-molecular-weight chelator, which triggers the production of PC's by phytochelatin synthase (M_w 95,000) that create metal-PC complexes, and these complexes are transported into vacuoles by an ABC-type transporter, thereby enhancing the plant's tolerance to HMs (Grill et al., 1989; Srivastava, 2016). In *A. thaliana*, upregulation of a critical enzyme in GSH production led to alleviate GSH concentration levels and improved tolerance of HMs like Co, Zn and Ni. Similarly, tolerance towards different HM's (Ni, Co, Zn, Cd) was seen in *A. thaliana* when *Thlaspi goesingense* gene serine acetyltransferase (TgSAT) was overexpressed in the former which led to three fold increase in GSH levels (Freeman et al., 2004).

4. Factors influencing metal uptake in plants

Thermodynamic activities and solubility of metal ions are vital factors that governs the HMs availability through rhizosphere in plants, because root uptake requires a soluble ion. In addition, thermodynamic activity refers to the effective concentration of a heavy metal ion in a system, considering its interactions with other ions and molecules whereas actual uptake refers to the amount of HMs absorbed by plant roots and translocated into tissues. This depends on metal bioavailability, root absorption mechanisms, and plant metabolism (Shukla et al., 2019). It has been observed that with an increase in thermodynamic activity the bioavailability thereby increasing the uptake of HMs (Kalis, 2006). Metal solubility is subjected to alterations due to the physiochemical properties of soil including composition of organic and inorganic solutes, oxidation-reduction potential and soil pH. Subsequently, soil Cation Exchange Capacity (CEC) plays a crucial role in determining the bioavailability of HMs by regulating their retention, mobility, and uptake in soil-plant systems. High CEC soils, such as those rich in clay minerals and organic matter, strongly adsorb HMs like Pb²⁺, Cd²⁺, and Zn²⁺, reducing their solubility and plant availability. Conversely, low CEC soils, such as sandy soils, provide fewer binding sites, increasing metal mobility and bioavailability, which can lead to environmental contamination and plant toxicity (Alloway, 2012).

Additionally, soil pH significantly influences CEC, as acidic conditions reduce metal binding and enhance solubility, while alkaline conditions promote metal precipitation and adsorption onto soil particles. To mitigate HMs risks, soil amendments like biochar, compost, and lime can be applied to increase CEC and regulate metal availability, improving soil health and reducing toxicity (Kumpiene et al., 2008). These soil characteristics are quite diverse and depend on a number of different geographic elements, including terrain, biological processes, parent rock, climate, and anthropogenic activities. As soil pH rises above 5.5–6, the solubility of many metals like Cd, Pb, and Cu generally decreases, making chelating agents like Ethylenediaminetetraacetic acid (EDTA) useful for promoting metal translocation in plant tissues and enhancing phytoextraction over extended exposure (Roy et al., 2005). Moreover, EDTA, a synthetic ligand, binds to heavy metals like Cd²⁺, Zn²⁺, and Pb²⁺ through its carboxylate and amine groups, keeping them in their soluble form even at higher pH levels, preventing precipitation as insoluble hydroxides, and enhancing their mobility in soil for plant uptake, which is essential for phytoextraction (Shahid et al., 2014). A list of binding agents and their thresholds against heavy metals in various plant species is given in Table 3.

Temperature also determines the solubility and diffusion rates of HMs. It has been observed that elevated temperatures can stimulate root metabolic activity and transpiration, leading to greater metal translocation to aerial parts of the plant (Yu et al., 2010). However, excessive temperatures may disrupt membrane integrity and metal transport proteins, altering uptake efficiency. Conversely, lower temperatures reduce root permeability and enzymatic activity, restricting metal absorption and slowing translocation. The impact of temperature also varies by metal type; for instance, it has been observed that in rice Cd uptake tends to increase with temperature (Ge et al., 2016).

A variety of plant factors, including root characteristics, root-soil interactions, physiological processes, species-specific traits, growth stages, nutrient availability, environmental stress, mycorrhizal associations, and climate conditions, influence the uptake of HMs from the soil by affecting metal absorption, transport, and detoxification mechanisms. Plants possess very precise and efficient ways for obtaining critical micronutrients from their micro-environment. These pathways are also involved in the absorption, transport, and storage of harmful elements, playing a crucial role in the process of bioremediation. HMs including Zn, Cu, Mn, Pb, Ni, and Co can be captured and accumulated by metal-accumulating plant species up to 100–1000 times more than non-accumulating plant species (Sheoran et al., 2010). Certain plant species, such as *Thlaspi caerulescens*—a well-known zinc (Zn) hyper-accumulator, can accumulate up to 10,000 mg/kg in their leaves. This is approximately 1000 times more than the Zn levels typically found in normal plants, which usually contain <10 mg/kg. In contrast,

non-accumulating plants growing in Zn-rich soils generally accumulate only about 30–100 mg/kg of Zn in their tissues (Asad, 2011).

5. Different types of metal transporters with their roles in HMs translocation

Hyperaccumulators use a range of metal transporter proteins, including those from the Zn regulated, Iron regulated transporter-like proteins (ZIP), Cation diffusion facilitator (CDF), Natural Resistance Associated Macrophage Protein (NRAMP) and P_{1B}-type HM ATPase (HMA) families, to facilitate metal uptake and maintain homeostasis, with these transporters playing a crucial role in regulating metal accumulation and tolerance at the cellular level (Fig. 2).

5.1. ZIP family

The roots and shoots of several dicot plant species are known to utilize transporters from the ZIP family, such as Iron regulated transporter (IRT) and Zinc Regulated Transporter (ZRT) proteins (Grotz and Gueriot, 2006). ZIP proteins have eight transmembrane domains crossing the plasma membrane with both the amino (-NH₂) and carboxy (-COO) terminal ends located on the outermost part of the membrane, and they vary in length from 309 to 476 amino acids due to a histidine-rich variable region, spanning from transmembrane domain III to domain IV, that resides in the cytoplasm (Gueriot, 2000). *AtIRT1*, *OsIRT1*, and *HvIRT1*, homologs of *A. thaliana*, *Oryza sativa*, and *Hordeum vulgare*, respectively, are crucial for Fe²⁺/Fe³⁺ and Mn²⁺ uptake, being located on the plasma membrane; under Zn-limiting conditions, the transcript levels of *A. thaliana* ZIP family transporters, including *AtZIP1* to *AtZIP5*, *AtZIP9* to *AtZIP12*, and *AtIRT3*, are elevated (Wintz et al., 2003; Talke et al., 2006; Van De Mortel et al., 2006).

5.2. CDF protein family

The CDF proteins, belonging to the class of cation efflux transporters crucial for HMs maintenance and tolerance, include plant CDFs, typically Metal Tolerance Proteins (MTPs) transporting Mn²⁺. Phylogenetic analysis divided the CDF family into three groups based on metal ion transport efficiency: (1) Mn²⁺-transporting CDFs, (2) Fe²⁺/Zn²⁺-transporting CDFs, and (3) Zn²⁺ and other metal ion (except Mn²⁺ or Fe²⁺) transporting CDFs. A study identified substrate-defined clades for Cd²⁺, Ni²⁺, and Co²⁺ transport within the CDF family using phylogenomic data (Montanini et al., 2007). CDF proteins play a role in binding or exporting cations like Zn from the cytoplasm to internal compartments through sequestration or efflux, affecting cation accumulation and tolerance, signal transduction, oxidative stress resistance, and protein

Table 3
Binding agents and their thresholds against heavy metals in various plant species.

S. No.	Plant species	Binding agents	Heavy metals	Concentration	Effect on plant	Refs.
1.	<i>Corchorus capsularis</i> L.	citric acid (CA)	Cu	2.0 mmol kg ⁻¹	Alleviate oxidative stress and Cu phytotoxicity	Parveen et al. (2020)
2.	<i>Helianthus annuus</i> L.	ethylenediamine-N,N'-disuccinic acid {(S,S)-EDDS}	Cd, U (Uranium)	5.0 mmol kg ⁻¹	Enhanced oxidative stress	Chen et al. (2020)
3.	<i>Macleaya cordata</i>	citric acid (CA)	U	10.0 mmol kg ⁻¹	Promoted dissolution of uranium, mitigated oxidative stress	Hu et al. (2019)
4.	<i>Oryza sativa</i> L.	Hydroxyiminodisuccinic acid (HIDS)	Fe, As	0.25 mmol	increased Fe uptake and growth of rice seedling As phytoextraction	Rahman et al. (2009)
5.	<i>Ricinus communis</i> L.	citric acid (CA)	Cr	5 mmol kg ⁻¹	Enhanced Cr absorption, improved plant growth	Ali et al. (2022)
6.	<i>Solanum nigrum</i>	N, N-bis glutamic acid (GLDA)	Cd	3.0 mmol kg ⁻¹	Improved Cd absorption	Teng et al. (2022)
7.	<i>Zea mays</i>	aspartate dibutyric acid ether (AES)	Cd	6.0 mmol kg ⁻¹	Increased biomass above the ground	Yang et al. (2021)




Fig. 2. Metal uptake and transport: Plants root cells take the heavy metals from the soil and through the symplast pathway. The heavy metals are transported to aerial parts of the plants using various types of transporters.

turnover (Barber-Zucker et al., 2019).

5.3. NRAMP family of proteins

The NRAMP family, which transports metal cations into the cytoplasm of plants, primarily functions to maintain iron homeostasis in *A. thaliana*, with several NRAMP genes in plants being expressed under Fe deficiency, indicating their role in nutrition, and are also expected to be present in intracellular membranes like the plastid envelope and vacuolar membrane (Thomine and Schroeder, 2004). Whereas, over-expression or disruption in *Arabidopsis* NRAMP genes results in alterations in sensitivity towards Fe or Cd. Transcriptomic analysis indicated *AtNRAMP3* upregulated at the time of leaf senescence, which implies its significant role in remobilization of HMs during leaf senescence (Zhu et al., 2021). *AtNRAMP1* demonstrates high affinity towards Mn and essential for a *Arabidopsis* growth in low Mn condition (Cailliatte et al., 2010). In addition, *A. halleri* and *T. caerulea*, which are metal hyperaccumulating species, have significantly higher expression levels of *NRAMP3* and *NRAMP4* compared to their homologs in *A. thaliana* (Oomen et al., 2009).

5.4. HMAs

These HMs transporters use ATP molecules to efflux cytoplasmic cations through biological membranes. On the basis of phylogenetic studies, HMAs cluster is subdivided into two classes, namely the Pb/Cd/Co/Zn type and Ag/Cu type (Wang et al., 2011). Therefore, *AtHMA4* code for Zn/Co/Cd/Pb P_{1B}-type ATPase is the first HMAs that was cloned and characterized in plants (Colangelo and Guerinot, 2006). Whereas, in *A. thaliana*, *AtHMA2* and *AtHMA4* confers translocation of Zn utilizing the xylem loading process and are located on plasma

membrane (Verret et al., 2004). *AtHMA4* homologs in *A. halleri* and *T. caerulea* possibly execute a similar function, resulting in a dramatic increase of *AhHMA4* and *TcHMA4* expression in roots which accounts for the influx/efflux of Zn in accumulator species (Krämer et al., 2007). *T. caerulea* and *A. halleri* utilize *HMA4* for the hyper-accumulation of Cd as well. Moreover, it has been affirmed that *HMA4* works as an efflux pump in roots and leaf cells conferring tolerance against Cd and Zn, which was confirmed by its heterologous expression in yeast (Colangelo and Guerinot, 2006).

6. Metal sequestration in vacuole by tonoplast

The vacuole in higher plants serves as a storage site for minerals, metabolites, and toxicants, while also playing a key role in long-term metal transport through its ability to dynamically modulate vacuolar sequestration capability (VSC), where it neutralizes and adjusts sequestration capacity in response to changing environmental conditions through the interaction of metal ion chelators and tonoplast-localized transporters (Clemens and Peršoh, 2009). Moreover, it paves the way for a successful biofortification of key nutrients or phytoremediation of the contaminants (Brunetti et al., 2011). Metal transporters in plants located in the tonoplast are imperative for the regulation of vacuolar sequestration capability and consequently the allocation of metal in different plant parts (Martinoia et al., 2012). When exposed with excess metal, metal transporters—identified in regulating metal sequestration in vacuoles—display increased cellular response. The enhanced expression of genes such as *MTP3* (Metal Transporter 3) (Arrivault et al., 2006), *VIT2* (Vacuolar Iron Transporter 2) (Zhang et al., 2012), *ZIF1* (Zinc- Induced Facilitator 1) (Haydon and Cobbett, 2007), *CAX4* (Cation Exchanger 4) (Mei et al., 2009), subsequently causes expansion in the VSC. The gene expression levels of *MTP1* increased

significantly in various metal hyperaccumulators (Ricachenevsky et al., 2013).

A subclass of chelators, called PCs are potentially very crucial in HMs detoxification by formation of metal complexes which thereby get trapped in the vacuoles. NA a different metal chelator that is primarily found in higher plants, changes the VSC of Fe and Zn and consequently their respective distribution. The root biomass of *A. helleri* contains significant levels of NA which allows the metal allocation of Zn to shoots (Deinlein et al., 2012). The VSC regulation is mediated by chelators and can be categorized into two categories: 1) The metal chelators tend to translocate to vacuoles which leads to enlargement of VSC of certain metals. 2) As a consequence of metal chelation, the vacuolar sequestration is reduced, hence, reducing the VSC of certain metals (Deinlein et al., 2012). In some Ni hyper-accumulators, it has also been shown that Histidine (His) has a prominent role in detoxification and translocation of Ni. It is concluded in a study that Ni in the form of Ni-His complex when supplied increases the rate of uptake in tonoplast vesicles of hyper-accumulator *T. arvensis*. According to their theory, His deposition in the roots of *T. caerulea* declines Ni's cytosolic chelation potential, thereby, enhancing Ni transport and its hyperaccumulation in shoots (Maestri et al., 2010).

7. HMs sequestration mechanisms in plants

HMs accumulation in multicellular plants involves a complex interplay of metal uptake, intracellular metal binding, and detoxification processes, which is further complicated by cell proliferation and tissue-specific mechanisms that can disrupt the balance of essential metal homeostasis and redox systems, leading to increased metal-induced stress (Foyer and Noctor, 2005). Various processes are involved in the deposition of metals in plants, including the mobilization and uptake of HMs from the soil, their sequestration through the creation of metal complexes such as chelators and storage in vacuoles. Metal transport can occur via apoplastic or symplastic pathways to reach shoots, aided by effective xylem loading. Once transported to leaves, metal ions are confined within tissue layers, and ultimately, deposited metals are stored in structures like trichomes (Clemens et al., 2002). The absorption and movement of metals in plant leaves are aided by transporters and metallochaperones. Several mechanisms for HM sequestration in plants are detailed below.

7.1. Phytochelatin (PC's)

PC's are low molecular weight peptides containing thiol group. GSH serves as a PC precursor where GSH is acted upon by phytochelatin synthetase (Hasanuzzaman et al., 2017). Reports have found that PC's play a part in plant defense against metal-related challenges and other stress conditions like salt, intense heat, UV radiation, herbicides (Zagorchev et al., 2013). Additionally, in HMs stress diagnosis, PCs are used as biomarkers. Whereas, HMs like Hg, gold (Au), Cu, silver (Ag), Cd and Zn induce biosynthesis of PCs. Further it has been observed that Cd is the strongest inducer in *Brassica napus* (Carrier et al., 2003). Moreover, PCs form complexes with metals, they are transported to the vacuoles. These complexes enable the extraction of HMs from cytoplasmic HMs sensitive enzymes. The transportation of metal-PC complexes from cytosol to vacuole is mediated by transporters of tonoplast which can be either ATP-dependent ABC transporters or metal/H⁺ antiporters (Kang et al., 2011). The sulfide and sulfite ions present in vacuoles are integrated in PC- HM complex, hence providing stability and stabilizing pH when dissociation of metal occurs. Roughly 90 % of total Cd of mesophyll cells remain in vacuoles in case of Cd-hyperaccumulator *Thlaspi caerulescens*. Vacuolar sequestration was therefore postulated in these plants as a key internal detoxifying mechanism for Cd in leaves (Ma et al., 2005).

7.2. Metallothionines (MTs)

MTs are cysteine-rich, low molecular weight proteins that sequester HMs by forming metal-thiolate clusters, and are found abundantly in eukaryotes (including mammals, plants, and fungi) as well as prokaryotes, differing from PCs, which are enzymatically produced, whereas MTs result from the mRNA translation process (Verkleij et al., 2003). MTs in plants show affinity for a broader range of HMs, including As and Zn, playing a crucial role in detoxifying these metals by sequestering them, maintaining intracellular metal ion homeostasis, and regulating HM transport (Guo et al., 2013). They actively participate in cellular functions such as ROS scavenging, redox maintenance, cell proliferation, and DNA repair, while multiple factors including osmotic stress, nutritional deficiency, hormone secretion, naturally occurring or induced tissue senescence, infections, or wounds can lead to the expression of MTs (Kumar et al., 2017). MTs are classified into four types on the basis of order of Cysteine residues: Type 1 – root expressing MTs, Type 2-shoot expressing MTs, Type 3 and Type 4-leaf and developing seeds expressing MTs. Further these subgroups are divided into isoforms. In a study of *Arabidopsis* it was depicted that MT isoforms (Type 1a, 2a, 2b and 3) are responsible for copper chelation whereas Type 4a and 4b MT isoforms function as Zinc binders (Grennan, 2011).

7.3. Stress proteins

Upon subjection to metal toxicity, plants stimulate novel proteins production. These proteins, commonly known as "Stress proteins," belongs to heat shock protein family (HSPs). HSPs are molecular transporters and facilitate correct protein folding and assembly. In various plant species upon exposure to Cd, the DNA isolated from cells subjected to Cd stress produces certain mRNA transcripts, that control biosynthesis of stress related proteins. Cys-rich membrane protein (*Arabidopsis* plant cadmium resistance, (*AtPCRS*) were identified in *A.thaliana*. Its homologs were also found in *Oryza sativa* and *Solanum lycopersicum*, that provides resistance to Cd exposure (Song et al., 2004). The expression of proteins that are related to pathogenesis also called as PR proteins, which show metal selectivity in plants when metal toxicity increases. For instance, when exposed to Cd, Zn, and Cu, *Lupinus luteus* plants produce PR-10 family proteins that are associated to pathogenesis (Przymusiński et al., 2004).

7.4. Proline

Proline is the most ubiquitous 5-carbon α -amino acid, functioning like a suitable metabolic osmolyte, free radical scavenger and antioxidant. Elevation in proline levels is a non-enzymatic response of higher plants towards biotic and abiotic stress conditions like HMs, oxidative stress and increased salinity (Szabados and Savaouré, 2010). HMs caused accumulation of proline in plants as a outcome of water imbalance, brought on by metal overload, rather than direct HM stress, hence proline working as an osmoregulator or osmoprotectant (Clemens, 2006). Primarily, proline acts as a ROS scavenger by hydroxyl radicals detoxification and singlet oxygen quenching. Experiments on *Brassica* sp. and lemongrass have demonstrated proline accumulation in roots after exposure to excessive concentration of HMs i.e., Pb, Cd, Hg and Cd (John et al., 2009; Handique and Handique, 2009).

8. Tolerance and toxicity to HMs at the cellular level

The exposure to HMs in plants can lead to genotoxic and oxidative stress responses, resulting in cellular damage and disruption. To counteract these harmful effects, plants have evolved two defense mechanisms: antioxidant enzymes (for example ascorbate peroxidase (APX), SOD, GR, and CAT) and the accumulation of various antioxidants like carotenoids, ascorbate, phenylpropanoids, GSH, proline, and alkaloids (Bhaduri and Fulekar, 2012). For example, it has been observed in

Sainfoin (*Onobrychis vicifolia*) the activity and function of three enzymes namely SOD, CAT and GPX increased in the leaves after contaminated with HM's like Cu and Pb (Beladi et al., 2011). However, HMs like Cd can still cause oxidative damage, disrupt lipid membranes, and contribute to reduced agricultural yield (Cho and Seo, 2005). Furthermore, ROS, despite their damaging potential, also play important roles as signalling messengers in various cellular processes, including cell development, differentiation, apoptosis, and stress response. Nevertheless, ROS-induced oxidative stress can lead to DNA damage and genome instability, further impacting plant development and agricultural productivity (Markkanen et al., 2012) (Fig. 3).

9. Environmental Implications of HMs

There is a widespread concern over HM contamination in both aquatic and terrestrial environments. HMs, originating from various sources (mining operations, industries, domestic sewage and agricultural runoff), are persistent and can accumulate in ecosystems, posing risks to organisms and human health (Singh and Steinnes, 2020). They enter the food chain, particularly affecting aquatic habitats, where sediments serve as reservoirs (Ali Azadi et al., 2018). Changes and reduction in pH can release HMs into water because reduction in pH causes a competition between metal ions and H^+ for binding sites in sediments which may dissolve the metal complexes into liberated metal ions in the water column (Nowrouzi et al., 2014). Soil contamination, often resulting from the use of fertilizers and various pollutants, highlights the importance of understanding the bioavailability of HMs in the soil to accurately assess their effects on ecosystems (Ali et al., 2019).

10. Nano-bioremediation of HMs contaminations

Significant amounts of pollutants have been dispersed around the world as a result of the world's rapid population increase, expanding industries, and developing urbanization. The main objective of scientific research is to create environmentally sustainable technologies that are

also commercially feasible. Bioremediation has proved to be a more efficient, environmentally beneficial, and cost-effective technique (Patel et al., 2022). It exploits microbial ability to remove or degrade environmental pollutant (Sharma et al., 2024). Though, this process is not suitable for such sites which are highly contaminated with HMs and salts because they are harmful for microbial survival. So, there is a need for exploring innovative technologies which are highly beneficial compared to existing methods. The usage of nanomaterials is expanding rapidly to address environmental concerns as a result of the outstanding advances in the field of nanotechnology. There are numerous ways to create nanoparticles (NPs), including sonication, pulse wire discharge, radiation, spray pyrolysis, and electro-deposition. However, each of these approaches uses a different set of chemicals, is costly, and creates a range of hazardous byproducts (Abdel-Shafy and Mansour, 2018). It is therefore necessary to develop more affordable and environmentally friendly methods for synthesizing NPs. The synthesis of NPs through biological processes is more advantageous for bioremediation processes because it is both economical and environmentally beneficial (Sivaraj et al., 2015). During the biological or green synthesis of NPs, metabolites of living organisms reduce metal ions to stable nanomaterials by using NADPH-dependent enzymes via an electron shuttle enzymatic metal reduction mechanism (El-Seedi et al., 2019). These metal NPs or nanomaterials (NMs) minimize pollution by absorbing a sizable amount of contaminants owing to its high surface tension and surface area features, respectively. The bioremediation process is accelerated by these nano-sized form of particles, which are able to reach contaminated areas more readily (Pete et al., 2021). Nanomaterials synthesized by fungi, bacteria, yeast and actinomycetes are widely used in industry, environmental remediation, remediation of uranium, solid waste remediation and HM remediation. Amongst all the microorganisms, fungi have been experimentally observed as the most momentous producer for NP synthesis, because fungal mycelia can tolerate harsh conditions and high pressure as compared to bacteria and plant extracts (Adebayo et al., 2021). Further fungi can grow easily and the extracellular secretions of fungi are composed of various reactive proteins which are easy to handle in




Fig. 3. Mechanisms involved in heavy metal sequestration: When heavy metals enter the plant system via transporters present in roots, they are distributed to whole plant body via symplastic and apoplastic mechanisms. At cellular level, these heavy metals enter the cell cytoplasm through plasma membrane. Upon reaching there, they tend to increase the ROS production that causes DNA damage, membrane disruption, increase in antioxidant enzyme synthesis. Also the expression of heavy metal responsive genes and enzymatic activity genes are upregulated in nucleus. Heavy metals forms complexes with stress proteins, phytochelatins, metallothionins and proline that are further transported to vacuole by ATPase transporters where heavy metal sequestration takes place.

downstream processing. It was reported that *Verticillium* sp. and *Fusarium oxysporum* on exposing with aqueous AgNO₃ solution produced intracellular and extracellular Ag NPs respectively (Khan et al., 2018; Yassin et al., 2021). Amongst all NPs, Ag-NPs, iron- NPs, carbon- NPs, nanoscale zero-valent iron (nZVI) are abundantly used in bioremediation experiment (Yadav et al., 2017). When utilized on heavily polluted sites containing persistent pollutants like Cr and As, zero valent nanoscale metal ions like Fe and bimetallic NPs like Fe/Ni and Fe/Pd have shown good results since they have very low toxicity and cost effective (Thomé et al., 2015). HMs pollution affects quality of aquatic bodies and soil. Although HMs are non-degradable, these can be modified by methylation, complexation, and absorption. Mostly the HMs polluted sites are found to be a good reservoir of various HMs tolerant microbes. Therefore, microbes are used to remediate and detoxify some metals and reduce metal pollution through bioremediation (Igiri et al., 2018). Fungi of the genera *Penicillium* and *Rhizopus* have been recognized to remove toxic HMs from solutions (Ghosh et al., 2023). As per reports, *Aspergillus niger* assist in removing lead and chromium ions from aqueous solutions (Li et al., 2016; Xu et al., 2021). Fungus based nanomaterials are more effective in remediation of various metal ions from aqueous and soil ecosystems. Fungi have biochemical and ecological capacity to remove environmental toxins associated with HM, radio nuclides and metalloids through chemical modifications (Fomina and Gadd, 2014). Agricultural pollution, a global concern, stems from practices such as fertilizer and pesticide application, soil tillage, and introduces harmful pollutants like nitrates, phosphates, pesticides, and HMs into the environment, posing significant hazards (Khan et al., 2018). Nano bioremediation is one such approach that exploits advantage of nanotechnology or nanomaterial along with bioremediation benefits. This combination of biotechnology and nanotechnology has proven to be a successful method for cleaning up pollutants. Additionally, it can be used in conjunction with bacteria or microbial enzymes.

11. Phytoremediation

Phytoremediation is the process whereby plants and their associated soil microbes collinearly work to reduce the concentrations or toxic effects of contaminants in the environment. It encompasses various methods, including phytoextraction, phytodegradation, phytovolatilization, and phyto-stabilization. Except for phytodegradation, which is employed only in the degradation of organic pollutants to non-toxic compounds, the other three methods can be employed for the removal of heavy metals from the soil (Kukreja and Goutam, 2012). Microbe-assisted phytoremediation is the decontamination of pollutants in the rhizosphere carried out by an interaction of plants and microbes. It can be carried out by one of the two ways: either by using natural existing population of microbes in the soil or by inoculation with the desired microbes that are able to break down pollutants (Asemoloye et al., 2019). To facilitate rhizospheric bacterial colonization, plants roots secrete copious amounts of bioactive chemicals such as root exudates (enzymatic and non-enzymatic) including flavanones, proteins, carbohydrates, enzymes, amino acids, nucleotides, organic acids and phenolics which are utilized by bacteria (Al-Ameeri and Al Sarawi, 2017). Bacteria release a variety of substances, including organic acids, polysaccharides, metal chelators (siderophores), organic acids, and biosurfactants. These substances all help in HMs detoxification by increasing bioavailability and reducing soil pH, which limits HMs mobility (Rajkumar et al., 2012). *Micrococcus luteus*, which is resistant to As, engaged in siderophore production, nitrogen sequestration, and phosphate solubilization, resulting in increased grapevine yield and reduced As toxicity, even under high arsenic concentrations (Pinter et al., 2017). The total dried weight and leaf surface area of pepper plants inoculated with AM fungus increased in response to high Cu in soil (Ruscitti et al., 2017). Mobilization and assimilation of Hg and As improved in *Brassica juncea* and *Lupinus albus* growing in soils contaminated with these metals when thiosulfate was added together with HM

tolerant plant growth promoting (PGP) microorganisms (Franchi et al., 2017). AMF and rhizobacteria release cysteine-rich peptide molecules referred as MTs that are structurally identical to PCs and aid in the accumulation of HMs (Miransari, 2011). The application of AMF *Glomus mosseae* to the *Schedonorus arundinaceus* (tall fescue) plants lowers the movement of Ni to the plant and enhances the amount of MT and ABC gene expression (Shabani et al., 2016). Rhizosphere microbes produce extracellular polymeric substances (EPS) that contain a significant amount of anion functional groups, which aid in the biosorption and subsequent removal or recovery of metals, for example *Trichoderma harzianum* and *Bacillus subtilis* fungal-bacteria biofilms were able to reduce Zn, Pb and Cd assimilation in potato tissue which can be employed as viable solution in rhizoremediation of HMs (Henagamage et al., 2022). In one experiment, *Sesbania sesban* legume plant seedlings were subjected to HMs like Cr and Cd with the bacterium *Bacillus anthracis* PM 21. It was found that this bacterium promoted the growth of plants biomass by ACC-deaminase expression, Indole-3-acetic acid (IAA), exopolysaccharide production and elevated functions of antioxidants in HMs environments (Ali et al., 2021).

12. Genetic modified plant in HMs remediation

Utilizing transgenic plants for HMs removal is providing new direction in environmental clean-up. HMs can be removed by inserting gene specific promoters in plants enhancing the transcription of existing genes. For example, the overexpression of the *Vicia sativa* caffeoyl-CoA O-methyltransferase (*VsCCoAOMT*) gene in *Arabidopsis thaliana* showed the potential of genetically modified (GM) crops in phytoremediation applications, as these transgenics exhibit enhanced Cd tolerance and increased biomass production (Xia et al., 2018). Similarly, HMs ATPase (HMA5) is a Cd exchanger channel, present in plants for cadmium exchange between shoots and roots. Whereas, PTHMA5 isolated from *Populus trichocarpa*, when overexpressed in *Nicotiana tabacum* leads to Cd accumulation suggesting its potential in phytoremediation (Wang et al., 2018). Additionally, Endophytic bacterial genes are quite useful for transforming plants to enhance their metal absorption. Endophytic bacterial genes, such as *CUP* and *bphC*, were used to produce first transgenic nettle (*Urtica dioica*) plants, which demonstrated a 33 % reduction in polychlorinated biphenyls and an 8 % reduction in HMs (Viktorova et al., 2016). Plants possess inherent detoxification mechanisms, notably glutathione (GSH) homeostasis; a potential redox compound, that links intracellular bioenergetic signaling pathways. Gamma- glutamyl-cysteine synthetase- glutathione synthetase gene (Gamma-GCS-GS) is recently discovered in *Sulfolobus thermophiles* that has high tolerance level for HMs (Liedschulte et al., 2010). Moreover, glutathione synthetase gene overexpressed in *Beta vulgaris* (sugar beet) had high ability to retain HMs like Cd, Cu, Zn, and showed high GSH and PC activities upon exposure to HM stress in comparison to wild type (Liu et al., 2015). There are also certain functional genes along with metal transporter proteins that could be targeted for remediation purposes. Furthermore, Nahar et al. (2017), successfully overexpressed the arsenic reductase2 gene (*AtACR2*) in tobacco isolated from *A. thaliana* leading to high As accumulation in roots whereas lower concentration in the shoots. This indicates that the *ACR2* gene may be useful in As phytoremediation and in the production of transgenic crops. A list of genes involved in heavy metal uptake in plants is given in Table 4. However, the deployment of GM crops in environmental remediation raises biosafety concerns, ecological risks, and regulatory challenges. In addition, biosafety concerns include the potential for horizontal gene transfer to non-target organisms, the development of herbicide resistance in weeds due to gene flow, and the unintended effects of transgene expression on plant metabolism and ecological interactions. Ecological risks include the potential for altered plant-microbe interactions, changes in soil microbial communities, and the disruption of ecosystem services. Furthermore, regulatory frameworks for GM crops vary significantly across countries, posing

Table 4
Genes of plants involved in HM uptake.

Sl. No	Genes	Plants	HM	Refs.
1.	<i>Nramp</i>	<i>Arabidopsis</i> sp., <i>Oryza sativa</i>	Cd and divalent metals	Belouchi et al. (1995); Alonso et al. (1999); Thomine et al. (2003)
2.	<i>AtNRAMP6</i>	<i>Arabidopsis thaliana</i>	Fe	Li et al. (2019)
3.	<i>AhZIP6</i>	<i>Arabidopsis halleri</i>	Cd	Spielmann et al. (2020)
4.	<i>CAD1</i>	<i>Arabidopsis</i> sp.	Cd	Ha et al. (1999)
5.	<i>AuRT1</i>	<i>Arabidopsis</i> sp.	Fe	DalCorso et al. (2008)
6.	<i>AtATM3</i>	<i>Arabidopsis</i> sp.	Pb and Cd	Kim et al. (2006)
7.	metallothionein (MT) genes	<i>Arabidopsis thaliana</i>	Cu	Khandekar & Leisner (2011)
8.	<i>COPT/Ctr</i> (<i>COPT1-7</i>)	<i>Arabidopsis thaliana</i>	Cu	Jan et al. (2016)
9.	<i>Cdr3</i>	<i>Arabidopsis</i> sp.	Cd	Sethy & Ghosh (2013)
10.	<i>VsCCoAOMT</i>	<i>Arabidopsis</i> sp.	Cd	Xia et al. (2018)
11.	<i>TcHMA4</i>	<i>Arabidopsis thaliana</i> <i>Thlaspi caerulescens</i>	Cd, Zn, Pb, and Cu	Papoyan & Kochian (2004)
12.	<i>TaHMA2</i>	<i>Arabidopsis</i> sp.	Zn and Cd	Qiao et al. (2018)
13.	<i>LuABCC9</i>	<i>Arabidopsis</i> sp.	Cd	El-Sappah et al. (2023)
14.	<i>SLMJ524</i>	<i>Arabidopsis</i> sp.	Cd	Li et al. (2022)
15.	<i>GGCT2</i>	<i>Arabidopsis</i> sp.	As	Paulose et al. (2013)
16.	<i>CaGrx</i>	<i>Arabidopsis thaliana</i>	As, Cr and Cd	Kumar et al. (2020)
17.	<i>AzMT2</i>	<i>Azolla filiculoides</i>	Ni and Cd	Schor-Fumbarov et al. (2005)
18.	<i>NRAMP</i>	<i>Brassica napus</i>	Zn, Mn, Cd and Pb	Meng et al. (2017)
19.	<i>GSHI</i> and <i>GSHII</i>	<i>Brassica juncea</i>	Cd	Liang Zhu et al. (1999)
20.	<i>HMA4</i>	<i>Brassica juncea</i>	Cd	Wang et al. (2019)
21.	<i>BjHMA4</i>	<i>Brassica juncea</i>	Cd	Wang et al. (2019)
22.	<i>BjYSL7</i>	<i>Brassica juncea</i>	Cd	Wang et al. (2013)
23.	<i>BjHO-1</i>	<i>Brassica juncea</i>	Hg	Li et al. (2012)
24.	<i>CsMTP8.2</i>	<i>Camellia sinensis</i>	Fe, Mn and Zn	Zhang et al. (2020)
25.	<i>CaMYB</i>	<i>Capsicum annuum</i>	Cd, Cu, and Pb	Xie et al. (2022)
26.	<i>GmHMA3</i>	<i>Glycine max</i>	Cd	Wang et al. (2018)
27.	<i>GmaMTPs</i>	<i>Glycine max</i>	Zn	El-Sappah et al. (2023)
28.	<i>HvHMA1</i>	<i>Hordeum vulgare</i>	Cu	Bernal et al. (2007)
29.	<i>LmSAP</i>	<i>Lobularia maritima</i>	Cd, Cu, Mn and Zn	Saad et al. (2018)
30.	<i>HMA4</i>	<i>Liriodendron tulipifera</i>	Zn,	Adams et al. (2011)
31.	<i>PCS1</i> <i>HMA</i>	<i>Linum usitatissimum</i>	Cd	El-Sappah et al. (2023)
32.	<i>PvSR2</i>	<i>Nicotiana tabacum</i>	Cd	Chai et al. (2003)
33.	<i>TaPCS1</i>	<i>Nicotiana glauca</i>	Pb and Cd	Gisbert et al. (2003)
34.	<i>OshMA3</i> <i>OsMTP1</i> <i>OsABCC1</i>	<i>Oryza sativa</i>	Cd, Zn and As	Menguer et al. (2013); Sasaki et al. (2014); Song et al. (2014)
35.	<i>HMA</i> (<i>HMA1</i> to <i>HMA9</i>)	<i>Oryza sativa</i>	Zn, Pb and Cd	Takahashi et al. (2011)
36.	<i>OshMA2</i> and <i>OshMA3</i>	<i>Oryza sativa</i>	Cd and Cu	Kim et al. (2014)
37.	<i>OsMTP1</i>	<i>Oryza sativa</i> IR64	Cd	Das et al. (2016)
38.	<i>OsMTP11</i>	<i>Oryza sativa</i>	Mn	Zhang and Liu (2017)
39.	<i>H655</i> and <i>H767</i>	<i>Oryza sativa</i>	Hg	Wang et al. (2023)
40.	<i>RsMYB1</i>	<i>Petunia</i> sp.	Cu, Zn, Mn and Cr	Ai et al. (2018)

Table 4 (continued)

Sl. No	Genes	Plants	HM	Refs.
41.	<i>ScMT2-1-3</i> ,	<i>Saccharum officinarum</i>	Cd and Cu	Guo et al. (2013)
42.	<i>SpHMA3</i>	<i>Sedum plumbizincicola</i>	Cd	Liu et al. (2017)
43.	<i>StMATE</i>	<i>Solanum tuberosum</i>	Cu	Huang & He (2021)
44.	<i>TcHMA3</i>	<i>Thlaspi caerulescens</i>	Cd	Ma et al. (2005); Ueno et al. (2011)
45.	ZIP family (<i>ZAT1</i> , <i>ZAT2</i> , <i>ZAT3</i>)	<i>Thlaspi caerulescens</i>	Cd, Zn and Mn	(Pence et al. (2000); Lombi et al. (2002)
46.	<i>ZmHMA3</i>	<i>Zea mays</i>	Cd	Tang et al. (2021)
47.	<i>ZmST1</i>	<i>Zea mays</i>	Cd, Zn, and Cu	Nocito et al. (2006)
48.	<i>ZjMT</i>	<i>Ziziphys jujuba</i>	Cd	Li et al. (2016)

challenges for the development and commercialization of these technologies. Public perception and acceptance of GM crops are also critical factors that can influence their adoption. Therefore, comprehensive risk assessments, stringent regulatory oversight, and transparent communication with stakeholders are essential to ensure the safe and responsible application of GM crops in HMs remediation. Whereas, future research should focus on developing strategies to mitigate these risks, such as the use of gene containment technologies and the development of site-specific and self-limiting transgenes. Additionally, the long-term ecological impacts of GM crops deployment should be thoroughly evaluated through field trials and monitoring studies.

13. Microbial based bioremediation of HMs

Heavy metal detoxification via microbial-based bioremediation has become popular as an cost effective and eco- friendly technique. This method uses of the innate ability of various microbes, such as bacteria, fungi, or algae, to degrade certain heavy metals from polluted areas. Since these microbes are flourishing in places affected by heavy metals, they are ideal candidates for microbial bioremediation due to their distinct metabolic characteristics. Microbial bioremediation can occur in several ways, including biosorption, bioaccumulation, biotransformation, and biomineralization (Tang et al., 2024; Priya et al., 2022; Pande et al., 2022).

According to Tripathi et al. (2023), biosorption is the process by which metal ions attach to microbial cell walls, containing functional groups with a high affinity for metal ions, such as carboxyl, amino, and hydroxyl. As this process is very effective and does not require the metabolic input from the cell so the dead cells can also be utilised for the process. A study demonstrated that dead cells of *Bacillus cereus* RC-1 shows higher Cd(II) biosorption capacity than live cells (Huang et al., 2013). Recent studies have highlighted the biosorption potential of green and red algae for heavy metals such as Pb, Cu, Cd, Zn, and Cr (Romera et al., 2007; Sari and Tuzen, 2008). However, despite their ability to remove these metals from aqueous solutions, their performance is significantly lower compared to brown algae (such as *Sargassum* sp.) (He and Chen 2014).

In contrast to absorption, bioaccumulation is the process by which microbes absorb and accumulate metals within their cells using cellular energy with the involvement of metal-binding proteins and an efflux mechanism (Mishra and Malik, 2013). Microbes are also involved in the enzymatic conversion of hazardous metals into less dangerous or immobile forms by the process of biotransformation. For example, several bacteria have the ability to decrease the bioavailability of deadly hexavalent chromium (Cr⁶⁺) by transforming it into less hazardous trivalent form Cr³⁺ (Juwarkar and Yadav, 2010).

Biomineralization is another process in which metals are immobilized in stable forms within the soils through microbial metabolic activities (Li et al., 2023). Microbes by lowering the toxicity, mobility, and

bioavailability of HMs, help to lessen their negative impacts on the environment. There are various microbial species that has shown their remarkable capabilities in the HMs bioremediation. For example, strains of *Pseudomonas* show resistance towards the Cd as they are involved in the absorption of the cadmium (Chellaiyah, 2018; Naz et al., 2016). *Bacillus* species are useful in the metal uptake and the transformation in the HMs contaminated environments (Wróbel et al., 2023; Butterfield et al., 2016). Whereas, *Aspergillus niger* and *Penicillium* exhibited the bio-sorption of HMs due to the nature of their cell walls that have the metal binding efficiency (El-Mahdy et al., 2021). Although, Use of microbe in a consortium can be utilized to detoxify several metals at once so it could be helpful in lethal contamination. Additionally, reduced levels of secondary pollutants, cost effective and recovery of metal for the industrial use are just a few benefits of microbial based bioremediation.

However, there are some limitations in the broad implementation of the microbial based bioremediation. There are several environmental elements that determine the effectiveness of microbial activity include pH, temperature, and the presence of other pollutants (Ayilara and Babalola, 2023). In addition, some HMs are poisonous to microbes, and this may restrict their ability to absorb and transform the metals into less harmful form. These constraints can be overcome by using adaptive evolution and genetic engineering practices to enhance metal-removal effectiveness and microbial resistance. To overcome these constraints, current studies have concentrated on using adaptive evolution and genetic engineering methods to improve metal-removal effectiveness and microbial resistance.

14. Conclusion and future outlook

The increasing concentrations of HMs in the environment poses a significant threat to the health and safety of the living organisms. As a result, it is crucial for plants to develop adaptive strategies that enhance their tolerance to HM stress and reduce the risk of HM transfer into the food chain. HMs appears to have diverse mechanisms to exert toxicity, while plants, in turn, deploy diverse defence responses to counteract their effects. When exposed to excessive HMs, plants exhibit distinct molecular and biochemical tolerance strategies. These strategies primarily aim to prevent the accumulation of toxic metals within the cells by the mechanisms such as apoplastic entrapment, chelation of metals by cytosolic ligands, and sequestrations into vacuole. Furthermore, the exogenous treatment of different organic or inorganic compounds has emerged as a promising approach to mitigate HM toxicity in plants. Exploring the interactions between these compounds and the plant systems, as well as their role in ameliorating HM stress, presents a potential area of future research.

CRedit authorship contribution statement

Himani Agarwal: Writing – original draft. **Divya Chaudhary:** Writing – original draft. **Himanshi Aggarwal:** Writing – original draft. **Chhavi Karala:** Writing – original draft. **Niharika Purkait:** Writing – original draft. **Neha Sharma:** Software, Investigation. **Arti Mishra:** Software, Methodology. **Vaibhav Mishra:** Software, Methodology. **Ajay Kumar:** Software, Methodology. **PrashantKumar Singh:** Visualization, Software, Investigation. **Laurent Dufossé:** Writing – review & editing, Supervision, Conceptualization. **NaveenChandra Joshi:** Writing – review & editing, Supervision, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

DST-SERB start-up grants provided funding for this research to Naveen Joshi (File no SRG/2020/002237).

Data availability

Data will be made available on request.

References

- Ali, S., Mfarrej, M.F.B., Rizwan, M., Hussain, A., Shahid, M.J., Wang, X., Nafees, M., Waseem, M., Alharby, H.F., 2022. Microbe-citric acid assisted phytoremediation of chromium by castor bean (*Ricinus communis* L.). *Chemosphere* 296, 134065.
- Abdel-Shafy, H.I. & Mansour, M.S. (2018). Green synthesis of metallic nanoparticles from natural resources and food waste and their environmental application. *Green Met. Nanoparticles Synth. Charact. Their Appl.*, 321–385.
- Adams, J.P., Adeli, A., Hsu, C.Y., Harkess, R.L., Page, G.P., Depamphilis, C.W., Schultz, E. B., Yuceer, C., 2011. Poplar maintains zinc homeostasis with HM genes HMA4 and PCS1. *J. Exp. Bot.* 62 (11), 3737–3752. <https://doi.org/10.1093/jxb/err025>.
- Adebayo, E.A., Azeez, M.A., Alao, M.B., Oke, A.M., Aina, D.A., 2021. Fungi as veritable tool in current advances in nanobiotechnology. *Heliyon* 7 (11).
- Ai, T.N., Naing, A.H., Yun, B.W., Kim, C.K., 2018. Overexpression of RsMYB1 enhances anthocyanin accumulation and HM stress tolerance in transgenic petunia. *Front. Plant Sci.* 9, 1388.
- Al-Ameeri, D.K., Al Sarawi, M., 2017. Rhizoremediation of contaminated soils by comparing six roots species in Al-Wafra, State of Kuwait. *J. Bioremediation Biodegrad.* 8 (384), 2.
- Alonso, J.M., Hirayama, T., Roman, G., Nourizadeh, S., Ecker, J.R., 1999. EIN2, a bifunctional transducer of ethylene and stress responses in Arabidopsis. *Science* 284 (5423), 2148–2152. <https://doi.org/10.1126/science.284.5423.2148> (1979).
- Ali Azadi, N., Mansouri, B., Spada, L., Sinkakarimi, M.H., Hamesadeghi, Y., Mansouri, A., 2018. Contamination of lead (Pb) in the coastal sediments of north and south of Iran: a review study. *Chem. Ecol.* 34 (9), 884–900.
- Ali, H., Khan, E., Ilahi, I., 2019. Environmental chemistry and ecotoxicology of hazardous HMs: environmental persistence, toxicity, and bioaccumulation. *J. Chem.* 2019 (1), 6730305. <https://doi.org/10.1155/2019/6730305>.
- Ali, J., Ali, F., Ahmad, I., Rafique, M., Munis, M.F.H., Hassan, S.W., Sultan, T., Ifrikhar, M., Chaudhary, H.J., 2021. Mechanistic elucidation of germination potential and growth of *Sesbania sesban* seedlings with *Bacillus anthracis* PM21 under HMs stress: an *in vitro* study. *Ecotoxicol. Environ. Saf.* 208, 111769.
- Alloway, B.J., 2012. Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability, 22. Springer Science & Business Media.
- Annamalainathan, K., Joseph, J., Alam, B., Satheesh, P.R., Jacob, J., 2013. Seasonal changes in xylem sap flow rate in mature rubber plants. *J. Plant Crop* 41 (3), 343–349.
- Arrivault, S., Senger, T., Krämer, U., 2006. The Arabidopsis metal tolerance protein AtMTP3 maintains metal homeostasis by mediating Zn exclusion from the shoot under Fe deficiency and Zn oversupply. *Plant J.* 46 (5), 861–879. <https://doi.org/10.1111/j.1365-313X.2006.02746.x>.
- Asad, S.A., 2011. Interactions Between Heavy Metals and Glucosinolates as Defense Mechanisms in *Thlaspi Caerulescens* (Doctoral Dissertation. University of Nottingham).
- Asemoloye, M.D., Jonathan, S.G., Ahmad, R., 2019. Synergistic plant-microbes interactions in the rhizosphere: a potential headway for the remediation of hydrocarbon polluted soils. *Int. J. Phytoremediation* 21 (2), 71–83.
- Aslam, M., Aslam, A., Sheraz, M., Ali, B., Ulhassan, Z., Najeeb, U., Zhou, W., Gill, R.A., 2021. Lead toxicity in cereals: mechanistic insight into toxicity, mode of action, and management. *Front. Plant Sci.* 11, 587785. <https://doi.org/10.3389/fpls.2020.587785>.
- Ayilara, M.S., Babalola, O.O., 2023. Bioremediation of environmental wastes: the role of microorganisms. *Front. Agron.* 5, 1183691.
- AjM, B., 1989. Terrestrial higher plants which hyper accumulate metallic elements. A review of their distribution, ecology and phytochemistry. *Biorecovery* 1 (2), 81–126.
- Backer, A.J.M., Brooks, R.R., 1989. Terrestrial higher plants which hyper accumulate metallic elements. A review of their distribution, ecology and phytochemistry. *Biorecovery* 1 (2), 81–126.
- Bali, A.S., Sidhu, G.P.S., 2021. Arsenic acquisition, toxicity and tolerance in plants-from physiology to remediation: a review. *Chemosphere* 283, 131050.
- Barber-Zucker, S., Hall, J., Mangapuram, S.V., Kass, I., Kolusheva, S., MacMillan, F., Zarivach, R., Henn, A., 2019. Metal binding to the dynamic cytoplasmic domain of the cation diffusion facilitator (CDF) protein MamM induces a 'locked-in' configuration. *FEBS J.* 286 (11), 2193–2215.
- Beladi, M., Habibi, D., Kashani, A., Paknejad, F., Nooralvandi, T., 2011. Phytoremediation of lead and Copper by Sainfoin (*Onobrychis vicifolia*): role of antioxidant enzymes and biochemical biomarkers. *Am. Eurasian J. Agric. Environ. Sci.* 10 (3), 440–449.
- Belouchi, A., Cellier, M., Kwan, T., Saini, H.S., Leroux, G., Gros, P., 1995. The macrophage-specific membrane protein nramp controlling natural resistance to infections in mice has homologues expressed in the root system of plants. *Plant Mol. Biol.* 29, 1181–1196. <https://doi.org/10.1007/BF00020461>.
- Bernal, M., Testillano, P.S., Alfonso, M., Carmen Risueño, M., Picorel, R., Yruela, I., 2007. Identification and subcellular localization of the soybean copper P1B-ATPase

- GmHMA8 transporter. *J. Struct. Biol.* 158, 46–58. <https://doi.org/10.1016/j.jsb.2006.10.016>.
- Bonanno, G., Giudice, R.L., 2010. HM bioaccumulation by the organs of *Phragmites australis* (common reed) and their potential use as contamination indicators. *Ecol. Indic.* 10 (3), 639–645.
- Bhaduri, A.M., Fulekar, M.H., 2012. Antioxidant enzyme responses of plants to HM stress. *Rev. Environ. Sci. Bio/Technol.* 11, 55–69.
- Brunetti, P., Zanella, L., Proia, A., De Paolis, A., Falasca, G., Altamura, M.M., Sanita di Toppi, L., Costantino, P., Cardarelli, M., 2011. Cadmium tolerance and phytochelatin content of *Arabidopsis* seedlings over-expressing the phytochelatin synthase gene *AtPCS1*. *J. Exp. Bot.* 62 (15), 5509–5519. <https://doi.org/10.1093/jxb/err228>.
- Broadhurst, C.L., Chaney, R.L., 2016. Growth and metal accumulation of an *alysium murale* Ni hyperaccumulator ecotype co-cropped with *alysium montanum* and perennial ryegrass in serpentine soil. *Front. Plant Sci.* 7, 451.
- Butterfield, C.N., Lee, S.W., Tebo, B.M., 2016. The role of bacterial spores in metal cycling and their potential application in metal contaminant bioremediation. *Microbiol. Spectr.* 4 (2), 10–1128.
- Cailliatte, R., Schikora, A., Briat, J.F., Mari, S., Curie, C., 2010. High-affinity manganese uptake by the metal transporter *nram1* is essential for *Arabidopsis* growth in low manganese conditions. *Plant Cell* 22 (3), 904–917. <https://doi.org/10.1105/tpc.109.073023>.
- Chen, L., Long, C., Wang, D., Yang, J., 2020. Phytoremediation of cadmium (Cd) and uranium (U) contaminated soils by *Brassica juncea* L. enhanced with exogenous application of plant growth regulators. *Chemosphere* 242, 125112.
- Campos, N.V., Arcaño-Silva, S., Freitas-Silva, L., de Araújo, T.O., Souza-Fernandes, D.P., Azevedo, A.A., 2018. Arsenic hyperaccumulation in *Pityrogramma calomelanos* L. (Link): adaptive traits to deal with high metalloid concentrations. *Environ. Sci. Pollut. Res.* 25, 10720–10729.
- Clemens, S., Deinlein, U., Ahmadi, H., Höreth, S., Uraguchi, S., 2013. Nicotianamine is a major player in plant Zn homeostasis. *Biometals* 26, 623–632.
- Carrier, P., Barylá, A., Havaux, M., 2003. Cadmium distribution and microlocalization in oilseed rape (*Brassica napus*) after long-term growth on cadmium-contaminated soil. *Planta* 216 (6), 939–950. <https://doi.org/10.1007/s00425-002-0947-6>.
- Cervantes, C., Campos-García, J., Devars, S., Gutiérrez-Corona, F., Loza-Tavera, H., Torres-Guzmán, J.C., Moreno-Sánchez, R., 2001. Interactions of chromium with microorganisms and plants. *FEMS Microbiol. Rev.* 25 (3), 335–347.
- Chai, T., Chen, Q., Zhang, Y., Dong, J., An, C., 2003. Cadmium resistance in transgenic tobacco plants enhanced by expressing bean HM-responsive gene *PvSR2*. *Sci. China Ser. C Life Sci.* 46, 623–630.
- Chellaiyah, E.R., 2018. Cadmium (heavy metals) bioremediation by *Pseudomonas aeruginosa*: a minireview. *Appl. Water Sci.* 8, 154. <https://doi.org/10.1007/s13201-018-0796-5>.
- Cho, U.H., Seo, N.H., 2005. Oxidative stress in *Arabidopsis thaliana* exposed to cadmium is due to hydrogen peroxide accumulation. *Plant Sci.* 168 (1), 113–120. <https://doi.org/10.1016/j.plantsci.2004.07.021>.
- Clemens, S., Palmgren, M.G., Krämer, U., 2002. A long way ahead: understanding and engineering plant metal accumulation. *Trends Plant Sci.* 7 (7), 309–315. [https://doi.org/10.1016/S1360-1385\(02\)02295-1](https://doi.org/10.1016/S1360-1385(02)02295-1).
- Clemens, S., 2006. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie* 88 (11), 1707–1719. <https://doi.org/10.1016/j.biochi.2006.07.003>.
- Clemens, S., Peršoh, D., 2009. Multi-tasking phytochelatin synthases. *Plant Sci.* 177 (4), 266–271. <https://doi.org/10.1016/j.plantsci.2009.06.008>.
- Cobbett, C.S., 2000. Phytochelatin and their roles in heavy metal detoxification. *Plant Physiol.* 123 (3), 825–832.
- Colangelo, E.P., Guerinet, M.L., 2006. Put the metal to the petal: metal uptake and transport throughout plants. *Curr. Opin. Plant Biol.* 9 (3), 322–330. <https://doi.org/10.1016/j.pbi.2006.03.015>.
- Curie, C., Cassin, G., Couch, D., Divol, F., Higuchi, K., Le Jean, M., Misson, J., Schikora, A., Czernic, P., Mari, S., 2009. Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. *Ann. Bot.* 103 (1), 1–11. <https://doi.org/10.1093/aob/mcn207>.
- Dahmani-Muller, H., Van Oort, F., Gélis, B., Balabane, M., 2000. Strategies of HM uptake by three plant species growing near a metal smelter. *Environ. Pollut.* 109 (2), 231–238.
- DalCorso, G., Farinati, S., Maistri, S., Furini, A., 2008. How plants cope with cadmium: staking all on metabolism and gene expression. *J. Integr. Plant Biol.* 50 (10), 1268–1280.
- Das, N., Bhattacharya, S., Maiti, M.K., 2016. Enhanced cadmium accumulation and tolerance in transgenic tobacco overexpressing rice metal tolerance protein gene *OsMTP1* is promising for phytoremediation. *Plant Physiol. Biochem.* 105, 297–309.
- Deepalakshmi, A.P., Ramakrishnaiah, H., Ramachandra, Y.L., Kumar, N.N., 2014. Leaves of higher plants as indicators of HM pollution along the urban roadways. *Int. J. Sci. Technol.* 3 (6), 340–346.
- Deinlein, U., Weber, M., Schmidt, H., Rensch, S., Trampczynska, A., Hansen, T.H., Husted, S., Schjoerring, J.K., Talke, I.N., Krämer, U., Clemens, S., 2012. Elevated nicotianamine levels in *Arabidopsis halleri* roots play a key role in zinc hyperaccumulation. *Plant Cell* 24 (2), 708–723. <https://doi.org/10.1105/tpc.111.095000>.
- Demtröder, L., Narberhaus, F., Masepohl, B., 2019. Coordinated regulation of nitrogen fixation and molybdate transport by *molybdenum*. *Mol. Microbiol.* 111 (1), 17–30.
- Dhanapal, A.R., Thiruvengadam, M., Vairavanathan, J., Venkidasamy, B., Easwaran, M., Ghorbanpour, M., 2024. Nanotechnology approaches for the remediation of agricultural polluted soils. *ACS Omega* 9 (12), 13522–13533. <https://doi.org/10.1021/acsomega.3c09776>.
- Droppa, M., Horváth, G., 1990. The role of copper in photosynthesis. *CRC Crit. Rev. Plant Sci.* 9 (2), 111–123.
- Ducic, T., Polle, A., 2005. Transport and detoxification of manganese and copper in plants. *Braz. J. Plant Physiol.* 17, 103–112.
- El-Seedi, H.R., El-Shabasy, R.M., Khalifa, S.A., Saeed, A., Shah, A., Shah, R., Iftikhar, F.J., Abdel-Daim, M.M., Omri, A., Hajrahand, N.H., Sabir, J.S., 2019. Metal nanoparticles fabricated by green chemistry using natural extracts: biosynthesis, mechanisms, and applications. In: *RSC Adv.*, 9, pp. 24539–24559.
- El-Sappah, A.H., Abbas, M., Rather, S.A., Wani, S.H., Soaud, N., Noor, Z., Qiulan, H., Eldomiati, A.S., Mir, R.R., Li, J., 2023. Genome-wide identification and expression analysis of metal tolerance protein (MTP) gene family in soybean (*Glycine max*) under HM stress. *Mol. Biol. Rep.* 50 (4), 2975–2990.
- El-Mahdy, O.M., Mohamed, H.I., Mogazy, A.M., 2021. Biosorption effect of *Aspergillus niger* and *penicillium chrysosporium* for Cd-and Pb-contaminated soil and their physiological effects on *Vicia faba* L. *Environ. Sci. Pollut. Res.* 28 (47), 67608–67631.
- Fomina, M., Gadd, G.M., 2014. Biosorption: current perspectives on concept, definition and application. *Bioresour. Technol.* 160, 3–14. <https://doi.org/10.1016/j.biortech.2013.12.102>.
- Foyer, C.H., Noctor, G., 2005. Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 17 (7), 1866–1875. <https://doi.org/10.1105/tpc.105.033589>.
- Franchi, E., Rolli, E., Marasco, R., Agazzi, G., Borin, S., Cosmina, P., Pedron, F., Rosellini, I., Barbaferri, M., Petruzzelli, G., 2017. Phytoremediation of a multi contaminated soil: mercury and arsenic phytoextraction assisted by mobilizing agent and plant growth promoting bacteria. *J. Soils Sediments* 17, 1224–1236.
- Freeman, J.L., Persans, M.W., Nieman, K., Albrecht, C., Peer, W., Pickering, I.J., Salt, D. E., 2004. Increased glutathione biosynthesis plays a role in nickel tolerance in *Thlaspi* nickel hyperaccumulators. *Plant Cell* 16 (8), 2176–2191.
- Gao, J., Sun, L., Yang, X., Liu, J.X., 2013. Transcriptomic analysis of cadmium stress response in the HM hyperaccumulator *Sedum alfredii* Hance. *PLoS One* 8 (6), e64643.
- Ge, L.Q., Cang, L., Liu, H., Zhou, D.M., 2016. Effects of warming on uptake and translocation of cadmium (Cd) and copper (Cu) in a contaminated soil-rice system under Free Air Temperature Increase (FATI). *Chemosphere* 155, 1–8.
- Grill, E., Löffler, S., Winnacker, E.L., Zenk, M.H., 1989. Phytochelatin, the heavy-metal-binding peptides of plants, are synthesized from glutathione by a specific γ -glutamylcysteine dipeptidyl transpeptidase (phytochelatin synthase). In: *Proc. Natl. Acad. Sci.*, 86, pp. 6838–6842.
- Ghosh, S., Rusyn, I., Dmytruk, O.V., Dmytruk, K.V., Onyeaka, H., Gryzenhout, M., Gaffoor, Y., 2023. Filamentous fungi for sustainable remediation of pharmaceutical compounds, HM and oil hydrocarbons. *Front. Bioeng. Biotechnol.* 11 h t t p s : / / d o i . o r g / 1 0 . 3389/fbioe.2023.1106973.
- Gisbert, C., Ros, R., De Haro, A., Walker, D.J., Bernal, M.P., Serrano, R., Navarro-Aviñó, J., 2003. A plant genetically modified that accumulates Pb is especially promising for phytoremediation. *Biochem. Biophys. Res. Commun.* 303 (2), 440–445.
- Gorny, J., Billon, G., Lesven, L., Dumoulin, D., Madé, B., Noiriel, C., 2015. Arsenic behavior in river sediments under redox gradient: a review. *Sci. Total Environ.* 505, 423–434. <https://doi.org/10.1016/j.scitotenv.2014.10.011>.
- Grennan, A.K., 2009. Identification of genes involved in metal transport in plants. *Plant Physiol.* 149 (4), 1623–1624. <https://doi.org/10.1104/pp.109.900287>.
- Grennan, A.K., 2011. Metallothioneins, a diverse protein family. *Plant Physiol.* 155 (4), 1750–1751. <https://doi.org/10.1104/pp.111.900407>.
- Grotz, N., Guerinet, M.L., 2006. Molecular aspects of Cu, Fe and Zn homeostasis in plants. *Biochim. Biophys. Acta Mol. Cell Res.* 1763 (7), 595–608. <https://doi.org/10.1016/j.bbamcr.2006.05.014>.
- Guerinet, M.L., 2000. The ZIP family of metal transporters. *Biochim. Biophys. Acta (BBA)-Biomembr.* 1465 (1–2), 190–198.
- Guo, J., Xu, L., Su, Y., Wang, H., Gao, S., Xu, J., Que, Y., 2013. ScMT2-1-3, a metallothionein gene of sugarcane, plays an important role in the regulation of heavy metal tolerance/accumulation. *Biomed. Res. Int.* (1), 904769, 2013.
- Ha, S.B., Smith, A.P., Howden, R., Dietrich, W.M., Buggs, S., O'Connell, M.J., Goldsbrough, P.B., Cobbett, C.S., 1999. Phytochelatin synthase genes from *Arabidopsis* and the yeast *Schizosaccharomyces pombe*. *Plant Cell* 11 (6), 1153–1163.
- Haag-Kerwer, A., Schäfer, H.J., Heiss, S., Walter, C., Rausch, T., 1999. Cadmium exposure in *Brassica juncea* causes a decline in transpiration rate and leaf expansion without effect on photosynthesis. *J. Exp. Bot.* 50 (341), 1827–1835.
- Haider, F.U., Liqun, C., Coulter, J.A., Cheema, S.A., Wu, J., Zhang, R., Wenjun, M., Farooq, M., 2021. Cadmium toxicity in plants: impacts and remediation strategies. *Ecotoxicol. Environ. Saf.* 211, 111887.
- Handique, G.K., Handique, A.K., 2009. Proline accumulation in lemongrass (*Cymbopogon flexuosus* Stapf.) due to HM stress. *J. Environ. Biol.* 30 (2), 299–302.
- Hall, J.L., 2002. Cellular mechanisms for HM detoxification and tolerance. *J. Exp. Bot.* 53 (366), 1–11. <https://doi.org/10.1093/jxb/53.366.1>.
- Haydon, M.J., Cobbett, C.S., 2007. Transporters of ligands for essential metal ions in plants: research review. *New Phytol.* 174 (3), 499–506. <https://doi.org/10.1111/j.1469-8137.2007.02051.x>.
- Hasanuzzaman, M., Nahar, K., Anee, T.I., Fujita, M., 2017. Glutathione in plants: biosynthesis and physiological role in environmental stress tolerance. *Physiol. Mol. Biol. Plants* 23, 249–268.
- He, J., Chen, J.P., 2014. A comprehensive review on biosorption of heavy metals by algal biomass: materials, performances, chemistry, and modeling simulation tools. *Bioresour. Technol.* 160, 67–78.

- Henagamage, A.P., Peries, C.M., Seneviratne, G., 2022. Fungal-bacterial biofilm mediated HM rhizo-remediation. *World J. Microbiol. Biotechnol.* 38 (5), 85.
- Huang, Y., He, G., 2021. Genome-wide identification of MATE gene family in potato (*Solanum tuberosum* L.) and expression analysis in HM stress. *Front. Genet.* 12, 650500.
- Huang, F., Dang, Z., Guo, C.L., Lu, G.N., Gu, R.R., Liu, H.J., Zhang, H., 2013. Biosorption of Cd (II) by live and dead cells of *Bacillus cereus* RC-1 isolated from cadmium-contaminated soil. *Colloids Surf. B Biointerfaces* 107, 11–18.
- Huang, B., Xin, J., Dai, H., Liu, A., Zhou, W., Yi, Y., Liao, K., 2015. Root morphological responses of three hot pepper cultivars to Cd exposure and their correlations with Cd accumulation. *Environ. Sci. Pollut. Res.* 22, 1151–1159.
- Hu, N., Lang, T., Ding, D., Hu, J., Li, C., Zhang, H., Li, G., 2019. Enhancement of repeated applications of chelates on phytoremediation of uranium contaminated soil by *Macleaya cordata*. *J. Environ. Radioact.* 199, 58–65.
- Igiri, B.E., Okoduwa, S.I., Idoko, G.O., Akabuogu, E.P., Adeyi, A.O., Ejiogu, I.K., 2018. Toxicity and bioremediation of heavy metals contaminated ecosystem from tannery wastewater: a review. *J. Toxicol.* 2018 (1), 2568038.
- Jan, S., Parray, J.A., Jan, S. & Parray, J.A. (2016). HM uptake in plants. *Approaches HM Toler. Plants*, 1–18.
- Júnior, C.A.L., de Sousa Barbosa, H., Galazzi, R.M., Koolen, H.H.F., Gozzo, F.C., Arruda, M.A.Z., 2015. Evaluation of proteome alterations induced by cadmium stress in sunflower (*Helianthus annuus* L.) cultures. *Ecotoxicol. Environ. Saf.* 119, 170–177.
- John, R., Ahmad, P., Gadgil, K. & Sharma, S. (2009). HM toxicity: effect on plant growth, biochemical parameters and metal accumulation by *Brassica juncea* L.
- Juwarkar, A.A., Yadav, S.K., 2010. Bioaccumulation and biotransformation of heavy metals. *Bioremediation Technology: Recent Advances*. Springer Netherlands, Dordrecht, pp. 266–284.
- Kang, J., Park, J., Choi, H., Burla, B., Kretschmar, T., Lee, Y., Martinoia, E., 2011. Plant ABC transporters. *Arab. Book/Am. Soc. Plant Biol.* 9, e0153.
- Kumar, S.S., Kadier, A., Malyan, S.K., Ahmad, A., & Bishnoi, N.R. (2017). Phytoremediation and rhizoremediation: uptake, mobilization and sequestration of heavy metals by plants. *Plant-microbe interactions in agro-ecological perspectives: volume 2: microbial interactions and agro-ecological impacts*, 367–394.
- Kalis, E.J., 2006. *Chemical Speciation and Bioavailability of Heavy Metals in Soil and Surface Water*. Wageningen University and Research.
- Kafle, A., Timilsina, A., Gautam, A., Adhikari, K., Bhattacharai, A. & Aryal, N., (2022). Phytoremediation: mechanisms, plant selection and enhancement by natural and synthetic agents. *Environ. Adv.* 8, 100203.
- Khandekar, S., Leisner, S., 2011. Soluble silicon modulates expression of Arabidopsis thaliana genes involved in copper stress. *J. Plant Physiol.* 168 (7), 699–705.
- Khan, A.U., Malik, N., Khan, M., Cho, M.H., Khan, M.M., 2018. Fungi-assisted silver nanoparticle synthesis and their applications. *Bioprocess. Biosyst. Eng.* 41, 1–20. <https://doi.org/10.1007/s00449-017-1846-3>.
- Kim, D.Y., Bovet, L., Kushnir, S., Noh, E.W., Martinoia, E., Lee, Y., 2006. AtATM3 is involved in HM resistance in Arabidopsis. *Plant Physiol.* 140 (3), 922–932.
- Kim, Y.H., Khan, A.L., Kim, D.H., Lee, S.Y., Kim, K.M., Waqas, M., Jung, H.Y., Shin, J.H., Kim, J.G., Lee, J.J., 2014. Silicon mitigates HM stress by regulating P-type HM ATPases, *Oryza sativa* low silicon genes, and endogenous phytohormones. *BMC Plant Biol.* 14, 1–13.
- Kohda, Y.H.T., Qian, Z., Chien, M.F., Miyauchi, K., Endo, G., Suzui, N., Yin, Y.G., Kawachi, N., Ikeda, H., Watabe, H., Kikunaga, H., 2021. New evidence of arsenic translocation and accumulation in *Pteris vittata* from real-time imaging using positron-emitting ⁷⁴As tracer. *Sci. Rep.* 11 (1), 12149.
- Konlechner, C., Türktas, M., Langer, I., Vaculik, M., Wenzel, W.W., Puschenreiter, M., Hauser, M.T., 2013. Expression of zinc and cadmium responsive genes in leaves of willow (*Salix caprea* L.) genotypes with different accumulation characteristics. *Environ. Pollut.* 178, 121–127.
- Krämer, U., Cotter-Howells, J.D., Charnock, J.M., Baker, A.J., Smith, J.A.C., 1996. Free histidine as a metal chelator in plants that accumulate Ni. *Nature* 379 (6566), 635–638.
- Krämer, U., Talke, I.N., Hanikenne, M., 2007. Transition metal transport. *FEBS Lett.* 581 (12), 2263–2272. <https://doi.org/10.1016/j.febslet.2007.04.010>.
- Kroh, G.E., Pilon, M., 2020. Regulation of iron homeostasis and use in chloroplasts. *Int. J. Mol. Sci.* 21 (9), 3395.
- Krzyszowska, M., 2011. The cell wall in plant cell response to trace metals: polysaccharide remodeling and its role in defense strategy. *Acta Physiol. Plant.* 33, 35–51.
- Kukreja, S., Goutam, U., 2012. Phytoremediation: a new hope for the environment. *Front. Recent Dev. Plant Sci.*, pp. 149–171.
- Kumar, A., Dubey, A.K., Kumar, V., Ansari, M.A., Narayan, S., Kumar, S., Pandey, V., Shirke, P.A., Pande, V., Sanyal, I., 2020. Over-expression of chickpea glutaredoxin (CaGrx) provides tolerance to HMs by reducing metal accumulation and improved physiological and antioxidant defence system. *Ecotoxicol. Environ. Saf.* 192, 110252.
- Kumpiene, J., Lagerkvist, A., Maurice, C., 2008. Stabilization of As, Cr, Cu, Pb and Zn in soil using amendments – A review. *Waste Manag.* 28 (1), 215–225.
- Li, J., Wang, Y., Zheng, L., Li, Y., Zhou, X., Li, J., Gu, D., Xu, E., Lu, Y., Chen, X., Zhang, W., 2019. The intracellular transporter AtNRAMP6 is involved in Fe homeostasis in Arabidopsis. *Front. Plant Sci.* 10, 1–11. <https://doi.org/10.3389/fpls.2019.01124>.
- Li, Z., Wang, F., Bai, T., Tao, J., Guo, J., Yang, M., Wang, S., Hu, S., 2016. Lead immobilization by geological fluorapatite and fungus *Aspergillus niger*. *J. Hazard. Mater.* 320, 386–392. <https://doi.org/10.1016/j.jhazmat.2016.08.051>.
- Li, Q., Sun, W., Chen, C., Dong, D., Cao, Y., Dong, Y., Yu, L., Yue, Z., Jin, X., 2022. Overexpression of histone demethylase gene *SlJM524* from tomato confers Cd tolerance by regulating metal transport-related protein genes and flavonoid content in Arabidopsis. *Plant Sci.* 318, 111205.
- Li, H., Jiang, M., Che, L.L., Nie, L., Yang, Z.M., 2012. BjHO-1 is involved in the detoxification of HM in India mustard (*Brassica juncea*). *Biomaterials* 25, 1269–1279.
- Li, S., Wu, X., Xie, J., 2023. Biomineralization technology for solidification/stabilization of heavy metals in ecosystem: status and perspective. *Front. Ecol. Evol.* 11, 1189356.
- Liedtschulte, V., Wächter, A., Zsigang, A., Rausch, T., 2010. Exploiting plants for glutathione (GSH) production: uncoupling GSH synthesis from cellular controls results in unprecedented GSH accumulation. *Plant Biotechnol. J.* 8 (7), 807–820.
- Liu, D., An, Z., Mao, Z., Ma, L., Lu, Z., 2015. Enhanced heavy metal tolerance and accumulation by transgenic sugar beets expressing *Streptococcus thermophilus* StGCS-GS in the presence of Cd, Zn and Cu alone or in combination. *PLoS One* 10 (6), e0128824.
- Liu, H., Zhao, H., Wu, L., Liu, A., Zhao, F.J., Xu, W., 2017. HM ATPase 3 (HMA3) confers cadmium hypertolerance on the cadmium/zinc hyperaccumulator *Sedum plumbizincicola*. *New Phytol.* 215 (2), 687–698.
- Lombi, E., Tearall, K.L., Howarth, J.R., Zhao, F.J., Hawkesford, M.J., McGrath, S.P., 2002. Influence of iron status on cadmium and zinc uptake by different ecotypes of the hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol.* 128, 1359–1367. <https://doi.org/10.1104/pp.010731>.
- Lan, X.Y., Yan, Y.Y., Yang, B., Li, X.Y., Xu, F.L., 2018. Differential expression of proteins in the leaves and roots of cadmium-stressed microsorium pteropus, a novel potential aquatic cadmium hyperaccumulator. *Sci. Total Environ.* 642, 1369–1377.
- Lawal, K.K., Ekeleme, I.K., Onuigbo, C.M., Ikpeazu, V.O., Obiekezie, S.O., 2021. A review on the public health implications of heavy metals. *World J. Adv. Res.* 10 (3), 255–265. <https://doi.org/10.30574/wjarr.2021.10.3.0249>.
- Lavres, J., Castro Franco, G., de Sousa Câmara, G.M., 2016. Soybean seed treatment with nickel improves biological nitrogen fixation and urease activity. *Front. Environ. Sci.* 4, 37.
- Ma, F., Ueno, D., Zhao, F.J., McGrath, S.P., 2005. Subcellular localisation of Cd and Zn in the leaves of a Cd-hyperaccumulating ecotype of *Thlaspi caerulescens*. *Planta* 220 (5), 731–736.
- Maestri, E., Marmioli, M., Visioli, G., Marmioli, N., 2010. Metal tolerance and hyperaccumulation: costs and trade-offs between traits and environment. *Environ. Exp. Bot.* 68 (1), 1–13. <https://doi.org/10.1016/j.envexpbot.2009.10.011>.
- Markkanen, E., van Loon, B., Ferrari, E., Parsons, J.L., Dianov, G.L., Hübscher, U., 2012. Regulation of oxidative DNA damage repair by DNA polymerase λ and MutYH by cross-talk of phosphorylation and ubiquitination. *Proc. Natl. Acad. Sci.* 109 (2), 437–442.
- Martin, S.R., Llugany, M., Barceló, J., Poschenrieder, C., 2012. Cadmium exclusion a key factor in differential Cd-resistance in *Thlaspi arvense* ecotypes. *Biol. Plant.* 56 (4), 729–734.
- Martinoia, E., Meyer, S., De Angeli, A., Nagy, R., 2012. Vacuolar transporters in their physiological context. *Annu. Rev. Plant Biol.* 63 (1), 183–213.
- Mehes-Smith, M., Nkongolo, K., Cholewa, E., 2013. Coping mechanisms of plants to metal contaminated soil. *Environ. Change Sustain.* 54, 53–90.
- Mei, H., Cheng, N.H., Zhao, J., Park, S., Escareno, R.A., Pittman, J.K., Hirschi, K.D., 2009. Root development under metal stress in Arabidopsis thaliana requires the H+ cation antiporter CAX4. *New Phytol.* 183 (1), 95–105.
- Meng, J.G., Zhang, X.D., Tan, S.K., Zhao, K.X., Yang, Z.M., 2017. Genome-wide identification of Cd-responsive NRAMP transporter genes and analyzing expression of NRAMP 1 mediated by miR167 in Brassica napus. *Biomaterials* 30, 917–931. <https://doi.org/10.1007/s10534-017-0057-3>.
- Menguer, P.K., Farthing, E., Peaston, K.A., Ricachenevsky, F.K., Fett, J.P., Williams, L.E., 2013. Functional analysis of the rice vacuolar zinc transporter OsMTP1. *J. Exp. Bot.* 64, 2871–2883. <https://doi.org/10.1093/jxb/ert136>.
- Mengoni, A., Cecchi, L., Gonnelli, C., 2011. Nickel hyperaccumulating plants and allysum bertolonii: model systems for studying biogeochemical interactions in serpentine soils. *Bio-Geo Interactions in Metal-Contaminated Soils*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 279–296.
- Miransari, M., 2011. Hyperaccumulators, arbuscular mycorrhizal fungi and stress of HMs. *Biotechnol. Adv.* 29, 645–653. <https://doi.org/10.1016/j.biotechadv.2011.04.006>.
- Mishra, A., Malik, A., 2013. Recent advances in microbial metal bioaccumulation. *Crit. Rev. Environ. Sci. Technol.* 43 (11), 1162–1222.
- Miwa, K., Fujiwara, T., 2010. Boron transport in plants: co-ordinated regulation of transporters. *Ann. Bot.* 105 (7), 1103–1108. <https://doi.org/10.1093/aob/mcq044>.
- Montanini, B., Blaudez, D., Jeandroz, S., Sanders, D., Chalot, M., et al., 2007. Phylogenetic and functional analysis of the Cation Diffusion Facilitator (CDF) family: improved signature and prediction of substrate specificity. *BMC Genom.* 8. <https://doi.org/10.1186/1471-2164-8-107>.
- Morris, E.C., Griffiths, M., Golebiowska, A., Mairhofer, S., Burr-Hersey, J., Goh, T., Von Wangenheim, D., Atkinson, B., Sturrock, C.J., Lynch, J.P., Vissenberg, K., 2017. Shaping 3D root system architecture. *Curr. Biol.* 27 (17), R919–R930.
- Mousavi, S.R., Galavi, M. & Rezaei, M. (2013). Zinc (Zn) importance for crop production – a review.
- Mukherjee, S., Ghosh, K.K., Chanda, A., 2023. Marine mineral resources. *Environmental Oceanography and Coastal Dynamics: Current Scenario and Future Trends*. Springer, Cham, pp. 269–324. https://doi.org/10.1007/978-3-031-34422-0_14. International Publishing.
- Nahar, N., Rahman, A., Nawani, N.N., Ghosh, S., Mandal, A., 2017. Phytoremediation of arsenic from the contaminated soil using transgenic tobacco plants expressing ACR2 gene of Arabidopsis thaliana. *J. Plant Physiol.* 218, 121–126.
- Naz, T., Khan, M.D., Ahmed, I., Rehman, S.U., Rha, E.S., Malook, I., Jamil, M., 2016. Biosorption of heavy metals by *Pseudomonas* species isolated from sugar industry. *Toxicol. Ind. Health* 32 (9), 1619–1627.

- Nim, Y.S., Wong, K.B., 2019. The maturation pathway of nickel urease. *Inorganics* 7 (7), 85 (Basel).
- Nishida, S., Tsuzuki, C., Kato, A., Aisu, A., Yoshida, J., Mizuno, T., 2011. AtIRT1, the primary iron uptake transporter in the root, mediates excess Ni accumulation in *Arabidopsis thaliana*. *Plant Cell Physiol.* 52 (8), 1433–1442. <https://doi.org/10.1093/pcp/pcr089>.
- Nocito, F.F., Lancilli, C., Crema, B., Fourcroy, P., Davidian, J.C., Sacchi, G.A., 2006. HM stress and sulfate uptake in maize roots. *Plant Physiol.* 141 (3), 1138–1148.
- Nowrouzi, M., Mansouri, B., Nabizadeh, S., Pourkhabbaz, A., 2014. Analysis of heavy metals concentration in water and sediment in the Hara biosphere reserve, southern Iran. *Toxicol. Ind. Health* 30 (1), 64–72.
- Oomen, R.J., Wu, J., Lelièvre, F., Blanchet, S., Richaud, P., Barbier-Brygoo, H., Aarts, M. G., Thomine, S., 2009. Functional characterization of NRAMP3 and NRAMP4 from the metal hyperaccumulator *Thlaspi caerulescens*. *New Phytol.* 181 (3), 37–650. <https://doi.org/10.1111/j.1469-8137.2008.02694.x>.
- Pande, V., Pandey, S.C., Sati, D., Bhatt, P., Samant, M., 2022. Microbial interventions in bioremediation of heavy metal contaminants in agroecosystem. *Front. Microbiol.* 13, 824084. <https://doi.org/10.3389/fmicb.2022.824084>.
- Paulose, B., Chhikara, S., Coomey, J., Jung, H.I., Vatamaniuk, O., Dhankher, O.P., 2013. A γ -glutamyl cyclotransferase protects *Arabidopsis* plants from HM toxicity by recycling glutamate to maintain glutathione homeostasis. *Plant Cell* 25 (11), 4580–4595.
- Papoyan, A., Kochian, L.V., 2004. Identification of *Thlaspi caerulescens* genes that may be involved in HM hyperaccumulation and tolerance. Characterization of a novel HM transporting ATPase. *Plant Physiol.* 136 (3), 3814–3823.
- Patel, A.K., Singhania, R.R., Albarico, F.P.J.B., Pandey, A., Chen, C.W., Dong, C.D., 2022. Organic wastes bioremediation and its changing prospects. *Sci. Total Environ.*, 153889 <https://doi.org/10.1016/j.scitotenv.2022.153889>.
- Peng, K., Luo, C., You, W., Lian, C., Li, X., Shen, Z., 2008. Manganese uptake and interactions with cadmium in the hyperaccumulator—*Phytolacca Americana* L. *J. Hazard. Mater.* 154 (1–3), 674–681.
- Pence, N.S., Larsen, P.B., Ebbs, S.D., Latham, D.L.D., Lasat, M.M., Garvin, D.F., Eide, D., Kochian, L.V., 2000. The molecular physiology of HM transport in the Zn/Cd hyperaccumulator *Thlaspi caerulescens*. *Proc. Natl. Acad. Sci. U. S. A.* 97, 4956–4960. <https://doi.org/10.1073/pnas.97.9.4956>.
- Pete, A.J., Bharti, B., Benton, M.G., 2021. Nano-enhanced bioremediation for oil spills: a review. *ACS ES T Eng.* 1 (6), 928–946. <https://doi.org/10.1021/acesteng.0c00217>.
- Phillips, D.P., Human, L.R.D., Adams, J.B., 2015. Wetland plants as indicators of HM contamination. *Mar. Pollut. Bull.* 92 (1–2), 227–232.
- Pierart, A., Shahid, M., Séjalon-Delmas, N., Dumat, C., 2015. Antimony bioavailability: knowledge and research perspectives for sustainable agricultures. *J. Hazard. Mater.* 289, 219–234. <https://doi.org/10.1016/j.jhazmat.2015.02.011>.
- Pinter, L.F., Salomon, M.V., Berli, F., Bottini, R., Piccoli, P., 2017. Characterization of the As (III) tolerance conferred by plant growth promoting rhizobacteria to *in vitro*-grown grapevine. *Appl. Soil Ecol.* 109, 60–68. <https://doi.org/10.1016/j.apsoil.2016.10.003>.
- Priya, A.K., Gnanasekaran, L., Dutta, K., Rajendran, S., Balakrishnan, D., Soto-Moscoso, M., 2022. Biosorption of heavy metals by microorganisms: evaluation of different underlying mechanisms. *Chemosphere* 307, 135957. <https://doi.org/10.1016/j.chemosphere.2022.135957>.
- Przymusiński, R., Rucińska, R., Gwóźdź, E.A., 2004. Increased accumulation of pathogenesis-related proteins in response of lupine roots to various abiotic stresses. *Environ. Exp. Bot.* 52 (1), 53–61. <https://doi.org/10.1016/j.envexpbot.2004.01.006>.
- Panda, S.K., Baluška, F., Matsumoto, H., 2009. Aluminum stress signaling in plants. *Plant Signal. Behav.* 4 (7), 592–597.
- Parveen, A., Saleem, M.H., Kamran, M., Haider, M.Z., Chen, J.T., Malik, Z., Rana, M.S., Hassan, A., Hur, G., Javed, M.T., Azeem, M., 2020. Effect of citric acid on growth, ecophysiology, chloroplast ultrastructure, and phytoremediation potential of jute (*Corchorus capsularis* L.) seedlings exposed to copper stress. *Biomolecules* 10 (4), 592.
- Poschenrieder, C., Barceló, J., Uversky, V.N. & Kretsinger, E.P.R., (2013). Aluminum in plants. In *Encyclopedia of metalloproteins* (34–38).
- Qiao, K., Gong, L., Tian, Y., Wang, H., Chai, T., 2018. The metal-binding domain of wheat HM ATPase 2 (TaHMA2) is involved in zinc/cadmium tolerance and translocation in *Arabidopsis*. *Plant Cell Rep.* 37, 1343–1352.
- Rajkumar, M., Sandhya, S., Prasad, M.N.V., Freitas, H., 2012. Perspectives of plant-associated microbes in heavy metal phytoremediation. *Biotechnol. Adv.* 30 (6), 1562–1574.
- Rahman, M.A., Hasegawa, H., Kadohashi, K., Maki, T., Ueda, K., 2009. Hydroximinodisuccinic acid (HIDS): a novel biodegradable chelating ligand for the increase of iron bioavailability and arsenic phytoextraction. *Chemosphere* 77 (2), 207–213.
- Raskin, I., Smith, R.D., Salt, D.E., 1997. Phytoremediation of metals: using plants to remove pollutants from the environment. *Curr. Opin. Biotechnol.* 8 (2), 221–226.
- Ravet, K., Pilon, M., 2013. Copper and iron homeostasis in plants: the challenges of oxidative stress. *Antioxid. Redox. Signal.* 19 (9), 919–932.
- Ricachenevsky, F.K., Menguer, P.K., Sperotto, R.A., Williams, L.E., Fett, J.P., 2013. Roles of plant metal tolerance proteins (MTP) in metal storage and potential use in biofortification strategies. *Front. Plant Sci.* 4, 144.
- Romera, E., González, F., Ballester, A., Blázquez, M.L., Muñoz, J.A., 2007. Comparative study of biosorption of heavy metals using different types of algae. *Bioresour. Technol.* 98 (17), 3344–3353.
- Romdhane, L., Panozzo, A., Radhouane, L., Dal Cortivo, C., Barion, G., Vamerli, T., 2021. Root characteristics and metal uptake of maize (*Zea mays* L.) under extreme soil contamination. *Agronomy* 11 (1), 178.
- Roy, S., Labelle, S., Mehta, P., Mihoc, A., Fortin, N., Masson, C., Leblanc, R., Châteauneuf, G., Sura, C., Gallipeau, C., Olsen, C., 2005. Phytoremediation of HM and PAH-contaminated brownfield sites. *Plant Soil* 272 (1/2), 277–290. <https://www.jstor.org/stable/42951704>.
- Ruscitti, M., Arango, M., Beltrano, J., 2017. Improvement of copper stress tolerance in pepper plants (*Capsicum annuum* L.) by inoculation with arbuscular mycorrhizal fungi. *Theor. Exp. Plant Physiol.* 29, 37–49. <https://doi.org/10.1007/s40626-016-0081-7>.
- Rubino, F.M., 2015. Toxicity of glutathione-binding metals: a review of targets and mechanisms. *Toxics* 3 (1), 20–62.
- Rutschow, H.L., Baskin, T.I., Kramer, E.M., 2011. Regulation of solute flux through plasmodesmata in the root meristem. *Plant Physiol.* 155 (4), 1817–1826. <https://doi.org/10.1104/pp.110.168187>.
- Saad, R.B., Hsouna, A.B., Saibi, W., Hamed, K.B., Brini, F., Ghneim-Herrera, T., 2018. A stress-associated protein, LmsAP, from the halophyte *Lobularia maritima* provides tolerance to HMs in tobacco through increased ROS scavenging and metal detoxification processes. *J. Plant Physiol.* 231, 234–243.
- Sable, H., Kumar, V., Singh, V., Rustagi, S., Chahal, S., Chaudhary, V., 2024. Strategically engineering advanced nanomaterials for heavy-metal remediation from wastewater. *Coord. Chem. Rev.* 518, 216079. <https://doi.org/10.1016/j.ccr.2024.216079>.
- Sari, A., Tuzen, M., 2008. Biosorption of cadmium (II) from aqueous solution by red algae (*Ceramium virgatum*): equilibrium, kinetic and thermodynamic studies. *J. Hazard. Mater.* 157 (2–3), 448–454.
- Sasaki, A., Yamaji, N., Ma, J.F., 2014. Overexpression of OsHMA3 enhances Cd tolerance and expression of Zn transporter genes in rice. *J. Exp. Bot.* 65, 6013–6021. <https://doi.org/10.1093/jxb/eru340>.
- Schor-Fumbarov, T., Goldsbrough, P.B., Adam, Z., Tel-Or, E., 2005. Characterization and expression of a metallothionein gene in the aquatic fern *azolla filiculoides* under HM stress. *Planta* 223, 69–76.
- Schat, H., Vooijs, R., 1997. Multiple tolerance and co-tolerance to HMs in *Silene vulgaris*: a co-segregation analysis. *New Phytol.* 136, 489–496.
- Srivastava, N. (2016). Role of phytochelatin in phytoremediation of heavy metals contaminated soils. *Phytoremediation: Management of Environmental Contaminants*, Volume 3, 393–419.
- Sethy, S.K., Ghosh, S., 2013. Effect of HMs on germination of seeds. *J. Nat. Sci. Biol. Med.* 4 (2), 272.
- Spielmann, J., Ahmadi, H., Scheepers, M., Weber, M., Nitsche, S., Carnol, M., Bosman, B., Kroymann, J., Motte, P., Clemens, S., Hanikenne, M., 2020. The two copies of the zinc and cadmium ZIP6 transporter of *Arabidopsis halleri* have distinct effects on cadmium tolerance. *Plant Cell Environ.* 43, 2143–2157. <https://doi.org/10.1111/pce.13806>.
- Shahid, M., Austruy, A., Echevarria, G., Arshad, M., Sanaullah, M., Aslam, M., Nadeem, M., Nasim, W., Dumat, C., 2014. EDTA-enhanced phytoremediation of heavy metals: a review. *Soil Sediment Contam. Int. J.* 23 (4), 389–416.
- Sharma, N., Koul, M., Joshi, N.C., Dufossé, L., Mishra, A., 2024. Fungal-bacterial combinations in plant health under stress: physiological and biochemical characteristics of the filamentous fungus *serendipita indica* and the actinobacterium *zhifengliuella* sp. *ISTPL4* under *in vitro* arsenic stress. *Microorganisms* 12 (2), 405.
- Singh, B.R., Steinnes, E., 2020. Soil and water contamination by HMs. *Soil Processes and Water Quality*. CRC Press, pp. 233–271.
- Sivaraj, R., Salam, H.A., Rajiv, P., Rajendran, V., 2015. *Green Nanotechnology: The Solution to Sustainable Development of Environment*, in: *Environmental Sustainability*. Springer, New Delhi, pp. 311–324.
- Shabani, L., Sabzalain, M.R., Mostafavi pour, S., 2016. Arbuscular mycorrhiza affects nickel translocation and expression of ABC transporter and metallothionein genes in *Festuca arundinacea*. *Mycorrhiza* 26, 67–76.
- Shahid, M., Dumat, C., Khalid, S., Niazi, N.K., Antunes, P.M., 2017. Cadmium bioavailability, uptake, toxicity and detoxification in soil-plant system. *Rev. Environ. Contam. Toxicol.* 241, 73–137. Volume.
- Sheoran, V., Sheoran, A.S., Poonia, P., 2010. Role of hyperaccumulators in phytoextraction of metals from contaminated mining sites: a review. *Crit. Rev. Environ. Sci. Technol.* 41 (2), 168–214. <https://doi.org/10.1080/10643380902718418>.
- Shukla, P.K., Misra, P., Maurice, N. & Ramteke, P.W. (2019). HM toxicity and possible functional aspects of microbial diversity in HM-contaminated sites. *Microbial Genomics in Sustainable Agroecosystems: Volume 2*, 255–317.
- Slatter, K.A. (2013). Ni Accumulation and Tolerance in *Berkheya Codii* and Its Application in Phytoremediation. Master's Thesis, University of Kwazulu, Natal, South Africa.
- Song, W.Y., Martinoia, E., Lee, J., Kim, D., Kim, D.Y., Vogt, E., Shim, D., Choi, K.S., Hwang, I., Lee, Y., 2004. A novel family of cys-rich membrane proteins mediates cadmium resistance in *Arabidopsis*. *Plant Physiol.* 135 (2), 1027–1039.
- Song, W.Y., Yamaki, T., Yamaji, N., Ko, D., Jung, K.H., Fujii-Kashino, M., An, G., Martinoia, E., Lee, Y., Feng, J., 2014. A rice ABC transporter, OsABC1, reduces arsenic accumulation in the grain. *Proc. Natl. Acad. Sci. U. S. A.* 111, 15699–15704. <https://doi.org/10.1073/pnas.1414968111>.
- Soongsombat, P., Kruatrachue, M., Chaiyarat, R., Pokethitayook, P., Ngernsarsarua, C., 2009. Lead tolerance and accumulation inpteris vittataandpityrogramma calomelanos, and their potential for phytoremediation of lead-contaminated soil. *Int. J. Phytoremediation* 11 (4), 396–412. <https://doi.org/10.1080/15226510802565634>.

- Sterckeman, T., Cazes, Y., Gonneau, C., Sirguey, C., 2017. Phenotyping 60 populations of *Noccaea caerulea* provides a broader knowledge of variation in traits of interest for phytoextraction. *Plant Soil* 418, 523–540.
- Sun, Y.B., Zhou, Q.X., Liu, W.T., An, J., Xu, Z.Q., Wang, L., 2009. Joint effects of arsenic and cadmium on plant growth and metal bioaccumulation: a potential Cd-hyperaccumulator and As-excluder *Bidens pilosa* L. *J. Hazard. Mater.* 165 (1–3), 1023–1028.
- Sytar, O., Kumar, A., Latowski, D., Kuczynska, P., Strzalka, K., Prasad, M.N.V., 2013. Heavy metal-induced oxidative damage, defense reactions, and detoxification mechanisms in plants. *Acta Physiol. Plant.* 35, 985–999.
- Szabados, L., Savouré, A., 2010. Proline: a multifunctional amino acid. *Trends Plant Sci.* 15 (2), 89–97. <https://doi.org/10.1016/j.tplants.2009.11.009>.
- Tang, B., Luo, M., Zhang, Y., Guo, H., Li, J., Song, W., Zhang, R., Feng, Z., Kong, M., Li, H., Cao, Z., 2021. Natural variations in the P-type ATPase HM transporter gene *ZmHMA3* control cadmium accumulation in maize grains. *J. Exp. Bot.* 72 (18), 6230–6246.
- Tang, H., Xiang, G., Xiao, W., Yang, Z., Zhao, B., 2024. Microbial mediated remediation of heavy metals toxicity: mechanisms and future prospects. *Front. Plant Sci.* 15, 1420408. <https://doi.org/10.3389/fpls.2024.1420408>.
- Talke, I.N., Hanikenne, M., Krämer, U., 2006. Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator *Arabidopsis halleri*. *Plant Physiol.* 142 (1), 148–167. <https://doi.org/10.1104/pp.105.076232>.
- Takahashi, R., Ishimaru, Y., Senoura, T., Shimo, H., Ishikawa, S., Arai, T., 2011. The OsNRAMP1 iron transporter is involved in Cd accumulation in rice. *J. Exp. Bot.* 62 (14), 4843–4850.
- Takashima, T., Hashimoto, K., Nakamura, R., 2012. Mechanisms of pH-dependent activity for water oxidation to molecular oxygen by MnO₂ electrocatalysts. *J. Am. Chem. Soc.* 134 (3), 1519–1527.
- Teng, Y., Li, Z., Yu, A., Guan, W., Wang, Z., Yu, H., Zou, L., 2022. Phytoremediation of cadmium-contaminated soils by *Solanum nigrum* L. enhanced with biodegradable chelating agents. *Environ. Sci. Pollut. Res.* 29 (37), 56750–56759.
- Thomé, A., Reddy, K.R., Reginatto, C., Cecchin, I., 2015. Review of nanotechnology for soil and groundwater remediation: Brazilian perspectives. *Water Air Soil Pollut.* 226, 1–20. <https://doi.org/10.1007/s11270-014-2243-z>.
- Thomine, S., Lelièvre, F., Debarbieux, E., Schroeder, J.I., Barbier-Brygoo, H., 2003. AtNRAMP3, a multispecific vacuolar metal transporter involved in plant responses to iron deficiency. *Plant J.* 34, 685–695. <https://doi.org/10.1046/j.1365-3113.2003.01760.x>.
- Thomine, S., & Schroeder, J.I. (2004). Plant metal transporters with homology to proteins of the NRAMP family. The NRAMP family. Edited by M. Cellier and P. Gros. Molecular Biology Intelligence Unit, Andes/Kluwer Series, 113–121.
- Tripathi, M., Singh, P., Singh, R., Bala, S., Pathak, N., Singh, S., Chauhan, R.S., Singh, P. K., 2023. Microbial biosorbent for remediation of dyes and heavy metals pollution: a green strategy for sustainable environment. *Front. Microbiol.* 14, 1168954.
- Ueno, D., Milner, M.J., Yamaji, N., Yokosho, K., Koyama, E., Clemencia Zambrano, M., Kaskie, M., Ebbs, S., Kochian, L.V., Ma, J.F., 2011. Elevated expression of *TcHMA3* plays a key role in the extreme Cd tolerance in a Cd-hyperaccumulating ecotype of *Thlaspi caerulescens*. *Plant J.* 66, 852–862. <https://doi.org/10.1111/j.1365-3113.2011.04548.x>.
- Van De Mortel, J.E., Almar Villanueva, L., Schat, H., Kwekkeboom, J., Coughlan, S., Moerland, P.D., Ver Loren van Themaat, E., Koornneef, M., Aarts, M.G., 2006. Large expression differences in genes for iron and zinc homeostasis, stress response, and lignin biosynthesis distinguish roots of *Arabidopsis thaliana* and the related metal hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol.* 142 (3), 1127–1147.
- Verbruggen, N., Hermans, C., Schat, H., 2009. Molecular mechanisms of metal hyperaccumulation in plants. *New Phytol.* 182 (3), 781. <https://doi.org/10.1111/j.1469-8137.2008.02748.x>. *New Phytologist*-781.
- Vaxevedidou, K., Giannikou, S., Pappasiopi, N., 2012. Microbial arsenic reduction in polluted and unpolluted soils from Attica, Greece. *J. Hazard. Mater.* 241–242, 307–315. <https://doi.org/10.1016/j.jhazmat.2012.09.048>.
- Verkleij, J.A.C., Sneller, F.E.C., Schat, H., 2003. Metallothioneins and phytochelatin: ecophysiological aspects. Sulphur in plants. Springer Netherlands, Dordrecht, pp. 163–176.
- Verret, F., Grivot, A., Auroy, P., Leonhardt, N., David, P., Nussaume, L., Vavasseur, A., Richaud, P., 2004. Overexpression of AtHMA4 enhances root-to-shoot translocation of zinc and cadmium and plant metal tolerance. *FEBS Lett.* 576 (3), 306–312.
- Viktorova, J., Jandova, Z., Madlenakova, M., Prouzova, P., Bartunek, V., Vrchoťova, B., Lovacka, P., Musilova, L., Macek, T., 2016. Native phytoremediation potential of *Urtica dioica* for removal of PCBs and heavy metals can be improved by genetic manipulations using constitutive CaMV 35S promoter. *PLoS One* 11 (12), e0167927.
- Wang, X., Winter, D., Ashrafi, G., Schleh, J., Wong, Y.L., Selkoe, D., Rice, S., Steen, J., LaVoie, M.J., Schwarz, T.L., 2011. PINK1 and Parkin target miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell* 147 (4), 893–906. <https://doi.org/10.1016/j.cell.2011.10.018>.
- Wang, Y., Wang, C., Liu, Y., Yu, K., Zhou, Y., 2018a. GmHMA3 sequesters Cd to the root endoplasmic reticulum to limit translocation to the stems in soybean. *Plant Sci.* 270, 23–29. <https://doi.org/10.1016/j.plantsci.2018.02.007>.
- Wang, X., Zhi, J., Liu, X., Zhang, H., Liu, H., Xu, J., 2018b. Transgenic tobacco plants expressing a P1B-ATPase gene from *Populus tomentosa* Carr. (PtoHMA5) demonstrate improved cadmium transport. *Int. J. Biol. Macromol.* 113, 655–661.
- Wang, J.W., Li, Y., Zhang, Y.X., Chai, T.Y., 2013. Molecular cloning and characterization of a Brassica juncea yellow stripe-like gene, BJYSL7, whose overexpression increases HM tolerance of tobacco. *Plant Cell Rep.* 32, 651–662.
- Wang, J., Liang, S., Xiang, W., Dai, H., Duan, Y., Kang, F., Chai, T., 2019. A repeat region from the Brassica juncea HMA4 gene BjHMA4R is specifically involved in Cd 2+ binding in the cytosol under low HM concentrations. *BMC Plant Biol.* 19, 1–16.
- Wang, S., Yao, H., Li, L., Du, H., Guo, P., Wang, D., Rennerberg, H., Ma, M., 2023. Differentially-expressed genes related to glutathione metabolism and HM transport reveals an adaptive, genotype-specific mechanism to Hg2+ exposure in rice (*Oryza sativa* L.). *Environ. Pollut.* 324, 121340.
- Wei, S., Zhou, Q., Wang, X., 2005. Identification of weed plants excluding the uptake of HMs. *Environ. Int.* 31 (6), 829–834.
- Wintz, H., Fox, T., Wu, Y.Y., Feng, V., Chen, W., Chang, H.S., Zhu, T., Vulpe, C., 2003. Expression profiles of *Arabidopsis thaliana* in mineral deficiencies reveal novel transporters involved in metal homeostasis. *J. Biol. Chem.* 278 (48), 47644–47653. <https://doi.org/10.1074/jbc.M309338200>.
- Wróbel, M., Śliwakowski, W., Kowalczyk, P., Kramkowski, K., Dobrzyński, J., 2023. Bioremediation of heavy metals by the genus *Bacillus*. *Int. J. Environ. Res. Public Health* 20 (6), 4964.
- Xia, Y., Liu, J., Wang, Y., Zhang, X., Shen, Z., Hu, Z., 2018. Ectopic expression of *Vicia sativa* caffeoyl-CoA O-methyltransferase (VsCCoAOMT) increases the uptake and tolerance of cadmium in *Arabidopsis*. *Environ. Exp. Bot.* 145, 47–53.
- Xie, Y.F., Zhang, R.X., Qin, L.J., Zhao, D.G., Xia, Z.M., 2022. Genome-wide identification and genetic characterization of the CaMYB family and its response to five types of HM stress in hot pepper (*Capsicum annuum* cv. CM334). *Plant Physiol. Biochem.* 170, 98–109.
- Xu, H., Hao, R.X., Xu, X.Y., Ding, Y., Lu, A.H., Li, Y.H., 2021. Removal of hexavalent chromium by *Aspergillus niger* through reduction and accumulation. *Geomicrobiol. J.* 38 (1), 20–28. <https://doi.org/10.1080/01490451.2020.1807659>.
- Yadav, K.K., Singh, J.K., Gupta, N., Kumar, V.J., 2017. A review of nanobioremediation technologies for environmental cleanup: a novel biological approach. *J. Mater. Environ. Sci.* 8 (2), 740–757.
- Yadav, A., Yadav, K., Ahmad, R., Abd-El salam, K.A., 2023. Emerging frontiers in nanotechnology for precision agriculture: advancements, hurdles and prospects. *Agrochemicals* 2 (2), 220–256. <https://doi.org/10.3390/agrochemicals2020016>.
- Yu, X.Z., Peng, X.Y., Xing, L.Q., 2010. Effect of temperature on phytoextraction of hexavalent and trivalent chromium by hybrid willows. *Ecotoxicology* 19, 61–68.
- Yang, Q., Yang, C., Yu, H., Zhao, Z., Bai, Z., 2021. The addition of degradable chelating agents enhances maize phytoremediation efficiency in Cd-contaminated soils. *Chemosphere* 269, 129373.
- Yamamura, S., Amachi, S., 2014. Microbiology of inorganic arsenic: from metabolism to bioremediation. *J. Biosci. Bioeng.* 118 (1), 1–9. <https://doi.org/10.1016/j.jbiosc.2013.12.011>.
- Yassin, M.A., Elgorban, A.M., El-Samawaty, A.E.R.M., Almunqedhi, B.M., 2021. Biosynthesis of silver nanoparticles using penicillium verrucosum and analysis of their antifungal activity. *Saudi J. Biol.* 28 (4), 2123–2127. <https://doi.org/10.1016/j.sjbs.2021.01.063>.
- Zhou, Y., Xue, M., Yang, Z., Gong, Y., Yuan, J., Zhou, C., Huang, B., 2013. High cadmium pollution risk on vegetable amaranth and a selection for pollution-safe cultivars to lower the risk. *Front. Environ. Sci. Eng.* 7, 219–230.
- Zagorchev, L., Seal, C.E., Kranner, I., Odjakova, M., 2013. A central role for thiols in plant tolerance to abiotic stress. *Int. J. Mol. Sci.* 14 (4), 7405–7432. <https://doi.org/10.3390/ijms14047405>.
- Zhang, M.K., Liu, Z.Y., Wang, H., 2010. Use of single extraction methods to predict bioavailability of HMs in polluted soils to rice. *Commun. Soil Sci. Plant Anal.* 41 (7), 820–831. <https://doi.org/10.1080/00103621003592341>.
- Zhang, Y., Xu, Y.H., Yi, H.Y., Gong, J.M., 2012. Vacuolar membrane transporters OsVIT1 and OsVIT2 modulate iron translocation between flag leaves and seeds in rice. *Plant J.* 72 (3), 400–410. <https://doi.org/10.1111/j.1365-3113.2012.05088.x>.
- Zhang, X., Li, Q., Xu, W., Zhao, H., Guo, F., Wang, P., Wang, Y., Ni, D., Wang, M., Wei, C., 2020. Identification of MTP gene family in tea plant (*Camellia sinensis* L.) and characterization of CsMTP8.2 in manganese toxicity. *Ecotoxicol. Environ. Saf.* 202, 110904. <https://doi.org/10.1016/j.ecoenv.2020.110904>.
- Zhang, M., Liu, B., 2017. Identification of a rice metal tolerance protein OsMTP11 as a manganese transporter. *PLoS One* 12 (4), e0174987.
- Zhao, F.J., Jiang, R.F., Dunham, S.J., McGrath, S.P., 2006. Cadmium uptake, translocation and tolerance in the hyperaccumulator *Arabidopsis halleri*. *New Phytol.* 172 (4), 646–654.
- Zhao, F.J., Lombi, E., Bredeon, T.M.S.P., 2000. Zinc hyperaccumulation and cellular distribution in *Arabidopsis halleri*. *Plant Cell Environ.* 23 (5), 507–514.
- Zhu, B., Gan, C., Gu, L., Du, X., Wang, H., 2021. Identification of NRAMP4 from *Arabis paniculata* enhance cadmium tolerance in transgenic *Arabidopsis*. *J. Genet.* 100 (2), 1–6.
- Liang Zhu, Y., Pilon-Smits, E.A., Jouanin, L., Terry, N., 1999. Overexpression of glutathione synthetase in Indian mustard enhances cadmium accumulation and tolerance. *Plant Physiol.* 119 (1), 73–80.
- Wang, Z., Zhang, Z., Zhang, J., Zhang, Y., Liu, H., Yan, S., 2012. Large-scale utilization of water hyacinth for nutrient removal in Lake Dianchi in China: the effects on the water quality, macrozoobenthos and zooplankton. *Chemosphere* 89 (10), 1255–1261.
- Zulfiqar, U., Farooq, M., Hussain, S., Maqsood, M., Hussain, M., Ishfaq, M., Ahmad, M., Anjum, M.Z., 2019. Lead toxicity in plants: impacts and remediation. *J. Environ. Manage* 250, 109557.
- Zhang, H., Miao, C., Huo, Z., Luo, T., 2022. Effects of zinc oxide nanoparticles transformation in sulfur-containing water on its toxicity to microalgae: physicochemical analysis, photosynthetic efficiency and potential mechanisms. *Water Res.* 223, 119030. <https://doi.org/10.1016/j.watres.2022.119030>.