

Effects of laboratory culture on compatibility between snails and schistosomes

A. Théron, C. Coustau, Anne Rognon, S. Gourbière, M.S. Blouin

► To cite this version:

A. Théron, C. Coustau, Anne Rognon, S. Gourbière, M.S. Blouin. Effects of laboratory culture on compatibility between snails and schistosomes. Parasitology, 2008, 135 (10), pp.1179-1188. 10.1017/S0031182008004745 . halsde-00330477

HAL Id: halsde-00330477 https://hal.science/halsde-00330477

Submitted on 14 Oct 2008 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	PARASITOLOGY 2008, 135(10):1179-1188
2	
3	
4	Effects of laboratory culture on compatibility between snails and schistosomes
5	
6	
7	
8	A. THERON ¹ , C. COUSTAU ² , A. ROGNON ¹ , S. GOURBIÈRE ³ , M. S. BLOUIN ^{1,4*}
9	
10	
11	¹ UMR 5244 CNRS-EPHE-UPVD, Biologie et Ecologie Tropicale et Méditerranéenne,
12	Université Via Domitia. 52 Av. Paul Alduy. 66860 Perpignan Cedex, France.
13	
14	² U547 Inserm, Institut Pasteur de Lille, 1 Rue du Prof. Calmette, BP 245. 59019 Lille Cedex,
15	France.
16	
17	³ EA 3680, Mathématiques Et Physique pour les Systèmes (MEPS), Université de Perpignan
18	Via Domitia, 52 Av. Paul Alduy. 66860 Perpignan Cedex, France.
19	
20	⁴ Dept. Zoology, Oregon State University, Corvallis, OR, 97331 USA
21	
22	
23	*Corresponding author: Tel: 541-737-2362; Fax : 541-737-0501; email:
24	blouinm@science.oregonstate.edu
25 25	
26	Running title: schistosome-snail compatibility
21	

28 SUMMARY

29 The genetic control of compatibility between laboratory strains of schistosomes and their snail 30 hosts has been studied intensively since the 1970s. These studies show (1) a bewildering 31 array of genotype-by-genotype interactions – compatibility between one pair of strains rarely 32 predicts compatibility with other strains, and (2) evidence for a variety of (sometimes 33 conflicting) genetic mechanisms. Why do we observe such variable and conflicting results? 34 One possibility is that it is partly an artifact of the use of laboratory strains that have been in 35 culture for many years and are often inbred. Here we show that results of compatibility trials 36 between snails and schistosomes - all derived from the same natural population – depend very 37 much on whether one uses laboratory-cultured or field-collected individuals. Explanations 38 include environmental effects of the lab on either host or parasite, and genetic changes in 39 either host or parasite during laboratory culture. One intriguing possibility is that genetic 40 bottlenecks during laboratory culture cause the random fixation of alleles at highly 41 polymorphic loci that control the matched/mismatched status of hosts and parasites. We show 42 that a simple model of phenotype matching could produce dose response curves that look very 43 similar to empirical observations. Such a model would explain much of the genotype-by-44 genotype interaction in compatibility observed among strains. 45

46

47 Keywords: Schistosoma mansoni, Biomphalaria glabrata, resistance, susceptibility,

48 compatibility, parasite, host

49 INTRODUCTION

50

51 Compatibility in host-parasite systems can be defined as the state in which a parasite can 52 establish infection and complete its development in a host (Basch, 1975). Thus, compatibility 53 is a joint trait of the parasite and host that probably depends on the genotypes of parasite and 54 host at many different loci. Consequently, there is ample opportunity for complex genotype-55 by-genotype interactions: the phenotype of the host (susceptible/resistant) depends on the genotype of the parasite, and the phenotype of the parasite (infective/noninfective) depends on 56 57 the genotype of the host (Lambrechts et al., 2006). Genotype-by-genotype interaction is 58 particularly apparent in the literature on compatibility between schistosomes and their snail 59 first intermediate hosts (Richards and Shade, 1987; Morand et al., 1996; Webster et al., 2004; 60 Theron and Cousteau, 2005). As an agent of human schistosomiasis, Schistosoma mansoni 61 and its new world snail host, Biomphalaria glabrata, have been the subjects of numerous 62 studies on the genetic basis of variation in compatibility between different laboratory strains 63 of snails and parasite (here we use the term "strain" to refer to a field isolate maintained in the laboratory for at least one generation). One consistent result, in study after study, is that snail 64 65 strains that are naturally or selected to be highly incompatible with one particular schistosome 66 strain, often remain highly compatible with other schistosome strains (Richards and Shade, 67 1987; Webster and Woolhouse, 1998; Webster et al., 2004). Susceptibility to particular 68 schistosome strains can be highly heritable in snails (Richards and Shade, 1987; Richards et 69 al., 1992; Webster et al., 2004). However, attempts to analyze the genetic basis of differences 70 among particular snail strains in their susceptibility to a given schistosome strain have 71 suggested a variety of mechanisms, including single or multi-locus inheritance, and either 72 dominance or recessiveness of the trait (Richards, 1975; Richards et al., 1992; Webster 2001). 73 To date, only one candidate locus has been found to associate with compatibility in either 74 parasite or host (Goodall et al., 2006; Bender et al., 2007).

75

Here we present compatibility studies conducted using lab and field samples of snails and schistosomes, all derived from the same natural population in Guadeloupe, West Indies. We show that results of compatibility trials depend very much on whether one uses laboratorycultured or field-collected individuals. We lay out possible hypotheses to explain some of these results in the hopes of stimulating new research directions.

81

82 MATERIALS AND METHODS

83 Brief overview of experiments

84 We tested snail-schistosome compatibility through four main types of experiments (table 1) :

85 (i) compatibility trails using lab strains of snails and schistosomes ; (ii) compatibility trails

86 using wild snails and wild schistosomes; (iii) compatibility trials involving first wild snails

87 and parasites and later their corresponding snails and schistosomes after each of the first three

88 generations of lab culture; finally (iv) compatibility trials using lab snails challenged by wild

- 89 schistosomes.
- 90

91 Sampling site

92 All host and parasite samples in this study originated from the same transmission site at Dans

93 Fond (DFO; N:16°18.500', W:061°30.720'), located in the marshy forest of Grande Terre

94 Island in Guadeloupe, West Indies (Theron and Pointier, 1995). This site has been the subject

95 of numerous studies on the ecology, dynamics and genetics of larval and adult *S. mansoni*,

96 their intermediate host, *B. glabrata*, and their definitive host, *Rattus rattus* (Sire *et al.*, 1999;

97 2001 *a,b*; Theron *et al.*, 2004; Prugnolle *et al.*, 2005*a,b*; 2006). Transmission is characterized

98 here by very low prevalence in snails (0.6% on average), contrasting with very high infection

rates (94%) and heavy schistosome loads (160 worms per rat on average) within the definitive

- 100 hosts.
- 101

102 Measuring compatibility: snail exposure, infection rates and intensities

103 The level of compatibility for a particular snail-schistosome combination is traditionnally 104 quantified as the proportion of snails infected after individual exposure to a fixed number of 105 miracidia, (usually in the range of 5-20 miracidia). Snail infection rates vary with the parasite 106 dose used (Theron et al., 1997), which makes it difficult to compare the outcomes of different 107 studies. Therefore, we evaluated compatibility between host and parasite populations or 108 strains by challenging individual snails with different numbers of miracidia. Although such 109 dose-response curves are labor-intensive to produce, they are much more informative about 110 the dynamics of compatibility between two strains than are single-dose challenges. All 111 challenge experiments described in this study were conducted by the same person (A. 112 Rognon), using the same protocols. For each experiment, snails (10-12 mm in diameter) were 113 exposed individually to a fixed number of miracidia in approximately 10 ml of water for 8 114 hours. Following exposure to miracidia, snails were replaced in their original containers until 115 their infection status was assessed.

117 The infected or uninfected status of the exposed snails was detected by two different 118 methods: (i) the shedding of cercariae 30 days post-exposure (the length of the prepatent 119 period); (ii) the presence of well developed mother sporocysts (MSp) in the head foot region 120 15 days post exposure. For the detection of mother sporocysts, the snails were fixed 15 days 121 post-exposure following the methods described by Theron and Gerard (1994). In brief, snails 122 were relaxed in pond water containing an excess crystalline menthol for 12 hours. The snail 123 body was removed and fixed in modified Raillet-Henry's solution. The number of MSp 124 present in each snail was determined following exhautive dissection of the head-foot zone. In 125 this technique, MSp's were readily observable as translucent white bodies within an opaque 126 grey tissue background (Fig. 1). For field-collected snails, the technique also allowed us to 127 distinguish *a posteriori* between snails infected during the experiment from those naturally 128 infected but undetectable because they were in the prepatent period during sampling. This 129 distinction is easily made by the presence of young daughter sporocysts in the hepatopancreas 130 of the snails at the time of dissection.

131

132 Compatibility trials using laboratory strains

Laboratory strains of snails and of schistosomes have been isolated from Dans Fond and
established in our laboratory at the University of Perpignan three times: in 1992, 2000 and
2005 (Table 1).

136

The snail strains were founded each time using 100-150 uninfected founders from the field. The snail populations were allowed to expand quickly and were then maintained at a census size in the hundreds of individuals. Snails in the lab are allowed to breed freely, and the lab populations are not exposed to any deliberate selection. Note that although *B. glabrata* are hermaphroditic, they preferentially outcross and show Hardy-Weinberg equilibrium at molecular markers in both laboratory and field populations (Prugnolle *et al.*, 2005*b*; unpub. data).

144

The schistosomes strains were founded each time using cercariae from 43-47 infected snails previously exposed to 20 miracidia hatched from eggs collected from the livers of six naturally-infected *R. Rattus*. In the lab, schistosome strains are passaged each generation through five mice and approximately 24 infected snails (8-10 miracidia at exposure).

149

- 150 Compatibility trails were performed for snail and parasite strains established in 1992 and
- 151 2005 and tested after 5 years (Theron *et al.*, 1997) and 2 years maintenance in the lab,
- respectively (labelled as "LAB 1997" and "LAB 2007" in Fig. 2A). Dose-response curves
- 153 were obtained by challenging individual snails (45-50 snails per treatment) with doses of 1, 2,
- 154 5, 10, 20, 30 and 50 miracidia. The doses of 20 and 5 miracidia were omitted for the LAB
- 155 1997 and LAB 2007 experiments, respectively. For both experiments snail infections rates
- and parasite intensities were evaluated by mother sporocyst count.
- 157
- 158 Compatibility trials using wild schistosomes and wild snails
- 159 In 2000 and 2005 we generated dose-response curves using schistosomes and snails collected
- 160 directly from the transmission site (i.e. never passaged in the lab). We refer to these
- 161 experiments as "DFO 2000" and "DFO 2005". All challenges were conducted at the INRA-
- 162 Duclos laboratory in Guadeloupe. The livers of heavily infected rats were crushed in a saline
- 163 solution, and the homogenates were passed through different filters to finally retrieve only
- 164 schistosome eggs. For each experiment, schistosome eggs were collected from six heavily
- 165 infected rats (> 100 worms/host) and placed in fresh water to hatch into miracidia.
- 166
- 167 Dose-response curves were obtained by challenging individual snails (50 snails per treatment)
- 168 with doses of 1, 2, 10, 20, 30 and 50 miracidia (Fig. 2B). For the DFO 2000 experiment, all
- 169 the snails were fixed 15 days post exposure and dissected to count numbers of mother
- 170 sporocysts per snail. Dissections showed no evidence of pre-patent infections in any of our
- 171 field-collected snails. Also, the prevalence of infection in the field averages only 0.6%.
- 172 Therefore we used the less-laborious cercarial shedding test for the DFO 2005 experiment on
- 173 the assumption that pre-patent infections would be so rare as to not appreciably influence our 174 results.
- 175

176 Compatibility during the first three generations of laboratory passage

- In this experiment we followed the change in compatibility between snails and parasites in going from the field through the first three generations of passage in the laboratory. Here we used a single challenge dose (20 miracidia), rather than generating a dose-response curve each time. In 2000, one hundred field-collected snails were individually exposed to 20 wild miracidia. Fifty of these snails were transported back to our laboratory in France, and cercariae shed by 43 of the 46 survivors were used to infect mice and establish a schistosome
- 183 strain. The remaining 50 wild exposed snails were fixed and dissected to measure infection

184 rates and numbers of mother sporocysts per snail. The lab strain of snails was established 185 from 100 uninfected, field-collected founders. After each of the first three generations of 186 culture in France, 50 lab snails were each challenged with 20 lab miracidia. These were also 187 fixed and dissected to measure infection rates and numbers of mother sporocysts per snail. 188 189 The lab strains isolated in 2005 were tested in the field, and then after each generation of 190 parasite passage, using a dose of 20 miracidia. We have continued testing them each 191 generation, using the same dose. 192 193 Compatibility trials using wild versus laboratory snails or parasites 194 It would be ideal to also have replicated trials in which lab parasites were used to challenge 195 196 only combination we were able to attempt was a single trial in which lab snails were 197 challenged with field parasites. After the laboratory strain of snail established in 2000 had

194 It would be ideal to also have replicated trials in which lab parasites were used to challenge 195 field snails, and field parasites used to challenge lab snails. Owing to logistical constraints the 196 only combination we were able to attempt was a single trial in which lab snails were 197 challenged with field parasites. After the laboratory strain of snail established in 2000 had 198 been in captivity for two years, 300 of these lab snails were transported to Guadeloupe to be 199 challenged with wild miracidia. We generated a dose-response curve using these snails and 198 the same protocols as for the wild-by-wild combinations. Infection rates and intensities 201 (number of mother sporocysts per snail) were measured by dissecting each snail. Because we 202 have not been able to replicate this experiment, we consider the results of this trial to be 203 preliminary. We present them here simply for the sake of completeness.

204

205 RESULTS

206

207 Laboratory schistosomes versus laboratory snails

The dose-response curve generated using the 1992 laboratory strains shows a rapid increase in infection rate up to a dose of 10-20 miracidia, but then levels off at about 65% infection (LAB

210 1997 in Fig. 2A). This result suggests that about 35% of the snails in the laboratory

211 population were completely resistant to infection, regardless of the number of miracidia used

to challenge. A very similar result was obtained with the 2005 lab isolates, except the curve

213 leveled off at about 45% infection (LAB 2007 in Fig. 2A).

214

215 Wild schistosomes versus wild snails

- 216 Results obtained for the two wild-by-wild dose-response curves (experiments DFO 2000 and
- 217 DFO 2005) are shown in Fig. 2B. Infection rates increased rapidly with increasing dose,
- reaching 90-100% at 20-30 miracidia, and 100% at the highest dose.
- 219

220 Change in compatibility in going from field isolates to laboratory strains

221 Among the 50 snails that were each exposed to 20 miracidia in Guadeloupe in 2000, 46 222 survived and 43 of those became infected (93.5%), shed cercariae, and were used to found the 223 schistosome colony in our lab in France. Among the 50 other snails exposed in the same 224 conditions and fixed, 48 harbored developed MSp's (96%). The compatibility of this pair of 225 strains using a dose of 20 miracidia dropped to 51.0%, 59.8% and 54.1% after 1, 2, and 3 226 passages through laboratory mice, respectively (Fig. 3). A similar pattern occurred with the 227 strains isolated in 2005. Again, infection rates dropped from near 100% when wild snails 228 were challenged with 20 wild miracidia apiece, to around 40% after a single generation of 229 passage in the lab. We continue to check the infection rate after each passage, and it remains 230 at about 40% to this day (mean ± 1 s.e. = 41.2 ± 2.8) – right in line with the results of the 231 dose-response curve (Fig. 2A).

232

233 Wild schistosomes versus laboratory snails

Infection rates of laboratory bred snails exposed to wild miracidia increased gradually to 95% at the highest dose (Fig. 2C). Interestingly, the shape of this curve differs from those of the wild-by-wild or lab-by-lab curves. Rather than quickly reaching a plateau, the infection rate increases almost linearly. So this result seems intermediate between those of the lab-by-lab and field-by-field trials. In this case lab snails appeared to be much more susceptible to groups of wild miracidia than to the same number of lab miracidia, yet not as susceptible as wild snails.

241

242 Infection intensities (establishment of mother sporocysts)

243 In the lab-by-lab trials, infection intensities rose gradually and then leveled off at

approximately 2-3 /snail after challenge by 10 or more miracidia (Fig. 2A). In contrast, in the

- 245 wild-by-wild challenge the mean number of mother sporocysts (MSp) that developed within
- infected snails increased steadily to an average of 6.6 ± 1.3 MSp/snail at the 50 miracidial
- 247 dose (Fig. 2B), with a maximum of 16 MSp found in one snail. Infection intensities in the
- 248 wild-schistosome by lab-snail trials (Fig. 2C) were lower than in the field-by-field trials,
- reaching a maximum of 3.4 MSp/snail at a miracidial dose of 50.

- 250 Figure 3 shows the change in number of MSp's per infected snail during the first three
- 251 generations of passage of the 2000 lab strains. Although a dose of 20 wild miracidia
- 252 produced an average of 5.37 MSp per infected snail in Guadeloupe, that number dropped to
- 253 2.04, 1.92 and 1.63 MSp/snail over three generations of passage in the lab. Thus, as with the
- 254 percentage of snails infected, the number of sporocysts per infected snail dropped
- 255 precipitously after the first generation of passage, and then remained relatively constant
- thereafter.
- 257

258 DISCUSSION

259 Loss of compatibility in going from field to lab

In the year-2000 field trials a challenge with 20 wild miracidia produced 95% infection. Yet after a single generation of passage in the lab (of schistosome through the new lab snails) the percent infection dropped to 51% and remained between 54% and 60% for the next two generations. The same result was obtained when we established new strains of parasite and snail in 2005 – almost 100% infection in the field dropped to about 40% after a single passage in lab, and stayed the same thereafter. The number of MSp established per infected snail also drops after laboratory passage (Fig. 3).

267

Thus it appears that in three independent isolations of schistosomes and snails from the same site, there was a massive drop in compatibility in going from field to lab. Even more intriguing is the observation that in two independent isolations of snails and parasites from the field, it appears that a large fraction of the lab snails became completely resistant to the lab strain of schistosomes (Fig. 2A). In stark contrast, we see 100% compatibility when wild snails are challenged with enough wild miracidia (above 20- 30 miracidia), and much higher rates and intensities of infection at the lower doses (Fig. 2B).

275

There are three main hypotheses to explain these results. (1) <u>Experimental artifact:</u> the lab-vslab trials were conducted in France, and the field-by-field trials were conducted in

278 Guadeloupe. Thus some experimental factor might have been consistently different between

the two settings. (2) <u>Environmental effects:</u> something about the laboratory environment

280 makes the parasite less infective and/or the snails more resistant. (2) <u>Genetic effects:</u> either

- the snail population, the parasite population, or both underwent genetic changes during
- 282 laboratory culture.
- 283

284 Experimental artifact

285 We took great pains to insure that the trials conducted in the lab in Guadeloupe were as 286 similar as possible to the trials conducted in the lab in France (same person did all the trials, 287 same equipment, and so on). Regardless, it is difficult to imagine how experimental artifact 288 could produce the plateaus seen in the lab-by-lab curves, which suggest the existence of a 289 completely resistant subset of the lab snail population. For example, if some factor (say the 290 water) reduced the average infectivity of miracidia in the lab in France (or resistance of the 291 snails), then the lab-by-lab curves would have shapes similar to those of the field-by-field 292 curves, but simply with a lower inflection point. Thus, we think that the fundamentally 293 different shapes of the two sets of curves really does reveal an interesting biological 294 phenomenon that deserves further study.

295

296 Environmental effects

297 It is well known that the susceptibility of snails can depend on, for example, their size, age 298 and physiological status (Anderson et al., 1982; Theron et al., 1998; Krist et al., 2004). So it 299 is quite plausible that something about being raised in the lab environment makes snails more 300 resistant. For example, perhaps nutritional status differs or some microbe or other stressor in 301 the lab tank keeps the immune system of lab snails ramped up (Hertel et al. 2002). But again, 302 it is difficult to imagine a mechanism that would generate a completely resistant subset of the 303 population, rather than just a higher average resistance per snail. An environmental effect on 304 the parasite is also possible, although again we do not have any candidate mechanism that 305 would render that strain of parasites completely unable to infect just a subset of the snail 306 population.

307

308 *Genetic effects*

309 (1) <u>Genetic change in the snails:</u> The lab breeding colony of snails is never challenged, so any
310 selected change must have been a correlated response to selection on some other trait that is
311 inadvertently under selection in the lab. Genetic drift remains a possibility, but the lab colony
312 was founded each time using more than 100 individuals, and remained large thereafter. Also,
313 lab strains of *B. glabrata* tend to retain substantial molecular genetic diversity (e.g. Mulvey
314 and Vrijenhoek, 1981; Campos *et al.*, 2002). Thus, it is hard to believe that just one or two
315 generations of drift in the snails caused the changes we observed (e.g. Fig. 3).

317 (2) <u>Genetic change in the schistosomes:</u> If selection is involved, one possible source is the
318 definitive host. Rats are the definitive host in nature, but the parasites are passed through
319 mice in the lab. It would be quite interesting if selection for performance in a novel definitive
320 host resulted in a correlated response in infectivity to snails. Such a result would suggest that
321 the parasite uses some common mechanism to infect both the intermediate and definitive
322 hosts.

323

324 An even more plausible genetic explanation is rapid genetic drift in the lab schistosome 325 isolate. The miracidium that infects a snail becomes a sporocyst that then undergoes a round 326 of asexual reproduction that culminates in the release of thousands of genetically identical 327 cercariae. In most labs, rodents are infected by being placed in water that contains cercariae 328 shed by several infected snails. Thus, there is a huge potential for non-random variance 329 among clones in transmission to the next generation if care is not taken to equalize the 330 cercarial contribution from each snail. Strong selection among clones in going from the 331 natural definitive host (rats or humans) to mice or hamsters in the lab would add an 332 additional, non-random component to the among-clone variance in transmission success (e.g. 333 we see high variance among clones in their inherent infectivity to mice; J. Boissier, pers. 334 comm.). Rodents are expensive to maintain, so most lab strains of S. mansoni are passed 335 through only a handful of rodents each generation. A mouse can support 50-100 adult 336 schistosomes at most. We have little idea what is the variance in family size (egg production) 337 among adult worms, and thus what is the effective number of breeders per host that 338 contributes to the next generation (Criscione and Blouin, 2005). Thus, most lab strains of 339 schistosomes probably undergo severe genetic bottlenecks beginning with the first generation 340 of lab passage. Consistent with this prediction, molecular genetic studies show that lab strains 341 of schistosomes have only a fraction of the allelic variation found in field samples, and they 342 also show the distorted allele frequency distributions typical of bottlenecked populations 343 (Cornuet and Luikart, 1997; Stohler et al., 2004).

344

345 A possible mechanism: matching alleles and bottlenecks

346 One hypothesis that would explain our results is that the success or failure of a challenge by

347 one parasite depends on the matched/mismatched status of the host and parasite genotypes, as

- in matching-alleles models (Agrawal and Lively, 2002; Basch, 1975; Theron and Coustau,
- 349 2005). Any one snail and miracidium may have a modest probability of matching. However,
- as a snail is challenged with increasing numbers of miracidia, the probability of at least one

351 match increases rapidly. Increasing the miracidial dose simply involves sampling a larger 352 fraction of the phenotypic diversity in the parasite population. Under this model all snails 353 eventually succumb when challenged by enough wild miracidia because the probability of at 354 least one match approached 1.0. Such a system of phenotype matching could then explain the 355 drop in compatibility after laboratory passage if the phenotypic variation is genetically based. 356 If laboratory passage involves a more severe genetic bottleneck and loss of alleles in the 357 parasite population than in the snail population, then a large fraction of the snails could 358 quickly become "resistant", owing simply to loss of compatible alleles in the schistosome 359 strain. Under this scenario those snails are not "resistant" to schistosomes per se. They are 360 simply not matched by any individuals in the now genetically-depauperate lab strain of 361 parasites.

362

Such a genetically-based system of phenotype matching could generate curves that look very much like those in figure 2A and 2B. If each miracidium has a low, but constant probability of matching a target snail, and if miracidia infect independently of each other, then the probability that a snail becomes infected when challenged by *x* miracidia can be written

368 (1) P(infection) =
$$(1 - (1 - p)^x)$$
,

369

370 where p = the probability of infection per miracidium. For example, in figure 4A we plotted 371 equation 1 with a value of p = 0.11 in the same panel as the wild-vs.-wild data from 2002. 372 Curves like those in Fig. 2B, in which there appears to be a completely resistant subset of 373 snails, can be generated simply by multiplying equation 1 by a constant, *f*, which is the 374 fraction of snails that can be infected. For example, in figure 4B we plotted the 1997 lab-vs.-375 lab data and equation (1) with f = 0.67 and p = 0.2.

376

One can also generate similar-looking curves via a matching genotypes model. Here the snail
becomes infected if at least one miracidium in the pool of challengers carries the same
(matching) genotype as the snail. The probability that a snail becomes infected when
challenged by *x* miracidia equals

382 (2) P(infection) =
$$f \sum_{i=1}^{N_a} (h_i)(1 - (1 - p_i)^x)$$
,

- where N_a = the number of genotypes present in the parasite population, h_i = the frequency of genotype *i* in the host population, p_i = the frequency of genotype *i* in the parasite population. If all snail alleles have a matching allele in the parasite population, then these curves will asymptote to 100% infection as *x*, the number of challenging miracidia, increases. If some snail genotypes are not matched by a genotype in the parasite population, then the curve will asymptote at *f*, the fraction of snails that can be matched.
- 390

The actual shapes of curves generated by equation 2 will obviously depend on the particular allele frequency distributions in parasite and host populations. We can, *a posteriori*, choose parameter values that make the curves from equation 3 fit some of our observed data quite well (Fig. 4). This does not mean we have proof for a matching genotypes model, but it does show that such a model remains a plausible explanation for our results.

396

397 Some possible consequences of using bottlenecked lab strains

398 If the bottleneck hypothesis is correct, then this could explain why studies to date show such a 399 variety of genetic mechanisms and of strain-by-strain interactions. The chance loss or fixation 400 of alleles at compatibility loci may cause different loci to appear overly important in 401 controlling compatibility in different pair-wise combinations. In other words, in different 402 studies a different set of loci may, by chance, explain a disproportionate amount of the 403 variance in compatibility between particular lab strains. Loci that might be particularly 404 misleading in that sense include loci encoding highly polymorphic systems of matching 405 alleles that hosts use to recognize invaders, and/or invaders use to mimic hosts (Zhang et al., 406 2004; Theron and Coustau 2005). Here the compatibility of any pair-wise combination could 407 depend largely on which matching alleles were lost by chance during the domestication of 408 each partner, and so would not predict compatibility with other strains, or with the original 409 field populations. For example, this hypothesis predicts that if we had selected the apparently 410 resistant fraction of snails revealed in Figure 2A, we would have rapidly obtained a snail 411 strain that is highly "resistant" to that particular schistosome strain (we are currently 412 conducting that experiment using the 2005 strains). However, in this case the proportion of 413 snails that did not acquire infection would not be sensus stricto resistant to S. mansoni. They 414 would simply be not matched by the reduced number of parasite genotypes contained within 415 that particular bottlenecked laboratory strain of schistosome. Those snails might remain

416 highly compatible with other schistosome strains that retain different alleles (cf Theron &

417 Coustau, 2005).

418

419 We are not arguing that all variation in compatibility results from polymorphic systems of 420 matching alleles. Compatibility is a complex process that involves many steps from 421 recognizing the invader to preventing its successful establishment (Loker et al., 2004). 422 Indeed, the only resistance locus identified to date (Goodall et al., 2006; Bender et al., 2007) 423 is clearly part of an effector mechanism – killing the parasite once it has been successfully 424 recognized by the host. But the idea that using inbred strains causes different loci to be 425 important in each strain-by-strain comparison would hold for any polymorphic, genetically-426 based mechanism of resistance.

427

428 Some caveats

429 We recognize that replicated trials of each reciprocal combination of field-by-lab would have 430 gone a long way towards identifying which species is responsible for the difference between 431 lab and field trials. The one combination we were able to attempt (lab-snails versus field-432 schistosomes) produced intriguing results in that infection rates were, overall, lower than in 433 the wild-by-wild combination, but there was no plateau. The higher infection rates are 434 consistent with the bottleneck hypothesis. But if parasite diversity was all that mattered, the 435 curve should have looked just like the field-by-field curves, not the more gradual increase 436 with dose that we observed. Taken at face value, this result suggests that something about the 437 snails also changed in the lab. But until this result can be replicated, we hesitate to make too 438 much of it.

439

440 Summary

441 To our knowledge, this is the first study to compare the results of compatibility trials using 442 lab-reared and field-collected individuals from the same source populations. We showed that 443 even a single generation of laboratory passage had a large effect on the results of 444 compatibility trials. Barring experimental artifact, possible explanations include 445 environmental effects on either host or parasite, and genetic change in either host or parasite. 446 We suggest one possible mechanism that could produce curves very much like those 447 observed. The bottleneck hypothesis could explain much of the highly variable strain-by-448 strain compatibilities that are so apparent in the literature on schistosomes and their snail 449 hosts.

451 ACKNOWLEDGEMENTS

- 452 We thank Hervé Mauleon (INRA, Domaine Duclos) for logistical support in Guadeloupe, and
- 453 Jean-Pierre Pointier for help with field samples. This work received financial support from
- 454 the CNRS (EDD), and the Ministère de l'Ecologie et du Développment Durable (Ecosystèms
- 455 Tropicaux n^o 752600). Thanks to C. Bayne, J. Tennessen, C. Adema and M. Steinauer for
- 456 comments on earlier drafts. M. Blouin's sabbatical at CNRS-UPVD was supported by the
- 457 University of Perpignan and Oregon State University.

- 458 REFERENCES
- 459
- 460 Agrawal, A. F. and Lively, C. M. (2003). Modelling infection as a two-step process
- 461 combining gene-for-gene and matching-allele genetics. *Proceedings. Biological sciences /*
- 462 *The Royal Society* **270**, 323-334.
- 463 Anderson, R. M., Mercer, J. G., Wilson, R. A. and Carter, N. P. (1982). Transmission of
- 464 Schistosoma mansoni from man to snail: experimental studies of miracidial survival and
- infectivity in relation to larval age, water temperature, host size and host age. *Parasitology* **85**
- 466 (**Pt 2**), 339-360.
- 467 **Basch, P. F.** (1975). An interpretation of snail-trematode infection rates: specificitity based
- 468 on concordance of compatible phenotypes. *Inernationalt Journal for Parasitology* **5**, 449-452.
- 469 Bayne, C. J., Hahn, U. K. and Bender, R. C. (2001). Mechanisms of molluscan host
- 470 resistance and of parasite strategies for survival. *Parasitology* **123 Suppl,** S159-167.
- 471 Campos, Y. R., Carvalho, O. S., Goveia, C. O. and Romanha, A. J. (2002). Genetic
- 472 variability of the main intermediate host of the Schistosoma mansoni in Brazil, Biomphalaria
- 473 glabrata (Gastropoda: Planorbidae) assessed by SSR-PCR. Acta Tropica 83, 19-27.
- 474 Cornuet, J. M. and Luikart, G. (1996). Description and power analysis of two tests for
- 475 detecting recent population bottlenecks from allele frequency data. *Genetics* **144**, 2001-2014.
- 476 Criscione, C. D. and Blouin, M. S. (2005). Effective sizes of macroparasite populations: a
- 477 conceptual model. *Trends in Parasitol* **21**, 212-217.
- 478 Goodall, C. P., Bender, R. C., Brooks, J. K. and Bayne, C. J. (2006). Biomphalaria
- 479 glabrata cytosolic copper/zinc superoxide dismutase (SOD1) gene: association of SOD1
- 480 alleles with resistance/susceptibility to Schistosoma mansoni. Molecular and Biochemical
- 481 *Parasitology* **147,** 207-210.
- 482 Hertel, L. A., Bayne, C. J. and Loker, E. S. (2002). The symbiont *Capsaspora owczarzaki*,
- 483 nov. gen. nov. sp., isolated from three strains of the pulmonate snail *Biomphalaria glabrata* is
- related to members of the Mesomycetozoea. *International Journal for Parasitology* **32**, 1183-
- 485 1191.
- 486 Krist, A. C., Jokela, J., Wiehn, J. and Lively, C. M. (2004). Effects of host condition on
- 487 susceptibility to infection, parasite developmental rate, and parasite transmission in a snail-
- 488 trematode interaction. *Journal of Evolutionary Biology* **17**, 33-40.
- 489 Lambrechts, L., Fellous, S. and Koella, J. C. (2006). Coevolutionary interactions between
- 490 host and parasite genotypes. *Trends in Parasitol* **22**, 12-16.

- 491 Loker, E. S., Adema, C. M., Zhang, S. M. and Kepler, T. B. (2004). Invertebrate immune
- 492 systems--not homogeneous, not simple, not well understood. *Immunological Reviews* 198, 10493 24.
- 494 Morand, S., Manning, S. D. and Woolhouse, M. E. (1996). Parasite-host coevolution and
- 495 geographic patterns of parasite infectivity and host susceptibility. *Proceedings. Biological*
- 496 sciences / The Royal Society 263, 119-128.
- 497 Mulvey, M. and Vrijenhoek, R. C. (1981). Genetic variation among laboratory strains of the
- 498 planorbid snail *Biomphalaria glabrata*. *Biochemical Genetics* **19**, 1169-1182.
- 499 Prugnolle, F., De Meeus, T., Pointier, J. P., Durand, P., Rognon, A. and Theron, A.
- 500 (2006). Geographical variations in infectivity and susceptibility in the host-parasite system
- 501 Schistosoma mansoni/Biomphalaria glabrata: no evidence for local adaptation. Parasitology
- 502 **133**, 313-319.
- 503 Prugnolle, F., Liu, H., De Meeus, T. and Balloux, F. (2005a). Population genetics of
- 504 complex life-cycle parasites: an illustration with trematodes. *International Journal For*
- 505 *Parasitology* **35,** 255-263.
- 506 Prugnolle, F., Theron, A., Pointier, J. P., Jabbour-Zahab, R., Jarne, P., Durand, P. and
- 507 **De Meeus, T.** (2005b). Dispersal in a parasitic worm and its two hosts: consequence for local 508 adaptation. *Evolution Int J Org Evolution* **59**, 296-303.
- 509 Richards, C. S. (1975). Genetic factors in susceptibility of *Biomphalaria glabrata* for
- 510 different strains of Schistosoma mansoni. *Parasitology* **70**, 231-241.
- 511 Richards, C. S., Knight, M. and Lewis, F. A. (1992). Genetics of Biomphalaria glabrata
- and its effect on the outcome of *Schistosoma mansoni* infection. *Parasitology Today* 8, 171-
- 513 174.
- 514 Richards, C. S. and Shade, P. C. (1987). The genetic variation of compatibility in
- 515 Biomphalaria glabrata and Schistosoma mansoni. Journal of Parasitol 73, 1146-1151.
- 516 Sire, C., Durand, P., Pointier, J. P. and Theron, A. (1999). Genetic diversity and
- 517 recruitment pattern of Schistosoma mansoni in a Biomphalaria glabrata snail population: a
- 518 field study using random-amplified polymorphic DNA markers. Journal of Parasitol 85, 436-
- 519 441.
- 520 Sire, C., Durand, P., Pointier, J. P. and Theron, A. (2001a). Genetic diversity of
- 521 Schistosoma mansoni within and among individual hosts (Rattus rattus): infrapopulation
- 522 differentiation at microspatial scale. *International Journal for Parasitology* **31**, 1609-1616.

- 523 Sire, C., Langand, J., Barral, V. and Theron, A. (2001b). Parasite (Schistosoma mansoni)
- and host (*Biomphalaria glabrata*) genetic diversity: population structure in a fragmented
 landscape. *Parasitology* 122, 545-554.
- 526 Stohler, R. A., Curtis, J. and Minchella, D. J. (2004). A comparison of microsatellite
- 527 polymorphism and heterozygosity among field and laboratory populations of *Schistosoma*
- 528 mansoni. International Journal for Parasitology **34**, 595-601.
- 529 Theron, A. and Coustau, C. (2005). Are Biomphalaria snails resistant to Schistosoma
- 530 mansoni? Journal of Helminthology **79**, 187-191.
- 531 Theron, A. and Gerard, C. (1994). Development of accessory sexual organs n Biomphalaria
- 532 glabrata as related to infection timing by Schistosoma mansoni: Consequences on the energy
- 533 utilisation patterns by the parasite. *The Journal of Molluscan Studies* **60**, 78-85.
- 534 Theron, A., Pages, J. R. and Rognon, A. (1997). Schistosoma mansoni: distribution patterns
- 535 of miracidia among *Biomphalaria glabrata* snail as related to host susceptibility and
- 536 sporocyst regulatory processes. *Experimental Parasitology* **85**, 1-9.
- 537 Theron, A. and Pointier, J. P. (1995). Ecology, dynamics, genetics and divergence of
- 538 trematode populations in heterogeneous environments: the model of *Schistosoma mansoni* in
- the insular focus of Guadeloupe. *Research Reviews in Parasitology* **55**, 49-64.
- 540 Theron, A., Rognon, A. and Pages, J. R. (1998). Host choice by larval parasites: a study of
- 541 Biomphalaria glabrata snails and Schistosoma mansoni miracidia related to host size.
- 542 Parasitology Research 84, 727-732.
- 543 Theron, A., Sire, C., Rognon, A., Prugnolle, F. and Durand, P. (2004). Molecular ecology
- 544 of *Schistosoma mansoni* transmission inferred from the genetic composition of larval and
- adult infrapopulations within intermediate and definitive hosts. *Parasitology* **129**, 571-585.
- 546 Webster, J. P. (2001). Compatibility and sex in a snail-schistosome system. *Parasitology*
- 547 **122,** 423-432.
- 548 Webster, J. P., Gower, C. M. and Blair, L. (2004). Do hosts and parasites coevolve?
- 549 Empirical support from the Schistosoma system. *American Naturalist* **164 Suppl 5**, S33-51.
- 550 Webster, J. P., Woolhouse, M.E.J. (1998). Selection and strain specificity of compatibility
- 551 between snail intermediate hosts and their parasitic schistosomes. *Evolution Int J Org*
- 552 *Evolution* **52**, 1627-1634.
- 553 Zhang, S. M., Adema, C. M., Kepler, T. B. and Loker, E. S. (2004). Diversification of Ig
- superfamily genes in an invertebrate. *Science* **305**, 251-254.
- 555

Table 1. Summary of experiments conducted.

_	_	
`	7	1
\sim	~	'

556

Strains compared	Year strains isolated from field	Year tested	Dose-response curve done?	Method used to verify infection ³
lab-vs-lab	1992	1997	Yes	Dissection
	2000	2000	No ¹	Dissection
	2005	2007	Yes ²	Dissection
wild-vs-wild	NA	2000	Yes	Dissection
	NA	2005	Yes	Shedding
wild parasites- vs-lab snails	2000 (snails)	2002	Yes	Dissection

558

¹challenges using 20 miracidia performed for first three generations of laboratory passage.

560

²challenges using 20 miracida performed every generation since the strains were established

562

³snails were either fixed and dissected to count number of mother sporocysts (Dissection) or
 were observed for cercarial shedding 30 days post exposure (Shedding).

565 **Figure legends**

- 566 Fig. 1. Head-foot region of a *Biomphalaria glabrata* fixed 15 days post-exposure to 20
- 567 miracidia of *Schistosoma mansoni*. Six well developed mother sporocysts of *Schistosoma*
- 568 *mansoni* are easily observable as white bodies within the snail tissue.
- 569
- 570 Fig. 2. Infection rates (\pm one standard error) of individual snails and mean number (bar \pm one
- 571 standard error) of mother sporocysts (MSp) per infected snail exposed to increasing doses of
- 572 *Schistosoma mansoni* miracidia (nMi). (A) Both parasites and snails were maintained in lab
- 573 for 5 years. N = 45 snails per dose. (B) Wild snails challenged with wild miracidia. Data
- 575 collected only in 2000. (C) Laboratory snails maintained for two years and then challenged

from 2 different trials in the DFO site. N = 39-50 surviving snails per dose. MSp data were

- 576 by wild miracidia. N = 39-45 surviving snails per dose.
- 577

- 578 Fig. 3. Snail infection rates (grey bar \pm one standard error) and mean number (black bar \pm
- 579 one standard error) of mother sporocysts (MSp) per infected snail after exposure to 20
- 580 miracidia. Snails and parasites were both obtained from the wild population (Field) in 2000,
- and were then tested after 1, 2, and 3 laboratory generations of passage.
- 582
- 583 Fig. 4. Example of how simple models of independent infection with low individual infection
- 584 probabilities can generate curves that look very similar to some empirical curves. (A) 2000
- 585 wild-by-wild data (circles) versus equation 1 parameterized with p = 0.11 (line). (B) 1997
- 186 lab-by-lab data (circles) versus equation 2 parameterized by f = 0.67 and p = 0.2 (line).



FIG. 1



FIG.2



FIG.3



Number of miracidia

FIG. 4



FIG. 1



FIG. 2 MS ID: PAR-2008-0080.R1 Title: Effects of laboratory culture on compatibility between snails and schistosomes



FIG.3 MS ID: PAR-2008-0080.R1 Title: Effects of laboratory culture on compatibility between snails and schistosomes



FIG. 4

MS ID: PAR-2008-0080.R1

Title: Effects of laboratory culture on compatibility between snails and schistosomes