



**HAL**  
open science

## Relative influences of DOC flux and subterranean fauna on microbial abundance and activity in aquifer sediments: new insights from $^{13}\text{C}$ -tracer experiments

Arnaud Foulquier, Laurent Simon, Franck Gilbert, François Fourel, Florian Malard, Florian Mermillod-Blondin

### ► To cite this version:

Arnaud Foulquier, Laurent Simon, Franck Gilbert, François Fourel, Florian Malard, et al.. Relative influences of DOC flux and subterranean fauna on microbial abundance and activity in aquifer sediments: new insights from  $^{13}\text{C}$ -tracer experiments. *Freshwater Biology*, 2010, 55 (7), pp.1560-1576. 10.1111/j.1365-2427.2010.02385.x . halsde-00520932

**HAL Id: halsde-00520932**

**<https://hal.science/halsde-00520932>**

Submitted on 7 Jun 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>  
Eprints ID: 6083

**To link to this article:** DOI: 10.1111/j.1365-2427.2010.02385.x  
URL: <http://dx.doi.org/10.1111/j.1365-2427.2010.02385.x>

**To cite this version:** Foulquier, Arnaud and Simon, Laurent and Gilbert, Franck and Fourel, François and Malard, Florian and Mermillod-Blondin, Florian *Relative influences of DOC flux and subterranean fauna on microbial abundance and activity in aquifer sediments: new insights from <sup>13</sup>C-tracer experiments*. (2010) *Freshwater Biology*, vol. 55 (n° 7). pp. 1560-1576. ISSN 1365-2427

Any correspondence concerning this service should be sent to the repository administrator: [staff-oatao@listes.diff.inp-toulouse.fr](mailto:staff-oatao@listes.diff.inp-toulouse.fr)

# Relative influences of DOC flux and subterranean fauna on microbial abundance and activity in aquifer sediments: new insights from $^{13}\text{C}$ -tracer experiments

ARNAUD FOULQUIER\*, LAURENT SIMON\*, FRANCK GILBERT<sup>†,‡</sup>, FRANCOIS FOUREL<sup>§</sup>, FLORIAN MALARD\* AND FLORIAN MERMILLOD-BLONDIN\*

\*UMR CNRS 5023, Ecologie des Hydrosystèmes Fluviaux, Université Claude Bernard Lyon 1, Villeurbanne Cedex, France

<sup>†</sup>Université de Toulouse, UPS, INP, EcoLab (Laboratoire d'écologie Fonctionnelle), Toulouse, France

<sup>‡</sup>CNRS, EcoLab, Toulouse, France

<sup>§</sup>UMR CNRS 5125, PaléoEnvironnements & PaléobioSphère, Université Claude Bernard Lyon 1, Villeurbanne Cedex, France

## SUMMARY

1. Aquifers are considered to be controlled bottom-up because of their dependence on organic matter supply from surface ecosystems. Microorganisms are generally assumed to form the base of the food web and to respond strongly to organic matter supply. Although the bottom-up control of microorganisms by carbon sources has been well documented, the potential top-down control of obligate groundwater invertebrates on microorganisms has never been addressed in alluvial aquifers.

2. The main aims of the present study were (i) to quantify the relative influences of the activity of a subterranean amphipod (*Niphargus rhenorhodanensis*) and the flux of dissolved organic carbon (DOC) on organic matter processing and microbial activity, biomass and abundance in slow filtration columns mimicking an alluvial aquifer, and (ii) to determine the feeding rate of *N. rhenorhodanensis* on sedimentary microbes by tracing the flux of a  $^{13}\text{C}$ -labelled source of DOC in batches (closed systems).

3. Slow filtration column experiments showed that microbial abundance, biomass and activity were primarily controlled by DOC flux, whereas the activity of *N. rhenorhodanensis* had only a slight effect on the microbial compartment. Modelling of carbon fluxes in the  $^{13}\text{C}$ -tracer experiments indicated that the feeding activity of the amphipod was too low to significantly modify microbial growth and activity.

4. Our experiments supported the hypothesis that groundwater ecosystems are controlled bottom-up. The small influence of *N. rhenorhodanensis* on the microbial compartment was probably linked to its slow metabolism. Our results highlight the need for further experiments to examine the relationship between metabolic rates of subterranean organisms and their role in ecosystem functioning.

*Keywords:* animal–microbe interactions, bottom-up control, carbon cycle, *Niphargus rhenorhodanensis*, top-down control

## Introduction

Aquatic subterranean ecosystems are heterotrophic, their functioning depending on the supply of allochthonous organic matter originating from surface ecosystems. The lack of photosynthesis (and primary production) in ground water implies that the food

---

Correspondence: Florian Mermillod-Blondin, UMR CNRS 5023, Ecologie des Hydrosystèmes Fluviaux, Université Claude Bernard Lyon 1, Bât Forel, 43 Bd 11 Novembre 1918, F-69622 Villeurbanne Cedex, France.  
E-mail: mermillo@univ-lyon1.fr

webs are truncated (Gibert & Deharveng, 2002). Theory predicts that heterotrophic and organic matter-limited environments such as groundwater ecosystems are controlled bottom-up because the quantity and quality of organic matter entering the system determine food web structure (Malard *et al.*, 1994; Notenboom, Plénet & Turquin, 1994; Datry, Malard & Gibert, 2005). The soil and vadose zone (between the ground surface and the water table) act as strong physical, chemical and biological filters that dramatically reduce the transfer of organic detritus (particulate organic matter – POM) from surface to ground water. Consequently, dissolved organic carbon (DOC) is the main source of food for life in aquifers (e.g. Starr & Gillham, 1993). Accordingly, Simon, Benfield & Macko (2003) showed that microorganisms which process DOC represented the main food resource for higher trophic levels in a karstic system. In the hyporheic zone, several studies (e.g. Findlay *et al.*, 1993; Claret & Fontvieille, 1997; Craft, Stanford & Pusch, 2002) have demonstrated the significance of DOC availability for microbial growth, with microorganisms being a potential food source for invertebrates (Bärlocher & Murdoch, 1989). In a phreatic aquifer, Datry *et al.* (2005) reported increased densities of groundwater invertebrates in zones enriched with DOC and suggested that a DOC-induced enrichment of microbial biomass stimulated the abundances of groundwater invertebrates.

Although this 'bottom-up view' of groundwater ecosystems is largely accepted, there have been no attempts to quantify the influence of the obligate groundwater fauna on the microbial compartment and DOC consumption and assimilation in alluvial groundwater ecosystems (Gibert *et al.*, 2009). In a recent review, Boulton *et al.* (2008) pointed out that obligate groundwater invertebrates should affect many of the microbial processes involved in the C and N cycles in a similar way to that observed for surface-dwelling aquatic invertebrates. In sediments, grazing activities of invertebrates may enhance biofilm productivity (e.g. Traunspurger, Bergtold & Goedkoop, 1997), whereas bioturbation activities (burrowing, gallery building, and faecal pellet production) may indirectly influence microbial communities through modifications of physicochemical conditions (Danielopol, 1989; Mermillod-Blondin & Rosenberg, 2006). However, clear evidence of the role

of invertebrates in microbial processes in sediment has essentially been limited to surface-dwelling species that lack many of the behavioural and physiological adaptations to food limitation shared by most obligate groundwater invertebrates (Hervant, Mathieu & Barré, 1999a). Thus, the main aim of the present study was to test whether a model obligate groundwater species, the amphipod *Niphargus rhenorhodanensis* Schellenberg had a significant influence on the microbial compartment and organic matter processing in aquifer sediments. This species was selected because it is the most abundant and widely distributed subterranean macroinvertebrate in the alluvial aquifer of the Rhône River (Dole-Olivier & Marmonier, 1992). Two laboratory experiments were developed (i) to quantify the relative influences of amphipod activities (top-down control) and the flux of DOC (bottom-up control) on the microbial compartment and DOC processing in slow filtration columns mimicking an alluvial aquifer habitat and (ii) to determine the feeding rate of *N. rhenorhodanensis* on attached bacteria by tracing the flux of a  $^{13}\text{C}$ -labelled source of DOC in batches.

## Methods

### *Collection of sediments*

Gravel collected from the Rhône River was sieved manually to select particle sizes ranging from 5 to 8 mm and then was cleaned with deionised water before being dried at 60 °C. For the two experiments, 10 kg of fresh sand was collected at a depth of 1 m below the bed of a gravel pit intersecting the water table of a glaciofluvial aquifer using the Bou-Rouch method (Bou & Rouch, 1967). DOC concentrations in ground water collected from this glaciofluvial aquifer ranged from 0.4 to 0.9 mg L<sup>-1</sup>. Groundwater temperature was 14.5 ± 0.3 °C, and dissolved oxygen (DO) concentrations ranged from 5.8 to 6.7 mg L<sup>-1</sup>. Particulate organic carbon contained in the collected sand was 0.42 ± 0.03 mg g<sup>-1</sup> of sediment dry mass, and the abundance of bacteria associated with sand was 2.1 10<sup>8</sup> bacteria g<sup>-1</sup> of sediment dry mass. After collection, the sand was manually sieved to select particle sizes ranging from 100–1000 µm and then kept in an oxygenated water bath in the laboratory two days before experiments started.

### Collection of subterranean organisms


The obligate groundwater amphipod *Niphargus rhenorhodanensis* (dry weight = 1.0–1.3 mg), which inhabits both karstic and alluvial aquifers, is an opportunistic species showing a generalist feeding strategy (Ginet, 1960; Danielopol, 1989). Individuals of *N. rhenorhodanensis* were collected using traps buried at a depth of 5 cm in the sediment of a ditch draining the alluvial aquifer of the Dombes Forest, France (45°58'28"N, 5°24'25"E, clade I in Lefebure *et al.*, 2007). To acclimatise animals to experimental conditions (temperature, granulometry and food), they were maintained in the laboratory for more than 15 days before the experiment started.

### Experiment I: relative influences of *N. rhenorhodanensis* and DOC flux on microbial compartment in slow filtration microcosms

Slow filtration column experiments were conducted to test for the influences of amphipod activity (top-down control) and the flux of DOC (bottom-up control) on microbial compartment and DOC processing in sediments. Columns had an inside diameter of 10 cm and consisted of two experimental modules (10 cm high)

topped by a third module of 5 cm in height (Fig. 1a). They were filled with inert gravel and fresh sand to a height of 15 cm. About 10 cm of water was left above the sediment surface. Masses of gravel (500, 500 and 400 g) and sand (150, 150 and 120 g) were alternately introduced into the columns to obtain a heterogeneous porous media. During sediment introduction, particle tracers (luminophores) were deposited at depths of 2.5–3.5 cm (1 g of yellow luminophores) and 6–7 cm (1 g of pink luminophores) below the sediment surface to estimate sediment reworking induced by amphipods (method described below).

Experiments were performed at constant temperature ( $15 \pm 0.5$  °C) in the dark. The columns were supplied from the top with reconstituted groundwater ( $96 \text{ mg L}^{-1} \text{ NaHCO}_3$ ,  $39.4 \text{ mg L}^{-1} \text{ CaSO}_4 \cdot 2\text{H}_2\text{O}$ ,  $60 \text{ mg L}^{-1} \text{ MgSO}_4 \cdot 7\text{H}_2\text{O}$  and  $4 \text{ mg L}^{-1} \text{ KCl}$ ,  $19 \text{ mg L}^{-1} \text{ Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ ) using a peristaltic pump controlling a constant infiltration flow rate of  $2 \text{ mL min}^{-1}$ . Supplied water was aerated to maintain concentrations of DO between  $8.5$  and  $9.5 \text{ mg L}^{-1}$  at the inlet of the columns. The experiment was designed to couple three modalities of DOC flux (i.e. 60, 180 and  $300 \mu\text{g h}^{-1}$  of C by modifying the concentration of easily degradable DOC sodium acetate supplied to columns) with two modalities of fauna (absence or



**Fig. 1** Schematic representation of slow filtration columns indicating the positions of sampling (a) and experimental design of the experiment I in slow filtration columns (b). SFC = slow filtration column.

presence of 10 individuals of *N. rhenorhodanensis*) to test for the interaction between DOC supply and the activity of *N. rhenorhodanensis* on organic matter processing and microbial parameters (Fig. 1b). A total of six experimental treatments were tested with three replicate columns per treatment. Dissolved acetate supplied to all columns was enriched with 0.2% of  $^{13}\text{C}$ -marked acetate ( $^{13}\text{C}_2\text{H}_4\text{O}_2$ , 99 atom%  $^{13}\text{C}$ ; Sigma–Aldrich, Saint-Quentin Fallavier, France) to determine the assimilation of DOC in attached bacteria and amphipods. The  $\delta^{13}\text{C}$  of the DOC solution supplied to columns was 135 ‰.

After 1 week of column stabilisation, 10 individuals of *N. rhenorhodanensis* were introduced in half of the columns ( $n = 3$  columns per DOC treatment). Amphipod density in slow filtration columns (i.e. 8.4 individuals  $\text{L}^{-1}$  of sediment) was similar to that measured in artificial substrata inserted in the bed sediment of streams ( $7.2 \pm 2.6$  individuals  $\text{L}^{-1}$  of sediment, Mathieu & Essafi-Chergui, 1990). Measurements of DOC and DO were made at four depths (5 cm above the sediment surface, and 1, 5 and 10 cm below the sediment surface) on days 0 (before addition of amphipods in the system), 6, 10, 14, 18, 22 and 26 after the addition of *N. rhenorhodanensis* to determine the removal rates of DOC and DO (aerobic respiration) in slow filtration columns (Fig. 1a). After the last measurements (day 26), columns were dismantled, the water layer was removed and columns were opened to sample sediment (Fig. 1a). The top 11 cm of fresh sediment were sampled in 0.5 cm thick slices. Each layer was sieved to remove gravel and invertebrates, homogenised, and a 1 g sub-sample was used for sediment reworking analyses (see below). Remaining fresh sediments corresponding to three layers (0–2, 4–6 and 9–11 cm below the sediment surface) were used to analyse the  $^{13}\text{C}/^{12}\text{C}$  ratio of bacteria (incorporation of acetate in bacteria, see model in Data analysis section), bacterial abundance (numbers of bacteria and active eubacteria), microbial activities (hydrolytic activity, dehydrogenase activity) and biochemical markers of biofilm biomass (carbohydrates and proteins). Carbohydrates and proteins were used to estimate biomasses of extracellular polymeric substances and bacterial cells, respectively (Foulquier A., Mermillod-Blondin F., Malard F. & Gibert J., *submitted*). Individuals of *N. rhenorhodanensis* were recovered from columns to analyse their  $^{13}\text{C}/^{12}\text{C}$  ratio and calculate the incorporation rate of  $^{13}\text{C}$ -enriched micro-

bial biomass in animals (see model in Data analysis section). After collection, *N. rhenorhodanensis* were kept 48 h in glass bowls with tap water to clear their gut contents before stable isotope analyses.

#### *Experiment II: incorporation of DOC in the food web*

We designed a second experiment with simplified experimental units to quantify the assimilation of DOC in attached bacteria and subterranean amphipods. A total of 36 erlenmeyer flasks were used to measure bacterial growth and the temporal dynamics of  $^{13}\text{C}/^{12}\text{C}$  ratio in bacteria and amphipods. Flasks (250 mL) were filled with 30 g dry mass of fresh sand (collected and treated as in Experiment I) and 160 mL of reconstituted ground water (ionic composition as in Experiment I). Reconstituted ground water was enriched with  $^{13}\text{C}$ -labelled acetate ( $^{13}\text{C}_2\text{H}_4\text{O}_2$ , 99 atom%  $^{13}\text{C}$ ) to fit an initial concentration of 25  $\text{mg L}^{-1}$  of DOC and a  $^{13}\text{C}/^{12}\text{C}$  ratio of 47 atom%. The concentration of DOC (25  $\text{mg L}^{-1}$ ) used in this experiment was higher than the DOC concentrations used in slow filtration columns (<3  $\text{mg L}^{-1}$ ). However, the application of a high concentration of  $^{13}\text{C}$ -labelled DOC was necessary to obtain significant  $^{13}\text{C}$  enrichments in bacteria and amphipods for the calculation of carbon assimilation rates. Four individuals of *N. rhenorhodanensis* were introduced in half of the flasks ( $n = 18$ ) to measure their capacity to modify DOC incorporation in attached bacteria and to feed and incorporate bacterial carbon in their tissues. At six dates during the experiment (0, 12, 24, 48, 72 and 96 h), three flasks of each treatment (absence or presence of *N. rhenorhodanensis*) were killed to determine DOC concentrations in the overlying water, number of bacteria and active eubacteria on the sediment, and the  $^{13}\text{C}/^{12}\text{C}$  ratios of the bacteria and amphipods (used for calculation of assimilation rates, see model in Data analysis section). As indicated for experiment I, collected amphipods were kept 48 h in glass bowls with tap water to clear their gut contents.

#### *Physical, chemical and microbial analyses*

*Sediment reworking analyses.* Sediment reworking by amphipods was assessed using luminophores (natural sediment particles of 350–500  $\mu\text{m}$  in size, dyed with a fluorescent paint) as particle tracers (Gerino *et al.*, 1998). Each sub-sample of fresh sediment

collected at the end of the experiment from 0.5 cm thick slices was lyophilised for luminophore counting. Detection of luminophores was performed under UV light (digital camera Olympus C-2500L; Olympus France, Rungis, France) for automatised counting on acquired images (image analysis software Image-Pro Plus). Luminophore quantities were reported in terms of quantity of dry sand collected in each slice.

*Dissolved oxygen and DOC analyses.* Dissolved oxygen measurements in slow filtration columns were made with an oxygen microsensors probe (Unisense, Aarhus, Denmark) directly connected to the four lateral water outlets, thereby preventing any contact with atmospheric oxygen. Water samples for DOC analyses were filtered through HAWP filters (porosity: 0.45  $\mu\text{m}$ ; Millipore, Billerica, MA, U.S.A.) and analysed with a total carbon analyser (multi N/C<sup>®</sup> 3100; Analytik Jena, Jena, Germany) based on thermocatalytic oxidation and MC-NDIR detection after removing inorganic carbon with hydrochloric acid (5  $\mu\text{L mL}^{-1}$ ) and CO<sub>2</sub> stripping under 15 min oxygen flow.

*Stable isotope analysis.* Sediment samples (with attached bacteria) and amphipods were oven dried at 50 °C for at least 48 h and then crushed using a mortar and pestle. About 500 mg of dry sediments were placed in pre-cleaned Oakridge centrifugation tubes and 2 M HCl was added for 12 h at room temperature to remove calcite. After centrifugation at 2500 g during 5 min, the supernatant was discarded; sediments were rinsed three times with ultrapure water and oven dried at 50 °C. An amount of 2.5 mg of dry sediments were weighted in tin capsules for stable isotope analysis. Dry amphipods were placed in glass vials and acidified with 400  $\mu\text{L}$  of 2 M HCl to remove calcium carbonate. Samples were oven dried at 50 °C without subsequent rinse. A total of 250  $\mu\text{g}$  of dry amphipod were weighted in tin capsules for stable isotope analysis.

Stable isotope ratios of carbon (<sup>13</sup>C/<sup>12</sup>C) were measured by continuous flow stable isotope ratio mass spectrometer (CF-IRMS) using a GVI Isoprime mass spectrometer interfaced with a Eurovector Euro-EA3028-HT elemental analyser. <sup>13</sup>C/<sup>12</sup>C ratios were expressed as  $\delta$  in part per thousand (‰) and referenced to V-PDB standard. The analytical precision achieved for tyrosine standards analysed along with the samples was better than 0.1‰ ( $\pm$  standard deviation).

*Microbial analyses.* The DNA intercalating dye (DAPI) and a Cy3 probe (EUB 338, eubacteria) were used on sediment samples to determine the total number of bacteria and the percentages of active eubacteria. Sediment samples (1 g) were fixed, homogenised, spotted on slides and hybridised according to Mermillod-Blondin *et al.* (2005). Slides were mounted with Citifluor solution (Citifluor Ltd, London, U.K.), and the preparations were examined at 1000 $\times$  magnification with a BH2-RFCA Olympus microscope fitted for epifluorescence with a high-pressure mercury bulb (50 W) and filter sets BP 405 (for DAPI) and BP 545 (for Cy3). Bacteria from the samples were analysed in 20 fields per sample with up to 30 cells per field. Numbers of DAPI and Cy3 bacteria were counted separately from the same field to determine the percentages of active eubacteria (% Cy3 bacteria/DAPI bacteria). Total numbers of bacteria were expressed as number of cells g<sup>-1</sup> dry weight (DW) of sediment.

The 2-(p-iodophenyl)-3-(p-nitrophenyl)-5-phenyl tetrazolium chloride (INT) was used to measure dehydrogenase activity as modified from Houridavignon, Relexans and Etcheher (1989). Sediment samples (1 g fresh weight) were incubated into a 0.02% INT solution (final solution) for 2 h at 15 °C and then filtered on a nylon membrane (0.22  $\mu\text{m}$ , MSI). Controls were prepared by adding formaldehyde (2% final) in INT solution. Extraction of INT formazan was made in vials containing 5 mL of methanol. Each vial was sonicated at 100 W during two periods of 60 s using a sonicator fitted with a 2-mm-diameter probe (Sonicator XL 2020; Misonix Inc., Farmingale, NY, U.S.A.) to increase solvent extraction yield (Maurines-Carboneill *et al.*, 1998). The INT formazan extract was measured by a spectrophotometer adjusted at 480 nm against control blank. The quantity of INT formazan was computed by using the molar extinction coefficient of 18 000 M<sup>-1</sup> cm<sup>-1</sup> at 480 nm and was expressed as  $\mu\text{mol}$  of INT h<sup>-1</sup> g<sup>-1</sup> DW of sediment.

Hydrolytic activity was estimated using the fluorescein diacetate (FDA) hydrolysis method (Jørgensen, Eriksen & Jensen, 1992). Sediment samples (1 g fresh weight) were placed into 3 mL of a pH 7.6 phosphate buffer solution with 0.15 mL of 4.8 mM FDA solution. The incubation was maintained for 1 to 3 h until a green colouration of fluorescein appeared. The reaction was stopped by freezing the sample after the addition of 4.5 mL of a solution of HgCl<sub>2</sub>

(400 mg L<sup>-1</sup>). Fluorescein concentration was estimated from the absorbance of the filtered supernatant (0.45 μm, Millipore) measured at 490 nm and was expressed as μmol of FDA h<sup>-1</sup> g<sup>-1</sup> DW of sediment.

Total proteins on sediment were measured according to the Lowry method modified by Peterson (1977), using the Sigma Protein Assay Kit (P 5656 Sigma Diagnostics, St Louis, MO, USA). Concentration was expressed as mg of protein g<sup>-1</sup> DW of sediment.

Carbohydrates were quantified on dry sediment (24 h at 60 °C in an oven) using the Dubois method (Dubois *et al.*, 1956). A homogenised sample of dry sediment (0.25 g) was mixed with 1 mL of a phenol solution (10%) and 5 mL of sulphuric acid (95%). Incubation was performed at room temperature in the dark for 1 h. Absorbance of the supernatant was measured at 495 nm, and carbohydrate content was expressed as mg of carbohydrates g<sup>-1</sup> DW of sediment.

#### Data analysis

*Experiment I.* The quantification of luminophore redistribution following sediment reworking by *N. rhenorhodanensis* was realised using a model inspired by the gallery-diffusion model of François *et al.* (2002). This model describes both the diffusive-like mixing of particles in the region of intense burrowing activity ( $D_b$ ) and the non-local mixing pattern ( $r$ ) associated with a biologically induced transfer of particles from one layer to another in a discontinuous pattern (i.e. a non-continuous transport; Boudreau, 1986; Meysman, Boudreau & Middelburg, 2003). Detailed information about this model can be found in Gilbert *et al.* (2007). Biodiffusion-like coefficients of particles obtained for yellow and rose luminophores were compared among DOC and fauna treatments using a two-way analysis of variance (ANOVA). Tukey's *post hoc* tests were performed to identify significant pairwise differences between treatments.

Concentrations of DO and DOC measured on day 26 at 4 depths in slow filtration columns were used to calculate the removal rate of DOC and DO in three sediment layers (i.e. 0–2, 2–5 and 5–10 cm below the sediment surface) as follows:

$$RR = (\Delta_C \times Q) / (V \times P) \quad (1)$$

Where:

RR: Removal rate of DOC and DO in mg h<sup>-1</sup> L<sup>-1</sup> sediment

$\Delta_C$ : Difference in DOC or DO concentration (mg L<sup>-1</sup>) between two consecutive layers

Q: water discharge (L h<sup>-1</sup>) through the sediment layer

V: volume of the sediment layer (L)

P: porosity

A three-way ANOVA was used to test for the effects of depth, DOC flux and amphipods on the removal rates of DO and DOC, microbial parameters (abundance, activities and biochemical markers) and <sup>13</sup>C/<sup>12</sup>C ratio of attached bacteria. A one-way ANOVA was used to compare the <sup>13</sup>C/<sup>12</sup>C ratio of amphipod tissues collected at the end of the experiment between DOC treatments. Tukey's *post hoc* tests were performed to identify significant pairwise differences between treatments.

*Experiment II.* Student's *t*-tests were performed on measurements made at 0 h to verify that there were no differences in DOC concentrations, bacterial counts and <sup>13</sup>C/<sup>12</sup>C ratios of attached bacteria between the 2 series of flasks ( $n = 3$ ) attributed to each fauna treatment before addition of *N. rhenorhodanensis*. After 0 h, a two-way ANOVA with fauna (presence or absence of *N. rhenorhodanensis*) and time (12, 24, 48, 72 and 96 h) was used to test for differences in the temporal dynamics of DOC, bacterial counts and <sup>13</sup>C/<sup>12</sup>C ratios of attached bacteria between fauna treatments. A one-way ANOVA was used to compare the <sup>13</sup>C/<sup>12</sup>C ratios of amphipod tissues among dates.

*Modelling of carbon fluxes.* The modelling of carbon fluxes between DOC, attached bacteria and *N. rhenorhodanensis* was performed using a similar approach as in Mauclair *et al.* (2003). Bacteria growth was assumed to follow Monod kinetics (Monod, 1942). The bacterial uptake of DOC ( $C_{DOC}$ ) over time  $t$  in the experimental system was described by the following differential equation:

$$\frac{dC_{DOC}}{dt} = -\frac{1}{Y_1} \mu_1 \frac{C_{DOC}}{K_S + C_{DOC}} B \quad (2)$$

where  $B$  was the carbon biomass of bacteria (g C bacteria),  $K_S$  was the Monod constant (mg C),  $\mu_1$  was the maximum growth rate of bacteria (day<sup>-1</sup>) and  $Y_1$  was the carbon conversion efficiency between acetate and bacterial biomass (g C bacteria g<sup>-1</sup> C<sub>DOC</sub>). The

feeding rate of amphipods on bacteria was assumed to depend on the bacterial biomass and we neglected biomass changes because of natural decay, which yielded the equation:

$$\frac{dN}{dt} = \mu_2 BN \quad (3)$$

with  $N$  as amphipod biomass (g C amphipods) and  $\mu_2$  as growth rate ( $\text{g}^{-1}$  C bacteria  $\text{day}^{-1}$ ).

Total change of bacteria biomass was then calculated as follows:

$$\frac{dB}{dt} = \left( \mu_1 \frac{C_{\text{DOC}}}{K_s + C_{\text{DOC}}} B \right) - \mu_3 B - \frac{1}{Y_2} \mu_2 BN \quad (4)$$

where  $\mu_3$  was the bacteria natural decay rate ( $\text{day}^{-1}$ ) and  $Y_2$  was the carbon conversion efficiency between bacteria and amphipod biomasses (g C amphipods  $\text{g}^{-1}$  C bacteria).

$^{13}\text{C}$  fluxes were modelled to describe changes in the  $^{13}\text{C}/^{12}\text{C}$  ratio of attached bacteria and amphipod tissues. We assumed that there was no isotopic fractionation between DOC and attached bacteria, and between bacteria and amphipods during C assimilation. This assumption was a reasonable simplification because C isotope fractionations during aerobic metabolism are generally low (0 to few ‰), thus negligible relatively to the high  $^{13}\text{C}$  enrichment of this study. The change of  $^{13}\text{C}$  mass in DOC was given by:

$$\frac{d^{13}C_{\text{DOC}}}{dt} = -\frac{1}{Y_1} \mu_1 \frac{^{13}C_{\text{DOC}}}{K_s + C_{\text{DOC}}} B \quad (5)$$

while the evolution through time of  $^{13}\text{C}$  in bacteria ( $^{13}C_B$ ) and amphipods ( $^{13}C_N$ ) was expressed as:

$$\frac{d^{13}C_B}{dt} = \left( \mu_1 \frac{^{13}C_{\text{DOC}}}{K_s + C_{\text{DOC}}} B \right) - \mu_3 ^{13}C_B - \frac{1}{Y_2} \mu_2 ^{13}C_B N \quad (6)$$

$$\frac{d^{13}C_N}{dt} = \mu_2 ^{13}C_B N \quad (7)$$

The carbon mass and the isotopic compositions of the DOC, bacteria and amphipods were computed at each time step by solving the system of differential equations (2–7) that described the carbon and  $^{13}\text{C}$  mass balances using a fourth-order Runge-Kutta numerical integration procedure (Press *et al.*, 1992). To compare the result of the model to the experimen-

tal data, the carbon isotope composition of total organic carbon attached to sediment (comprising the organic carbon contained in bacterial cells and bacterial exopolymeric substances) was calculated using a simple mass-balance equation:

$$\delta^{13}C_{\text{TOC}} = \frac{B\delta^{13}C_B + C_{\text{TOC}}^i \delta^{13}C_{\text{TOC}}^i}{B + C_{\text{TOC}}^i} \quad (8)$$

where  $C_{\text{TOC}}^i$  was the total organic carbon attached to sediments at the start of the experiment.


Cell bacterial biomass at the start of the experiment was estimated from DAPI counts using a conversion factor of 30 fg of C  $\text{cell}^{-1}$  (Fukuda *et al.*, 1998; Troussellier *et al.*, 1997). Invertebrates contained about 500  $\mu\text{g}$  of C per individual at  $t = 0$ .

Unknown parameters of the model ( $\mu_1$ ,  $\mu_2$ ,  $\mu_3$ ,  $K_s$ ,  $Y_1$  and  $Y_2$ , Table 1) were calculated by adjusting the simulation results to measured DOC concentrations,  $\delta^{13}\text{C}$  values of total organic carbon and  $\delta^{13}\text{C}$  of amphipod tissues. The quality of the fit between modelled and measured data was evaluated by calculating the sum of the squared differences between observed and calculated values.

For all parametric tests, homoscedasticity and data normality were checked using Levene's tests and Shapiro's tests, respectively. Whenever necessary, data were ln or square-root transformed to homogenise variances and fit data normality. For variables

**Table 1** Biodiffusion-like coefficients (Db) calculated for luminophores deposited in two depth layers of the columns for the six experimental treatments. DOC1 = 60  $\mu\text{g}$  C  $\text{h}^{-1}$ , DOC2 = 180  $\mu\text{g}$  C  $\text{h}^{-1}$ , and DOC3 = 300  $\mu\text{g}$  C  $\text{h}^{-1}$ . Values are means (SD).  $n = 3$  columns per treatment

Depth of luminophore deposition (cm)	Treatment	Db ( $\text{cm}^2 \text{ year}^{-1}$ )
2.5–3.5	DOC1 * control	0.76 (0.39)
	DOC1 * <i>N. renorhodanensis</i>	0.52 (0.32)
	DOC2 * control	0.36 (0.17)
	DOC2 * <i>N. renorhodanensis</i>	0.47 (0.40)
	DOC3 * control	0.15 (0.02)
	DOC3 * <i>N. renorhodanensis</i>	0.17 (0.05)
6.0–7.0	DOC1 * control	0.85 (0.19)
	DOC1 * <i>N. renorhodanensis</i>	1.07 (0.46)
	DOC2 * control	1.29 (0.66)
	DOC2 * <i>N. renorhodanensis</i>	0.84 (0.28)
	DOC3 * control	1.09 (0.76)
	DOC3 * <i>N. renorhodanensis</i>	0.84 (0.14)



**Fig. 2** DO and DOC removal rates, and microbial activities, biomasses and abundances measured at three layers in columns for the six tested treatments (3 DOC flux treatment × 2 animal treatments) in experiment I. Symbols for DOC supply: DOC1 = 60  $\mu\text{g C h}^{-1}$ , DOC2 = 180  $\mu\text{g C h}^{-1}$  and DOC3 = 300  $\mu\text{g C h}^{-1}$ . Symbols for depth layers: H1 = 0–2 cm, H2 = 4–6 cm and H3 = 9–11 cm below the sediment surface. Values are means  $\pm$  SD ( $n = 3$  columns per treatment).

expressed as percentages (% of active eubacteria), data were arcsine transformed. All statistical analyses were performed using Statistica 5™ (Statsoft, Tulsa, OK, U.S.A.).

## Results

### *Experiment I: relative influences of *N. rhenorhodanensis* and DOC flux on microbial compartment in slow filtration microcosms*

The luminophore vertical profiles were diffusive-like shapes, indicating a preferential occurrence of random mixing events over short distances in the columns. This induced the production of diffusion-like coefficients ( $D_b$ ) and the absence of non-local coefficients ( $r$ ) by the gallery-diffusor model. The diffusion-like coefficients ( $D_b$ ) were not significantly affected by the presence and activity of the amphipods (Table 1, two-way ANOVA, fauna effect,  $F_{(1,12)} < 0.4$ ,  $P > 0.5$  for the two investigated sediment layers).

Most variables showed no significant influence of *N. rhenorhodanensis* on the microbial compartment (Fig. 2, Table 2). Amphipods had only significant positive effects on DO removal rate and dehydrogenase activity in slow filtration columns, but these effects depended on depth as well as on 'depth  $\times$  DOC flux' interactions for DO removal rate (Table 2). For instance, *Niphargus rhenorhodanensis* increased DO removal rate by 97% in the top 2 cm of sediment columns receiving a DOC flux of  $60 \mu\text{g C h}^{-1}$ , whereas it had no significant effect ( $-5.4\%$ ) at the same depth in the treatment with a DOC flux of  $180 \mu\text{g C h}^{-1}$  (Fig. 2).

On the contrary, the increase in DOC flux had a significant positive effect (Table 2, Fig. 2) on microbial processes (DO removal, DOC removal, and dehydrogenase and hydrolytic activities), microbial biomass (proteins and carbohydrates) and bacterial counts (total number of cells and percentage of active eubacteria). In the top 2 cm of sediments, the DOC removal rate, DO consumption, protein concentration and bacterial abundance increased by 552, 389, 129 and 102%, respectively, as the DOC flux increased from 60 to  $180 \mu\text{g C h}^{-1}$ . The influence of DOC supply on microbial parameters decreased with increasing depth in slow filtration columns except for hydrolytic activity (Table 2). As a consequence of the influence of

vertical DOC fluxes on microorganisms, all microbial variables were positively correlated with the DOC removal rates measured in slow filtration columns (Table 3).

The  $\delta^{13}\text{C}$  value of total organic carbon attached to sediment also increased significantly with increasing DOC flux and decreased with depth (Fig. 3a, Table 2). This  $\delta^{13}\text{C}$  signature was highly correlated with the DOC removal rate and the microbial variables (Table 3) indicating that carbon from acetate was incorporated into bacterial biomass. *N. rhenorhodanensis* did not have a significant influence on  $\delta^{13}\text{C}$  of total organic carbon attached to sediment (Table 2) but  $\delta^{13}\text{C}$  values of amphipods showed a higher incorporation of organic carbon in their tissues for the highest DOC treatment (Fig. 3b, one-way ANOVA,  $F_{(3,9)} = 7.4$ ,  $P < 0.01$ ).

### *Experiment II: incorporation of DOC in the food web*


DOC decreased rapidly with time (Fig. 4a, Table 4) owing to consumption by the microbial compartment, the abundance of which concomitantly increased over time (Fig. 4b, Table 4). The percentage of active eubacteria also evolved significantly during the experiment but to a lesser extent than the total number of bacteria (Fig. 4c). The occurrence of *N. rhenorhodanensis* did not significantly affect the microbial compartment although amphipods significantly delayed the DOC decrease during the experiment (Fig. 4a, Table 4).

A significant increase in  $\delta^{13}\text{C}$  was measured in bacteria attached to sediment (Fig. 5b, two-way ANOVA, time effect,  $F_{(4,18)} = 67.3$ ,  $P < 0.001$ ) and tissues of amphipods (Fig. 5c, one-way ANOVA,  $F_{(5,12)} = 39.9$ ,  $P < 0.001$ ). The best model fit of DOC concentration (Fig. 5a) and  $^{13}\text{C}/^{12}\text{C}$  ratio of total organic carbon associated with sediments (Fig. 5b) and amphipods (Fig. 5c) was obtained with parameter values reported in Table 5. According to differences in DOC decrease with time between control and fauna treatments ('time  $\times$  fauna' effect in Table 4), a lower half velocity constant ( $K_s$ ) and a higher maximum growth rate of bacteria ( $\mu_1$ ) were estimated in the control treatment in comparison with the fauna treatment (Table 5). Although amphipods delayed DOC decrease and reduced bacterial growth rate, the conversion coefficient of DOC into bacterial carbon ( $Y_1$ ) was increased by 18% with *N. rhenorhodanensis*



**Table 3** Correlations between biogeochemical processes, microbial variables and sedimentary  $\delta^{13}\text{C}$  obtained in slow filtration columns (DOC RR: dissolved organic carbon removal rate; DO RR: dissolved oxygen removal rate; INT: dehydrogenase activity; FDA: hydrolytic activity, BA: bacterial abundance; % AE: percentage of active eubacteria)


Variables	DOC RR	DO RR	INT	FDA	Proteins	Carbohydrates	BA	% AE
DO RR	$R = 0.912$ $P < 0.001$							
INT	$R = 0.786$ $P < 0.001$	$R = 0.805$ $P < 0.001$						
FDA	$R = 0.368$ $P < 0.05$	$R = 0.425$ $P < 0.01$	$R = 0.420$ $P < 0.01$					
Proteins	$R = 0.529$ $P < 0.001$	$R = 0.609$ $P < 0.001$	$R = 0.580$ $P < 0.001$	$R = 0.873$ $P < 0.001$				
Carbohydrates	$R = 0.801$ $P < 0.001$	$R = 0.824$ $P < 0.001$	$R = 0.816$ $P < 0.001$	$R = 0.463$ $P < 0.01$	$R = 0.694$ $P < 0.001$			
BA	$R = 0.646$ $P < 0.001$	$R = 0.738$ $P < 0.001$	$R = 0.717$ $P < 0.001$	$R = 0.760$ $P < 0.001$	$R = 0.901$ $P < 0.001$	$R = 0.835$ $P < 0.001$		
% AE	$R = 0.565$ $P < 0.001$	$R = 0.652$ $P < 0.001$	$R = 0.624$ $P < 0.001$	$R = 0.747$ $P < 0.001$	$R = 0.897$ $P < 0.001$	$R = 0.757$ $P < 0.001$	$R = 0.908$ $P < 0.001$	
Sedimentary $\delta^{13}\text{C}$	$R = 0.866$ $P < 0.001$	$R = 0.957$ $P < 0.001$	$R = 0.779$ $P < 0.001$	$R = 0.485$ $P < 0.01$	$R = 0.630$ $P < 0.001$	$R = 0.772$ $P < 0.001$	$R = 0.740$ $P < 0.001$	$R = 0.661$ $P < 0.001$



**Fig. 3** Changes in  $\delta^{13}\text{C}$  of total organic carbon attached to sediment (a) and tissues of *N. rhenorhodanensis* (b) during the course of the experiment I. Values are means  $\pm$  SD ( $n = 3$  columns per treatment).

In slow filtration columns, we detected significant increases in oxygen consumption and dehydrogenase activity of bacteria with *N. rhenorhodanensis*. With a DOC flux of  $60 \mu\text{g h}^{-1}$ , amphipods produced an increase in oxygen consumption of  $1.3 \text{ mg of O}_2 \text{ h}^{-1} \text{ L}^{-1}$  of sediment in the uppermost 2 cm of sediments that could not be attributed to the sole respiration of 10 individuals of *N. rhenorhodanensis* ( $0.04 \text{ mg of O}_2 \text{ h}^{-1} \text{ L}^{-1}$  of sediment calculated with respiration data of Hervant *et al.*, 1997). Such results indicate that feeding and/or bioturbation activity of *N. rhenorhodanensis* apparently had a positive influence on respiratory activity of bacteria. In batches, the occurrence of amphipods also delayed the DOC decrease (increase in  $K_s$  and decrease in  $\mu_1$ ) and increased the efficiency of DOC incorporation into bacteria ( $Y_1$ ). The amphipod-induced modifications of carbon conversion efficiency  $Y_1$  in batches and dehydrogenase activity in slow filtration columns suggested that amphipods could have slightly influenced the physiology of the microbial communities. However, this effect of *N. rhenorhodanensis* on microbial function was not associated with a significant change in the biomass (proteins and carbohydrates) and abundances of bacteria (total number and number of active bacteria). Modelling of carbon fluxes in batches clearly indicated that the feeding rate of *N. rhenorhodanensis* on sedimentary biomass was too low to influence significantly the bacterial biomass (top-down effect). The fast and efficient assimilation of DOC by





**Fig. 5** Changes in DOC concentrations (a) and  $\delta^{13}\text{C}$  values of total organic carbon attached to sediment (b) and *N. rhenorhodanensis* (c) during experiment II. Values are means  $\pm$  SD ( $n = 3$  flasks per treatment). Curves represent modelling results with (dashed lines) and without *N. rhenorhodanensis* (plain lines).

were fed (only 2.5  $\mu\text{g}$  being incorporated in animal tissues) by amphipods during the same time period.

We did not detect any significant effect of *N. rhenorhodanensis* on the physical structure of sandy sediment. Amphipods did not influence the sediment transport coefficients (calculated with luminophore tracers), although they were observed to move in large

interstices of the sediment column (visual observations on inner wall of the columns) and were collected in the whole sediment columns at the end of the experiment. The lack of sediment reworking implied that the bioturbation by amphipods did not significantly modify the living conditions for microorganisms growing on sand particles. Thus, our experiments clearly indicated that the feeding and bioturbation activities of *N. rhenorhodanensis* were much too low to significantly affect microbial processes in sediments.

Our findings differ markedly from those obtained by Edler & Dodds (1996) and Cooney & Simon (2009) with cave animals. Edler & Dodds (1996) reported a stimulation by 300–400% of attached bacteria (both abundance and activity) by the subterranean isopod *Caecidotea tridentata*, whereas organic carbon addition (D-glucose) had no effect on bacteria. They suggested that microbial activity was limited by N, rather than C, in their experiments. In such conditions, the excretion of nitrogen by isopods could have stimulated microbial abundances and activities (Edler & Dodds, 1996). In our experiments, N was not limiting because  $\text{N-NO}_3^-$  concentration in water (10 mg of  $\text{N-NO}_3^- \text{L}^{-1}$ ) was more than tenfold higher than those used by Edler & Dodds (1996). Such comparison suggests that the influence of invertebrates on microbial processes may vary as a function of nutrient limitation. Significance of invertebrates on ecosystem processes may also depend on the functional traits of the species (Mermillod-Blondin & Rosenberg, 2006). If the amphipod *N. rhenorhodanensis* had slight effect in our experiments, Cooney & Simon (2009) reported that the cave amphipod *Gammarus minus* could reduce the bacterial production on rocks, offsetting the stimulatory effect as a result of glucose amendment. The contrasting influences of *G. minus* and *N. rhenorhodanensis* on bacterial activity were probably related to differences in metabolic activities between these two species. The ecophysiology of *N. rhenorhodanensis* and *G. minus* were studied by Hervant *et al.* (1997) and Hervant, Mathieu & Culver (1999b), respectively. At a temperature of 11 °C, the obligate groundwater amphipod *N. rhenorhodanensis* had a respiration rate of 12.9  $\mu\text{mol O}_2 \text{g}^{-1}$  of dry mass  $\text{h}^{-1}$  (Hervant *et al.*, 1997). Comparatively, the amphipod *Gammarus minus* studied by Cooney & Simon (2009) exhibited a threefold higher respiration rate in normoxia at the

**Table 5** Parameters used in the model and best estimated values from the fitting procedure in control and amphipod treatments

Parameters	Definition	Estimated value	
		Control	With amphipods
$K_S$	Monod constant (half velocity constant) (mg C)	2.55	2.66
$\mu_1$	Maximum growth rate for bacteria ( $\text{day}^{-1}$ )	1.17	1.11
$\mu_2$	Growth rate parameter for amphipods ( $\mu\text{g C bacteria day}^{-1}$ )	–	0.35
$\mu_3$	Rate of natural decay ( $\text{day}^{-1}$ )	<0.05	<0.05
$Y_1$	Carbon conversion efficiency between DOC and bacteria [g C (g C bacteria) $^{-1}$ ]	0.22	0.26
$Y_2$	Carbon conversion efficiency between bacteria and <i>N. rhenorhodanensis</i> [g C bacteria (g C amphipods) $^{-1}$ ]	–	0.06

same temperature (between 40 and 45  $\mu\text{mol O}_2 \text{g}^{-1}$  of dry mass  $\text{h}^{-1}$ , Hervant *et al.*, 1999b). Consequently, *G. minus* probably needs a threefold higher grazing activity than *N. rhenorhodanensis* to acquire its energy from microorganisms. Differences in metabolic demands could thus explain why *G. minus* can exert a higher control on microbial communities than *N. rhenorhodanensis*. In the same way, Kinsey *et al.* (2007) reported that *G. minus* produced a mean leaf litter breakdown of 0.14  $\text{g}^{-1}$  of leaf  $\text{g}^{-1}$  of dry animal  $\text{day}^{-1}$ , whereas recent laboratory experiments showed that *Niphargus rhenorhodanensis* had a 20-fold lower effect on leaf litter breakdown (0.007  $\text{g}^{-1}$  of leaf  $\text{g}^{-1}$  of dry animal  $\text{day}^{-1}$ , Simon Navel, unpubl. data).

The low metabolism of *N. rhenorhodanensis* is a characteristic shared by most subterranean species (Hüppop, 1985; Hervant *et al.*, 1997). Indeed, most subterranean invertebrates have developed efficient energy-saving strategies (low metabolism, storage of energy reserves) for fitting their physiological function to food-limited conditions prevailing in most aquifers (e.g. Hüppop, 1985; Danielopol *et al.*, 1994; Hervant & Renault, 2002). Hervant *et al.* (1997) also demonstrated that *N. rhenorhodanensis*, by depressing its metabolism, could survive without feeding for periods largely exceeding 200 days. We can therefore presume that the low influence of *N. rhenorhodanensis* on microorganisms in our experiments was linked to its low metabolic demand. However, we cannot exclude that the low influence of this omnivorous amphipod on microorganisms might have resulted from a low feeding preference for sedimentary bacteria. Although experiments showed that *N. rhenorhodanensis* could be maintained several years in laboratory under a large variety of foods (leaves,

sediment, meat) (Ginet, 1960; Mathieu, 1967), knowledge about the feeding preferences of this species are lacking and we only detected small quantities of faecal pellets with fine sediment in boils used for gut content clearing (personal observation). Further experiments in the laboratory and in the field (using  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  to reconstitute food webs) are therefore greatly needed to determine the feeding preferences of *N. rhenorhodanensis* in the alluvial ecosystems. Nevertheless, the present study highlighted that, in sand-gravel sediments characteristics of the alluvial aquifer of the Rhône River, *N. rhenorhodanensis* was not able to influence significantly dissolved organic matter processing (DOC removal rate). This result was linked to the lack of significant activity of the amphipods in sandy zones where microbial processes predominantly occurred in our experimental systems (Mermillod-Blondin *et al.*, 2001). Although *N. rhenorhodanensis* is a dominant hypogean species of the alluvial aquifer of the Rhône River, other less abundant subterranean organisms that are strict microbivores and effective sediment bioturbators would have more influence on microbial processes than amphipods. In surface environments, tubificid worms such as *Tubifex tubifex* or *Limnodrilus hoffmeisteri* are known to strongly affect organic matter processing in freshwater sediments (Pelegri & Blackburn, 1995; Mermillod-Blondin *et al.*, 2001). Therefore, experiments with strict subterranean tubificid worms of the genus *Trichodrilus* (Creuzé des Châtelliers *et al.*, 2009) will be needed to more widely explore the potential influence of subterranean fauna on microbial processes in alluvial sediments. Moreover, comparisons of physiological traits and roles of tubificid worms living in both surface and subterranean habitats would allow us to determine whether it exits a

relationship between the ecological status of the animals, their metabolic rates and their roles in ecosystem functioning.

Finally, our laboratory experiments did not confirm the assumption of Boulton *et al.* (2008) according to which subterranean amphipods could have a significant influence on organic matter processing in interstitial systems. Microbial activity was tightly linked to the flux of dissolved organic matter in slow filtration columns, but the feeding and burrowing activities of *N. rhenorhodanensis* were far too low for influencing microbial processes. We suggest that the feeding preferences and the low metabolism of *N. rhenorhodanensis* probably induced by a natural adaptation to low-food conditions might have severely reduce its ability to influence its environment. Testing our hypothesis of a relationship between metabolism and functional role of subterranean fauna needs to extend our experiments to a wide range of strict microbivore organisms, including species with multiple populations living in habitats with contrasted food supply.

### Acknowledgments

This study was funded by the French programme EC2CO-Cytrix from INSU/CNRS (project 'NAPCOD'), the Institut Fédératif de Recherche No. 41 (Bio-Environnement et Santé), the urban community of Lyon (project OTHU, <http://www.graie.org/othu/>) and the Rhône-Alpes Region. We thank two anonymous referees for helpful comments on the manuscript.

### References

- Bärlocher F. & Murdoch J. (1989) Hyporheic biofilm – a potential food source for interstitial animals. *Hydrobiologia*, **184**, 61–67.
- Bou C. & Rouch R. (1967) Un nouveau champ de recherches sur la faune aquatique souterraine. *Comptes-Rendus de l'Académie des Sciences, Paris*, **265**, 369–370.
- Boudreau B.P. (1986) Mathematics of tracer mixing in sediments: I. Spatially-dependent, diffusive mixing. *American Journal of Science*, **286**, 161–198.
- Boulton A.J. (2000) The functional role of the hyporheos. *Verhandlungen International Vereinigung Limnologie*, **27**, 51–63.
- Boulton A.J., Fenwick G.D., Hancock P.J. & Harvey M.S. (2008) Biodiversity, functional roles and ecosystem services of groundwater invertebrates. *Invertebrate Systematics*, **22**, 103–116.
- Claret C. & Fontvieille D. (1997) Characteristics of biofilm assemblages in two contrasted hydrodynamic and trophic contexts. *Microbial Ecology*, **34**, 49–57.
- Cooney T.J. & Simon K.S. (2009) Influence of dissolved organic matter and invertebrates on the function of microbial films in groundwater. *Microbial Ecology*, **58**, 599–610.
- Craft J.A., Stanford J.A. & Pusch M. (2002) Microbial respiration within a floodplain aquifer of a large gravel-bed river. *Freshwater Biology*, **47**, 251–261.
- Creuzé des Châtelliers M., Juget J., Lafont M. & Martin P. (2009) Subterranean aquatic oligochaeta. *Freshwater Biology*, **54**, 678–690.
- Danielopol D.L. (1989) Groundwater fauna associated with riverine aquifers. *Journal of the North American Benthological Society*, **8**, 18–35.
- Danielopol D.L., Creuzé Des Châtelliers M., Moeszlacher F., Pospisil P. & Popa R. (1994) Groundwater contamination and its impact on groundwater animals and ecosystems. In: *Groundwater Ecology* (Eds Gibert J., Danielopol D.L. & Stanford J.A.), pp. 217–243. Academic Press, UK.
- Datry T., Malard F. & Gibert J. (2005) Response of invertebrate assemblages to increased groundwater recharge rates in a phreatic aquifer. *Journal of the North American Benthological Society*, **24**, 461–477.
- Dole-Olivier M.-J. & Marmonier P. (1992) Patch distribution of interstitial communities: prevailing factors. *Freshwater Biology*, **27**, 177–191.
- Dubois M., Gilles K.A., Hamilton J.K., Rebers P.A. & Smith F. (1956) Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, **28**, 350–356.
- Edler C.C. & Dodds W.K. (1996) The ecology of a subterranean isopod, *Caecidotea tridentata*. *Freshwater Biology*, **35**, 249–259.
- Findlay S.E.G., Strayer D., Goumbala C. & Gould K. (1993) Metabolism of stream water dissolved organic carbon in the shallow hyporheic zone. *Limnology and Oceanography*, **38**, 1493–1499.
- François F., Gerino M., Stora G., Durbec J.-P. & Poggiale J.-C. (2002) A functional approach to sediment reworking by gallery-forming macrobenthic organisms: modelling and application with the polychaete *Nereis diversicolor*. *Marine Ecology Progress Series*, **229**, 127–136.
- Fukuda R., Ogawa H., Nagata T. & Koike I. (1998) Direct determination of carbon and nitrogen contents of natural bacterial assemblages in marine environments. *Applied and Environmental Microbiology*, **64**, 3352–3358.

- Gerino M., Aller R.C., Lee C., Cochran J.K., Aller J.Y., Green M.A. & Hirschberg D. (1998) Comparison of different tracers and methods used to quantify bioturbation during a spring bloom:  $^{234}\text{Th}$ , luminophores and chlorophyll a. *Estuarine, Coastal and Shelf Science*, **46**, 531–547.
- Ghiorse W.C. & Wilson J.T. (1988) Microbial ecology of the terrestrial subsurface. *Advances in Applied Microbiology*, **33**, 107–172.
- Gibert J. & Deharveng L. (2002) Subterranean ecosystems: a truncated functional biodiversity. *BioScience*, **52**, 473–481.
- Gibert J., Culver D.C., Dole-Olivier M.J., Malard F., Christman M.C. & Deharveng L. (2009) Assessing and conserving groundwater biodiversity: synthesis and perspectives. *Freshwater Biology*, **54**, 930–941.
- Gilbert F., Hulth S., Grossi V., Poggiale J.-C., Desrosiers G., Rosenberg R., Gerino M., François-Carcaillet F., Michaud E. & Stora G. (2007) Sediment reworking by marine benthic species from the Gullmar Fjord (Western Sweden): importance of faunal biovolume. *Journal of Experimental Marine Biology and Ecology*, **348**, 133–144.
- Ginet R. (1960) Écologie, éthologie et biologie de *Niphargus*. *Annales de Spéléologie*, **15**, 127–237.
- Goldscheider N., Hunkeler D. & Rossi P. (2006) Review: microbial biocenoses in pristine aquifers and an assessment of investigative methods. *Hydrogeology Journal*, **14**, 926–941.
- Hervant F. & Renault D. (2002) Long-term fasting and realimentation in hypogean and epigean isopods: a proposed adaptive strategy for groundwater organisms. *The Journal of Experimental Biology*, **205**, 2079–2087.
- Hervant F., Mathieu J., Barré H., Simon K. & Pinon C. (1997) Comparative study on the behavioral, ventilatory, and respiratory responses of hypogean and epigean crustaceans to long-term starvation and subsequent feeding. *Comparative Biochemistry and Physiology, Part A*, **118**, 1277–1283.
- Hervant F., Mathieu J. & Barré H. (1999a) Comparative study on the metabolic responses of subterranean and surface-dwelling amphipods to long-term starvation and subsequent refeeding. *The Journal of Experimental Biology*, **202**, 3587–3595.
- Hervant F., Mathieu J. & Culver D.C. (1999b) Comparative response to severe hypoxia and subsequent recovery in closely related amphipod populations (*Gammarus minus*) from cave and surface habitats. *Hydrobiologia*, **392**, 197–204.
- Houri-Davignon C., Relexans J.C. & Etcheher H. (1989) Measurements of actual electron transport system (ETS) activity in marine sediments by incubation with INT. *Environmental Technology*, **10**, 91–100.
- Hüppop K. (1985) The role of metabolism in the evolution of cave animals. *Bulletin of the National Speleological Society*, **47**, 136–146.
- Jørgensen P.E., Eriksen T. & Jensen B.K. (1992) Estimation of viable biomass in wastewater and activated sludge by determination of ATP, oxygen utilisation rate and FDA hydrolysis. *Water Research*, **26**, 1495–1501.
- Kinsey J., Cooney T.J. & Simon K.S. (2007) A comparison of the leaf shredding ability and influence on microbial films of surface and cave forms of *Gammarus minus* Say. *Hydrobiologia*, **589**, 199–205.
- Lefébure T., Douady C.J., Malard F. & Gibert J. (2007) Testing dispersal and cryptic diversity in a widely distributed groundwater amphipod (*Niphargus rhenorhodanensis*). *Molecular Phylogenetics and Evolution*, **42**, 676–686.
- Malard F., Reygrobellet J.L., Mathieu J. & Lafont M. (1994) The use of invertebrate communities to describe groundwater flow and contaminant transport in a fractured rock aquifer. *Archiv für Hydrobiologie*, **131**, 93–110.
- Marmonier P., Fontvieille D., Gibert J. & Vanek V. (1995) Distribution of dissolved organic carbon and bacteria at the interface between the Rhône river and its alluvial aquifer. *Journal of the North American Benthological Society*, **14**, 382–392.
- Marshall M.C. & Hall R.O.J. (2004) Hyporheic invertebrates affect N cycling and respiration in stream sediment microcosms. *Journal of the North American Benthological Society*, **23**, 416–428.
- Mathieu J. (1967) Recherches sur les températures létales et les températures d'acclimatation de *Niphargus longicaudatus* (Crustacés, Amphipodes) épigés et hypogés. Doctoral Thesis, University of Lyon, 76 pp.
- Mathieu J. & Essafi-Chergui K. (1990) Le peuplement aquatique interstitiel à l'interface karst/plaine alluviale. 1. Cas d'une alimentation en eau essentiellement karstique. *Mémoires de Biospéologie*, **17**, 113–122.
- Mauclair L., Pelz O., Thullner M., Abraham W.-R. & Zeyer J. (2003) Assimilation of toluene carbon along a bacteria-protist food chain determined by  $^{13}\text{C}$ -enrichment of biomarker fatty acids. *Journal of Microbiological Methods*, **55**, 635–649.
- Maurines-Carboneill C., Pernelle J.J., Morin L., Sachon G. & Leblon G. (1998) Relevance of the INT test response as an indicator of the ETS activity in monitoring heterotrophic aerobic bacterial populations in activated sludges. *Water Research*, **32**, 1213–1221.
- Mermillod-Blondin F. & Rosenberg R. (2006) Ecosystem engineering: the impact of bioturbation on biogeochemical processes in marine and freshwater benthic habitats. *Aquatic Sciences*, **68**, 434–442.

- Mermillod-Blondin F., Gerino M., Degrange V., Lensi R., Chassé J.-L., Rard M. & Creuzé des Châtelliers M. (2001) Testing the functional redundancy of *Limnodrilus* and *Tubifex* in hyporheic sediments: an experimental study in microcosms. *Canadian Journal of Fisheries and Aquatic Sciences*, **58**, 1747–1759.
- Mermillod-Blondin F., Gerino M., Sauvage S. & Creuzé des Châtelliers M. (2004) Influence of nontrophic interactions between benthic invertebrates on river sediment processes: a microcosm study. *Canadian Journal of Fisheries and Aquatic Sciences*, **61**, 1817–1831.
- Mermillod-Blondin F., Nogaro G., Detry T., Malard F. & Gibert J. (2005) Do tubificid worms influence the fate of organic matter and pollutants in stormwater sediments? *Environmental Pollution*, **134**, 57–69.
- Meysman F.J.R., Boudreau B.P. & Middelburg J.J. (2003) Relations between local, nonlocal, discrete and continuous models of bioturbation. *Journal of Marine Research*, **61**, 391–410.
- Monod J. (1942) *Recherches sur la Croissance des Cultures Bactériennes*. Hermann & Co, Paris.
- Notenboom J., Plénet S. & Turquin M.-J. (1994) Groundwater contamination and its impact on groundwater animals and ecosystems. In: *Groundwater Ecology* (Eds Gibert J., Danielopol D.L. & Stanford J.A.), pp. 477–504. Academic Press, UK.
- Pelegri S.P. & Blackburn T.H. (1995) Effects of *Tubifex tubifex* (Oligochaeta: tubificidae) on N-mineralization in freshwater sediments, measured with  $^{15}\text{N}$  isotopes. *Aquatic Microbial Ecology*, **9**, 289–294.
- Peterson G.L. (1977) A simplification of the protein assay method of Lowry *et al.* which is more generally applicable. *Analytical Biochemistry*, **83**, 346–356.
- Press W.H., Teukolsky S.A., Vetterling W.T. & Flannery B.P. (1992) *Numerical recipes in Fortran 77*, 2nd edn. Cambridge University Press, UK.
- Simon K.S., Benfield E.F. & Macko S.A. (2003) Food web structure and the role of epilithic biofilms in cave streams. *Ecology*, **84**, 2395–2406.
- Sobczak W.V. & Findlay S. (2002) Variation in bioavailability of dissolved organic carbon among stream hyporheic flowpaths. *Ecology*, **83**, 3194–3209.
- Starr R.C. & Gillham R.W. (1993) Denitrification and organic carbon availability in two aquifers. *Ground Water*, **31**, 934–947.
- Traunspurger W., Bergtold M. & Goedkoop W. (1997) The effects of nematodes on bacterial activity and abundance in a freshwater sediment. *Oecologia*, **112**, 118–122.
- Troussellier M., Bouvy M., Courties C. & Dupuy C. (1997) Variation of carbon content among bacterial species under starvation condition. *Aquatic Microbial Ecology*, **13**, 113–119.