



**HAL**  
open science

## First report of larval stages of *Fasciola hepatica* in a wild population of *Pseudosuccinea columella* from Cuba and the Caribbean.

Alain Gutierrez, A. A. Vázquez, Y. Hevia, J. Sánchez, A. C. Correa, S. Hurtrez-Boussès, J.-P. Pointier, A. Théron

### ► To cite this version:

Alain Gutierrez, A. A. Vázquez, Y. Hevia, J. Sánchez, A. C. Correa, et al.. First report of larval stages of *Fasciola hepatica* in a wild population of *Pseudosuccinea columella* from Cuba and the Caribbean.. *Journal of Helminthology*, 2011, 85 (1), pp.109-11. 10.1017/S0022149X10000350 . halsde-00608583

**HAL Id: halsde-00608583**

**<https://hal.science/halsde-00608583>**

Submitted on 7 Sep 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# First report of larval stages of *Fasciola hepatica* in a wild population of *Pseudosuccinea columella* from Cuba and the Caribbean

A. Gutiérrez<sup>1</sup>, A.A. Vázquez<sup>1</sup>, Y. Hevia<sup>1</sup>, J. Sánchez<sup>1</sup>, A.C. Correa<sup>2</sup>,  
S. Hurtrez-Boussès<sup>3</sup>, J.-P. Pointier<sup>4</sup> and A. Théron<sup>5\*</sup>

<sup>1</sup>Laboratorio de Malacologia, IPK, Apartado 601, Marianao 13, C. Habana, Cuba: <sup>2</sup>GEMI, UMR CNRS-IRD 2724 Centre de recherche IRD, 911, Avenue Agropolis BP 6450, 34394 Montpellier, France: <sup>3</sup>Département de Biologie-Ecologie, Université de Montpellier, 4 Place E. Bataillon, 34095 Montpellier Cedex 5, France: <sup>4</sup>USR 3278 CNRS-EPHE, CRIOBE, Université de Perpignan, 52 Av. Paul Alduy, 66860 Perpignan, France: <sup>5</sup>BETM, UMR 5244 CNRS-UPVD, Université de Perpignan, 52 Av. Paul Alduy, 66860 Perpignan, France

(Accepted 19 June 2010; First Published Online 19 July 2010)

## Abstract

A wild population of the lymnaeid snail *Pseudosuccinea columella* infected by larval stages of *Fasciola hepatica* was discovered in the Pinar del Río Province, Cuba. One of 100 snails was infected in a rice culture field. This is the first time this species has been found acting as intermediate host of *F. hepatica* under natural conditions, not only for Cuba but also for the Caribbean area.

## Introduction

Fasciolosis is a parasitic disease that affects humans and other mammals, especially ruminants, representing a medical and veterinary problem with a serious impact on public health and economy at different levels, from local to national level in several countries (Crossland *et al.*, 1977; Rangel & Martinez, 1994; Roberts & Suhardono, 1996; Claxton *et al.*, 1997; Mas-Coma *et al.*, 1999; Ortiz *et al.*, 2000). The parasite responsible for this disease is the cosmopolitan digenean trematode *Fasciola hepatica*, with a worldwide distribution, especially in regions where cattle-raising has been developed. The life cycle of *F. hepatica* includes the transmission through a freshwater snail intermediate host belonging to the Lymnaeidae family (see review in Hurtrez-Boussès *et al.*, 2001). In Cuba, only two lymnaeid species occur, *Fossaria cubensis* and *Pseudosuccinea columella*, but until now the former

had been the only species found to be naturally infected in the field by the parasite (Melcon & Perera, 1994; Gutiérrez *et al.*, 2003a, 2005a, b). Although no report on *P. columella* harbouring parasite larvae under natural conditions had previously been made for the Caribbean area, this snail is also considered an effective host of *F. hepatica* as it acts as a natural intermediate host in countries such as Brazil, Australia and Argentina (Ueta, 1980; Boray *et al.*, 1985; Oliveira *et al.*, 2002; Prepelitchi *et al.*, 2003) and it shows a high susceptibility to miracidial exposures under laboratory conditions (Léon-Dancel, 1970; Boray *et al.*, 1985; Gutiérrez *et al.*, 2002).

In Cuba, both species occur at high densities in a variety of freshwater bodies, but *F. cubensis* shows a broader distribution throughout the island compared to *P. columella*, which is found from Pinar del Río through Camagüey provinces and seems to be absent from the eastern provinces Las Tunas through Guantánamo (Rojas *et al.*, 2010). Also, recent studies showed that, at least in Cuba, *F. cubensis* occupies the most anthropic habitats and *P. columella* is found more frequently in rural sites

\*Fax: + 33 468 662 281  
E-mail: theron@univ-perp.fr

(Cañete *et al.*, 2004). Recently, we detected very high densities of *P. columella* in a rice culture area at the El Pilón locality, Pinar del Río province, and searched to see if some of these snails were naturally infected with intramolluscan stages of the parasite, since a number of cows were normally seen grazing in the area and cases of bovine and caprine fasciolosis had been reported in the neighbouring village of San Andrés. In this paper we report the first infection observed in naturally occurring *P. columella* from Cuba.

## Materials and methods

### *Snail sampling and examination*

The locality of El Pilón is situated west of Bahía Honda village in the municipality of the same name (22°50'45"N; 83°21'32"W). This area belongs to the mountain system Sierra de los Organos in the Pinar del Río province. The sampling site consisted of a field sown with rice, completely flooded over an area of nearly 300 m<sup>2</sup>. The depth of water did not exceed 0.5 m, with a bottom mainly of mud and plant sediments. Suitability for the development of snail populations is granted for most of the year, with food from aquatic vegetation and water. As cattle-raising occurs in many farms in these localities, the life cycle of *F. hepatica* may be ensured.

Snail samples were taken on 17 December 2007 along the shores of the rice field. Snails were hand-collected from the mud or at the water surface using forceps. Because the abundance of their populations was extremely high, only 100 snails were randomly collected and saved in plastic boxes for laboratory studies. The presence of trematode infection was detected first by stimulation of cercarial shedding using a light source, and, second, by visual inspection of living snails for recognition of intramolluscan parasite development using a stereoscope. The search for infection in living snails lasted for a period of 15 days following snail collection. After this period the remaining snails were crushed and soft parts were analysed carefully, searching for parasite larvae. Each snail found harbouring trematode larvae was scored as infected. The hepatopancreas of each infected snail was dissected to remove redial stages of trematodes, which were stocked at 80% ethanol for later molecular analysis.

### *Molecular identification of F. hepatica intramolluscan stages*

DNA extractions of parasite larvae were performed using DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. In order to identify *F. hepatica* we amplified the rDNA gene, ITS-1 using the following primers, Lim1657 (forward): 5'-CTGCCCTTTGTACACACCG; and ITS1-RIXO: 5'-TGGCTGCGTTCATCG (Almeyda-Artigas *et al.*, 2000). For the amplification we used 2 µl of DNA template in a 25 µl reaction volume, containing: 5 µl each of polymerase chain reaction (PCR) buffer, 1.5 mM MgCl<sub>2</sub>, 200 µM of each dNTP, 10 pmol of each primer and 1 U GoTaq DNA polymerase (Promega, Madison, Wisconsin, USA). Temperature cycling was: 2 min at 94°C, followed by 30 cycles for 30 s at 94°C, 50°C for 30 s, 72°C for 30 s and finally 7 min at 72°C.

The amplified products (5 µl) were verified on 1% agarose gels in Tris–acetate–EDTA buffer. DNA sequencing was performed by CoGenics Genome Express (Meylan, France) using PCR-amplified products as templates.

The obtained sequences were aligned using CLUSTAL-W implemented in BioEdit v.7.0.9 (Hall & BioEdit, 1999). We used BLASTN 2.2.22 (available at <http://www.ncbi.nlm.nih.gov/BLAST>) to detect homologue sequences.

## Results

Three of the 100 snails sampled were found to be infected by redia of trematodes. The complete ITS-1 sequence obtained from trematode larvae found in *P. columella* from Cuba was 600 pb long. After BLAST analysis, our sequence was compared with some rDNA homologue sequences of *F. hepatica* available at <http://www.ncbi.nlm.nih.gov> (AB207139, AB207140, AB207141, AB385611, AJ243016). One of the three larval parasites showed no nucleotide difference with these five sequences. This similarity proves that the redia found in one *P. columella* snail from Cuba was *F. hepatica*. As reported previously, ITS-1 remains an appropriate marker for identification of this trematode species (Mera y Sierra *et al.*, 2009).

## Discussion

In spite of being very susceptible to parasite infection, as shown in previous experimental exposures (Gutiérrez *et al.*, 2002), so far no specimen of *P. columella* had been found harbouring intramolluscan larvae of *F. hepatica* in the field in Cuba. El Pilón is the first Cuban locality where the presence of *P. columella* naturally infected by *F. hepatica* has been demonstrated. All snails collected at the El Pilón site, where infection by the parasite was found, had mantles showing the type of pigmentation considered to be a good marker of susceptible populations of *P. columella* to *F. hepatica* infection (see Gutiérrez *et al.*, 2003b).

The Caribbean area is a region where *F. cubensis* seemed to be the unique intermediate host of *F. hepatica*. This situation is probably due to the fact that this snail is known as a local species, probably occurring in this area for a long time. It was apparently already distributed all over Cuba when it was first described by L. Pfeiffer at the beginning of the 19th century (Pointier *et al.*, 2005). On the contrary, *P. columella* originates from North America and is known as an invasive species which presently has a worldwide distribution (Pointier, 2008). In Cuba *P. columella* was reported for the first time by A. Poey in the middle of the 19th century, but at the present time the species only occurs from Pinar del Río to Camagüey provinces (Rojas *et al.*, 2010). In the Caribbean area *P. columella* was reported from Puerto Rico (Van der Schalie, 1948), Dominican Republic (Gómez *et al.*, 1986), Venezuela (Malek & Chrosiecchowski, 1964) and Guadeloupe (Pointier, 2008) but was never recorded as an active intermediate host of *F. hepatica* in the field. However, as already pointed out, in several other invaded countries, such as Australia, Brazil and Argentina, *P. columella* has been reported to be naturally infected by *F. hepatica* (Ueta, 1980; Boray *et al.*, 1985; Prepelitchi *et al.*, 2003). Consequently, the discovery of naturally infected

*P. columella* in Cuba shows that this species must be seriously taken into account in the development of future control programmes of this parasitosis.

## References

- Almeyda-Artigas, R.J., Bargues, M.D. & Mas-Coma, S. (2000) ITS-2 rDNA sequencing of *Gnathostoma* species (Nematoda) and elucidation of the species causing human gnathostomiasis in the Americas. *Journal of Parasitology* **86**, 537–544.
- Boray, J.C., Fraser, G.C. & Williams, J.M. (1985) The occurrence of the snail *Lymnaea columella* on grazing areas in New South Wales and studies on its susceptibility to *Fasciola hepatica*. *Australian Veterinary Journal* **62**, 4–6.
- Cañete, R.Y., Sánchez, J.M., Wong, L. & Gutiérrez, A. (2004) Population dynamics of intermediate snail hosts of *Fasciola hepatica* and some environmental factors in San Juan y Martínez municipality, Cuba. *Memorias do Instituto Oswaldo Cruz* **3**, 257–262.
- Claxton, J., Zambrano, H., Ortiz, P., Amoros, C., Delgado, E., Escurra, E. & Clarkson, M. (1997) The epidemiology of fascioliasis in the inter-Andean valley of Cajamarca, Peru. *Parasitology International* **46**, 281–288.
- Crossland, N., Johnstone, A., Beaumont, G. & Bennett, M. (1977) The effect of chronic fasciolosis on the productivity of lowland sheep. *British Veterinary Journal* **133**, 518–524.
- Gómez, J., Vargas, M. & Malek, E.A. (1986) *Pseudosuccinea columella* (Lymnaeidae) in the Dominican Republic and transmission of fascioliasis in the Caribbean region. *The Nautilus* **100**, 66–68.
- Gutiérrez, A., Yong, M., Perera, G., Sánchez, J. & Théron, A. (2002) *Fasciola hepatica* (Trematoda: Digenea): its effect on the life history traits of *Pseudosuccinea columella* (Gastropoda: Lymnaeidae), an uncommon interaction. *Parasitology Research* **88**, 535–539.
- Gutiérrez, A., Pointier, J.-P., Fraga, J., Jobet, E., Modat, S., Pérez, R.T., Yong, M., Sánchez, J., Loker, E.S. & Théron, A. (2003a) *Fasciola hepatica*: identification of molecular markers for resistant and susceptible *Pseudosuccinea columella* snail hosts. *Experimental Parasitology* **105**, 211–218.
- Gutiérrez, A., Pointier, J.-P., Yong, M., Sánchez, J. & Théron, A. (2003b) Evidence of phenotypic differences between resistant and susceptible isolates of *Pseudosuccinea columella* (Gastropoda: Lymnaeidae) to *Fasciola hepatica*. *Parasitology Research* **90**, 129–134.
- Gutiérrez, A., Hernandez, D. & Sánchez, J. (2005a) Variation of snail's abundance in two water bodies harboring strains of *Pseudosuccinea columella* resistant and susceptible to *Fasciola hepatica* miracidial infection in Pinar del Río province, Cuba. *Memorias do Instituto Oswaldo Cruz* **7**, 725–727.
- Gutiérrez, A., Yong, M., Sánchez, J., Wong, M. & Pointier, J.-P. (2005b) Competition between *Fossaria cubensis* and two isolates (susceptible and resistant to *Fasciola hepatica*) of *Pseudosuccinea columella* under laboratory conditions. *Haliotis* **35**, 1–11.
- Hall, T.A. & BioEdit (1999) A user friendly biological sequence alignment editor and analysis program for Windows 95/98NT. *Nucleic Acids Symposium* **41**, 95–98.
- Hurtrez-Boussès, S., Meunier, C., Durand, P. & Renaud, F. (2001) Dynamics of host–parasite interactions: the example of population biology of the liver fluke (*Fasciola hepatica*). *Microbes and Infection* **3**, 841–849.
- Léon-Dancel, D. (1970) Life history of *Lymnaea columella* (Say) and its experimental infection with *Fasciola hepatica* (L.). *Journal of Agriculture of the University of Puerto Rico* **2**, 297–305.
- Malek, E.A. & Chrosiecowski, P. (1964) *Lymnaea* (*Pseudosuccinea*) *columella* from Venezuela, and notes on the distribution of *Pseudosuccinea*. *The Nautilus* **78**, 54–56.
- Mas-Coma, S., Bargues, M.D. & Esteban, J.G. (1999) Human fasciolosis. pp. 411–434 in Dalton, J.P. (Ed.) *Fasciolosis*. Wallingford, UK, CAB International.
- Melcon, R. & Perera, G. (1994) Estudios ecológicos de *Fossaria cubensis* en sitios de transmisión de *Fasciola hepatica* en las Tunas y La Habana. *Parasitología al Día* **18**, 126–132.
- Mera y Sierra, R., Artigas, P., Cuervo, P., Deis, E., Sidoti, L., Mas-Coma, S. & Bargues, M.D. (2009) Fascioliasis transmission by *Lymnaea neotropica* confirmed by nuclear rDNA and mtDNA sequencing in Argentina. *Veterinary Parasitology* **166**, 73–79.
- Oliveira, S., Fujii, T., Filha, E. & Martins, A. (2002) Ocorrencia de *Lymnaea columella* Say, 1817 infectada naturalmente por *Fasciola hepatica* (Linnaeus, 1758), no vale do Ribeira, São Paulo, Brasil. *Aquívos do Instituto de Biologia, São Paulo* **1**, 29–37.
- Ortiz, P., Cabrera, M., Jav, J., Claxton, J. & Williams, D. (2000) Human fascioliasis: prevalence and treatment in a rural area of Peru. *Infectious Disease Review* **1**, 42–46.
- Pointier, J.-P. (2008) *Guide to the freshwater mollusks of the Lesser Antilles*. 128 pp. Hackenheim, Conchbooks.
- Pointier, J.P., Yong, M. & Gutiérrez, A. (2005) *Guide to the freshwater mollusks of Cuba*. 120 pp. Hackenheim, Conchbooks.
- Prepelitchi, L., Kleiman, F., Pietrokovsky, S.M., Moriena, R.A., Racioppi, O., Alvarez, J. & Wisnivesky-Colli, C. (2003) First report of *Lymnaea columella* Say, 1817 (Pulmonata: Lymnaeidae) naturally infected with *Fasciola hepatica* (Linnaeus, 1758) (Trematoda: Digenea) in Argentina. *Memorias do Instituto Oswaldo Cruz* **7**, 889–891.
- Rangel, J. & Martínez, E. (1994) Perdidas económicas por decomisos de hígados y distribución geográfica de la fasciolosis bovina en el Estado Tabasco. *Veterinaria de México* **25**, 337–341.
- Roberts, J. & Suhardon (1996) Approaches to the control of fasciolosis in ruminants. *International Journal for Parasitology* **26**, 971–981.
- Rojas, L., Vásquez, A.A., Domenech, I. & Robertson, L.J. (2010) Fascioliasis: can Cuba conquer this emerging parasitosis? *Trends in Parasitology* **26**, 26–34.
- Ueta, M. (1980) Estudo morfométrico da concha de *Lymnaea columella* Say, 1817 (Mollusca, Gastropoda, Pulmonata). *Revista da Sociedade Brasileira de Medicina Tropical* **1**, 119–141.
- Van der Schalie, H. (1948) *The land and fresh water mollusks of Puerto Rico*. 70 pp. Miscellaneous Papers and Publications of the Museum of Zoology, University of Michigan.