



HAL
open science

Diatom-specific highly branched isoprenoids as biomarkers in Antarctic consumers

Aurélie Goutte, Yves Cherel, Marie-Noëlle Houssais, Vincent Klein, Catherine Ozouf-Costaz, Mireille Raccurt, Camille Robineau, Guillaume Massé

► To cite this version:

Aurélie Goutte, Yves Cherel, Marie-Noëlle Houssais, Vincent Klein, Catherine Ozouf-Costaz, et al.. Diatom-specific highly branched isoprenoids as biomarkers in Antarctic consumers. PLoS ONE, 2013, 8 (2), pp.e56504. 10.1371/journal.pone.0056504 . halsde-00790218

HAL Id: halsde-00790218

<https://hal.science/halsde-00790218v1>

Submitted on 10 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Diatom-Specific Highly Branched Isoprenoids as Biomarkers in Antarctic Consumers

Aurélie Goutte^{1*}, Yves Chere², Marie-Noëlle Houssais¹, Vincent Klein¹, Catherine Ozouf-Costaz³, Mireille Raccurt⁴, Camille Robineau¹, Guillaume Massé^{1,5}

1 Laboratoire d'océanographie et du climat: expérimentations et approches numériques, UMR 7159 CNRS, Université Pierre et Marie Curie, Paris, France, **2** Centre d'Etudes Biologiques de Chizé, UPR 1934 CNRS, Beauvoir sur Niort, France, **3** Muséum National d'Histoire Naturelle, UMR 7138 CNRS, Paris, France, **4** Université de Lyon; UMR 5023 CNRS, Villeurbanne, France, **5** CNRS & Université Laval, UMI 3376, Takuvik, Québec, Canada

Abstract

The structure, functioning and dynamics of polar marine ecosystems are strongly influenced by the extent of sea ice. Ice algae and pelagic phytoplankton represent the primary sources of nutrition for higher trophic-level organisms in seasonally ice-covered areas, but their relative contributions to polar marine consumers remain largely unexplored. Here, we investigated the potential of diatom-specific lipid markers and highly branched isoprenoids (HBIs) for estimating the importance of these two carbon pools in an Antarctic pelagic ecosystem. Using GC-MS analysis, we studied HBI biomarkers in key marine species over three years in Adélie Land, Antarctica: euphausiids (ice krill *Euphausia crystallorophias* and Antarctic krill *E. superba*), fish (bald notothen *Pagothenia borchgrevinki* and Antarctic silverfish *Pleuogramma antarcticum*) and seabirds (Adélie penguins *Pygoscelis adeliae*, snow petrels *Pagodroma nivea* and cape petrels *Daption capense*). This study provides the first evidence of the incorporation of HBI lipids in Antarctic pelagic consumers. Specifically, a di-unsaturated HBI (diene) of sea ice origin was more abundant in ice-associated species than in pelagic species, whereas a tri-unsaturated HBI (triene) of phytoplanktonic origin was more abundant in pelagic species than in ice-associated species. Moreover, the relative abundances of diene and triene in seabird tissues and eggs were higher during a year of good sea ice conditions than in a year of poor ice conditions. In turn, the higher contribution of ice algal derived organic matter to the diet of seabirds was related to earlier breeding and higher breeding success. HBI biomarkers are a promising tool for estimating the contribution of organic matter derived from ice algae in pelagic consumers from Antarctica.

Citation: Goutte A, Chere Y, Houssais M-N, Klein V, Ozouf-Costaz C, et al. (2013) Diatom-Specific Highly Branched Isoprenoids as Biomarkers in Antarctic Consumers. PLoS ONE 8(2): e56504. doi:10.1371/journal.pone.0056504

Editor: André Chiaradia, Phillip Island Nature Parks, Australia

Received: July 5, 2012; **Accepted:** January 15, 2013; **Published:** February 13, 2013

Copyright: © 2013 Goutte et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by the European Research Council (ICEPROXY, project number 203441) and the French Institut Paul-Emile Victor (IPEV projects N° 1010, G. Massé; N° 1124, G. Lecoindre; N° 131, C. Duchamp; N° 1039, J.-Y. Touleuc and N° 109, H. Weimerskirch). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: agoutte@gmail.com

Introduction

Shrinking sea ice threatens the structure, functioning and dynamics of polar marine ecosystems [1,2,3]. Many species rely on sea ice to complete their life history. At the base of the polar marine trophic web, some micro-algae, primarily pennate diatoms, bloom under and within sea ice in the spring [4,5,6]. A second source of primary production is pelagic phytoplankton, which bloom during the summer ice melts. Sea ice decay and breakup modifies the physical and chemical parameters of oceans, including water column stability, nutrient availability, salinity and solar radiation [7,8]. Several attempts have been made to estimate the relative contribution of ice algae and phytoplankton to primary production in Arctic and Antarctic waters [5,9]. On an annual basis, ice-related primary production contributes approximately 5 to 28% of the total production in the ice-covered waters of the Southern Ocean [5,6]. Although ice algae are a significant (direct or indirect) source of nutrition for zooplankton, krill, fish, seabirds and marine mammals [6], their relative contribution to polar marine ecosystems remains poorly understood. Recently, a method to trace carbon flow through Arctic marine trophic webs by analysing the stable isotopes of fatty acids was proposed [10].

Although elegant, the method presents some limitations because of the isotopic fractionation of some fatty acids during ingestion [11].

The aim of the present study was to evaluate the potential of some novel lipid markers, highly branched isoprenoids (HBIs), for estimating the contribution of organic matter derived from ice algae in pelagic consumers from Antarctica. C₂₅ HBIs are synthesised by a few diatom species [12]. In Antarctica, a di-unsaturated HBI isomer (diene, Figure 1 A) was identified in lipid fractions from diatom communities in sea ice, whereas other HBIs, such as a tri-unsaturated isomer (triene, Figure 1 B), were absent in sea ice samples but detected in ice edge and open ocean phytoplankton communities [13,14]. Recent investigations into the carbon isotopic composition of diene isolated from Antarctic sea ice and sediments revealed that this isomer was isotopically ¹³C enriched relative to the more unsaturated HBIs [13]. This enrichment is consistent with a sea ice origin and is likely derived from *Haslea* and/or *Navicula* species of diatoms, which live in the brine channels at the base of Antarctic sea ice and bloom in spring [12,13,14]. In contrast, the lower δ¹³C values of triene and other poly-unsaturated isomers are consistent with a phytoplanktonic origin [13]. Planktonic diatom species that bloom in open waters

near the marginal ice zone, such as *Rhizosolenia* spp., are likely responsible for the synthesis of triene and other poly-unsaturated HBI isomers [12,13,14]. Based on their occurrence in sediments and their specific origin, these and some other related HBIs markers have been used as proxies for sea ice reconstruction in the Arctic [15,16,17] and Antarctic [13,18]. In particular, the relative abundance of diene to triene (hereafter, the D/T ratio) has been successfully used to estimate the relative contributions of sea-ice-derived and planktonic diatoms to the total primary production in Antarctica [13]. Therefore, HBIs may prove useful for ecological studies at higher trophic levels as ice-based *versus* phytoplankton biomarkers in polar marine ecosystems. Three recent studies have detected HBI isomers in pelagic zooplankton and benthic macrofauna in the Arctic [19,20,21], demonstrating that these diatom-specific biomarkers are transferred to Arctic consumers and establishing their potential for determining the relative contribution of ice algae and pelagic phytoplankton to higher trophic levels.

Here, we investigated the potential of this approach for studying the Antarctic pelagic ecosystem. Our first aim was to determine whether HBIs are transferred to key pelagic species in Antarctica. We collected specimens of two krill species, two planktivorous fish species and three seabird species and analysed the presence and concentrations of HBIs. Our second aim was to identify whether the sea-ice-specific diene was more abundant in species preferentially feeding in ice-covered waters (cryopelagic species) than in pelagic species by comparing corresponding tissues among closely related taxa. Finally, we tested (i) whether Antarctic seabirds incorporate more diene during a year of extended pack ice than during a year of reduced ice and (ii) whether a high contribution of ice algal-derived organic matter predict key ecological parameters; i.e., earlier breeding and higher reproductive success.

Materials and Methods

1. Ethics statement

Ethical approval for all procedures was granted by the ethics committee of the Ministère de l'Environnement and the French Polar Research Institute (Institut Paul Emile Victor – IPEV),




Figure 1. Structure of the two HBIs biomarkers: (A) diene and (B) triene.

doi:10.1371/journal.pone.0056504.g001

which approved all our fieldwork. The experiments complied with the Code of Ethics of Animal Experimentation in the Antarctic.

2. Study site and specimen collection

Table 1 summarises information on specimen collection. Briefly, seabirds (see below) and the bald notothen *Pagothenia borchgrevinki* were studied in the Pointe Géologie Archipelago (Adélie Land, Antarctica, 66°40'S, 140°01'E, Figure 2 A), whereas the ice and Antarctic krill (*Euphausia crystallorophias* and *E. superba*, respectively) and the Antarctic silverfish *Pleuragramma antarcticum* were collected from the Research Icebreaker Astrolabe using an Isaacs-Kidd Midwater Trawl (IKMT). The Adélie Land area is ice covered seasonally, with land-fast ice typically forming in March/April and breaking up in November/December. The cruise was conducted in January 2011 in open waters (between 65°31'S/66°33'S and 139°59'E/143°60'E, Figure 2 B) during the Austral summer.

For the two euphausiids, oceanic Antarctic krill were collected at two locations over the continental slope (site AS605: 65°31'S, 140°00'E and site AS608: 65°39'S, 142°56'E, Figure 2 B), whereas neritic ice krill [22] were collected at two coastal stations (site AS601: 66°19'S, 139°59'E and site AS607: 66°33'S, 143°40'E, Figure 2 B). Krill exploit both sea ice and open water habitats throughout the year, but larval and postlarval krill are strongly associated with the underneath of the sea ice [23,24]. The Antarctic silverfish was collected at the site AS612 (66°18'S, 141°56'E, Figure 2 B) and is the predominant mid-water notothenioid fish in neritic Antarctic waters. During adulthood, it exploits open water habitats; however, juvenile developmental stages and spawning occur under and within sea ice [25,26]. Bald notothen *Pagothenia borchgrevinki* were sampled under the sea ice closed to Bernard Island (Figure 2 A) using a fishing rod in December 2010. This species is a common cryopelagic fish that forages underneath of the sea ice [26,27]. For the two fish species, muscle samples were excised.

Pointe Géologie Archipelago is a breeding area for approximately 36,000 pairs of Adélie penguins (*Pygoscelis adeliae*), 870 pairs of snow petrels (*Pagodroma nivea*) and 450 pairs of cape petrels (*Daption capense*) [28]. Seabirds were sampled at Petrels Island, the main island of the Pointe Géologie Archipelago, where the French Antarctic station, Dumont D'Urville is located (Figure 2 A). The snow petrel is confined to high Antarctic waters feeding primarily on fish among ice floes. The cape petrel has a wider latitudinal distribution and feeds primarily on crustaceans in Adélie Land [29,30,31]. All procellariids store stomach oil from the breakdown of ingested food and the differential digestion rates of proteins and lipids. We collected both abandoned petrel eggs and the stomach oil of chicks during the 2010/2011 breeding season (Table 1) for HBI analysis. The Adélie penguin is an ice-associated diver that feeds in areas of 20 to 80% ice, but it can also forage in the open sea, under pack ice and under coastal fast ice [32,33]. Abandoned eggs and dead chicks of Adélie penguins were collected (Table 1) and blood samples taken. The stomach; liver; kidneys; quadriceps, pectoralis and gastrocnemius muscles; and subcutaneous and retroperitoneal adipose tissues were sampled when possible. The sampling dates of chicks (from the 28th of December to the 21st of January) did not differ between 2009/2010 and 2010/2011 (paired t-test, $t = -1.874$, $df = 8$, $p = 0.098$). All samples were immediately frozen at -80°C until analysis at the LOCEAN/IPSL (Paris, France) laboratory.

3. HBI analyses

HBI analyses were conducted according to a previously described procedure [15]. Each biological tissue was freeze-dried and an internal standard (7-hexylnonadecane) was added to an




Figure 2. Map showing the location of the Pointe Géologie archipelago and all the sampling sites off Adélie Land, Antarctica. (A) Samples of Adélie penguin, snow petrel and cape petrel were collected on Petrel Island; bald notothen were fished under the sea closed to Bernard Island, Pointe Géologie archipelago. (B) Ice krill, Antarctic krill and Antarctic silverfish were collected off Adélie Land (see the method section for GPS locations). The high-resolution bathymetry of the areas is represented [48]. doi:10.1371/journal.pone.0056504.g002

aliquot of freeze-dried sample (approximately 1 g dry weight). The sample was saponified using a solution of KOH (4N, MeOH/H₂O, 80/20) for 2 hours at 80°C, and the non-saponifiable lipids (NSL) were extracted into hexane (3×10 mL). The extract was purified using open column chromatography (SiO₂ 50 g.g⁻¹ NSL; hexane 8 mL) to yield an apolar lipid fraction containing the HBIs. This hydrocarbon fraction was analysed using an Agilent 7890A gas chromatograph (GC) fitted with a 30 m fused silica Agilent J&C GC column (0.25 mm i.d., 0.25 μm film) coupled to a Agilent 5975C Series mass selective detector (MSD). A ramped oven temperature profile (40–300°C; 10°C/min) followed by an isothermal interval (300°C for 10 min) was used. Data were collected and analysed with Agilent Chemstation software. Individual compounds were identified and quantified using both scan (m/z 50–500 mass range) and selective ion monitoring (SIM)

techniques with a constant ionisation potential (70 eV). Mass spectral data were used to identify the occurrence of HBIs, whereas SIM chromatograms were used to quantify the abundances of diene (m/z 348.3), triene (m/z 346.3), tetraene (m/z 344.3) and pentaene (m/z 342.3) by peak area integration. Individual HBI isomers were identified by comparing their GC retention indices and mass spectra with those of previously authenticated standards. HBI relative abundances were calculated based on their individual GC-MS responses and those of the internal standard (7-hexylnonadecane, m/z 266). As we did not determine their individual response factors, HBI abundances were expressed in ng.g⁻¹ relative to the internal standard. Procedural blanks were analysed for each 20 samples to ensure the absence of contamination.

Table 1. Specimen information: common and scientific names of species, stage, tissue, sampling date and site.

Species	Stage	Tissue type	N	Collection date	Sampling site
KRILL					
Antarctic krill <i>Euphausia superba</i>	Adult	Whole body	27*	January 2011	Continental slope
Ice krill <i>Euphausia crystallographica</i>	Adult	Whole body	38*	January 2011	Adélie Land coast
FISH					
Antarctic silverfish, <i>Pleuragramma antarcticum</i>	Adult	Muscle	6	January 2011	Continental slope
Bald notothen, <i>Pagothenia borchgrevinki</i>	Adult	Muscle	9	December 2010	Pointe Géologie Archipelago
SEABIRD					
Adelie penguin, <i>Pygoscelis adeliae</i>	Chick	T1	12	Dec 2009–Jan 2010	Petrels island
Adelie penguin, <i>Pygoscelis adeliae</i>	Chick	T2	12	Dec 2010–Jan 2011	Petrels island
Adelie penguin, <i>Pygoscelis adeliae</i>	Egg	Whole egg	4	December 2011	Petrels island
Snow petrel, <i>Pagodroma nivea</i>	Egg	Whole egg	5	December 2010	Petrels island
Snow petrel, <i>Pagodroma nivea</i>	Egg	Whole egg	14	December 2007	Petrels island
Snow petrel, <i>Pagodroma nivea</i>	Chick	Stomach oil	13	January 2011	Petrels island
Cape petrel, <i>Daption capense</i>	Chick	Stomach oil	4	January 2011	Petrels island

*Pool of individuals.

T1: Subcutaneous adipose tissue, retroperitoneal adipose tissue, gastrocnemius muscle, pectoralis muscle, quadriceps muscle, liver, stomach content.

T2: Same tissues than T1, blood and kidney.

doi:10.1371/journal.pone.0056504.t001

4. Prey identification in Adélie penguin food samples

In the laboratory, chick stomach contents were placed in a large flat-bottomed tray. Sixteen stomachs contained prey items that were not completely digested. The most abundant prey items were identified to species in three to six sub-samples of each food sample. Prey were identified through examination of exoskeletons (for crustaceans) and of otoliths and bones (for fish). Preys items were counted in each sample. We estimated the proportions of fresh mass represented by each species.

5. Sea ice data

Daily remote-sensing sea ice concentration maps, retrieved from the Advanced Microwave Scanning Radiometer–Earth Observing System (AMSR-E), were obtained from the University of Hamburg, Institute of Oceanography from the web site ftp://ftp-projects.zmaw.de/seaice/AMSR-E_ASI_IceConc/no_landmask/hdf/s6250/ [34]. Sea ice concentration data were calculated with the ARTIST sea ice (ASI) concentration algorithm using AMSR-E 89 GHz brightness temperatures. We extracted daily sea-ice concentrations in the area within 137.5°E–142.5°E and 65°S–67°S (approximately 100 km from the Dumont D'Urville station) from September 2007 to March 2011.

6. Ecological data

Phenological and reproductive success data were obtained for seabirds via long term monitoring programs conducted by the Centre d'Etudes Biologiques de Chizé [35,36]. The peak dates of egg laying and hatching, as well as the total numbers of Adélie penguin breeding pairs, chicks in crèche and fledglings were monitored on Petrels Island. For snow petrels, the number of eggs, hatching date and number of fledglings were monitored daily for 282 nests in three colonies located on Petrels Island.

7. Statistical analyses

All statistical analyses were performed using R 2.14.2 [37]. It was recently proposed that the D/T ratio reflects the relative contribution of sea ice and planktonic algae to sediments [13,17]. However, HBIs may degrade or be metabolised while incorporated within tissue. Because of its additional double bond, the triene is likely to degrade much quicker than the diene, we first analysed diene and triene separately, then analysed D/T ratios. Significantly higher D/T ratios were all validated by either a significantly higher diene value and/or a significantly lower triene value. We therefore decided to present the D/T ratio results as a proxy of the relative contribution of sea ice and planktonic derived organic matter to consumer diets. We used Wilcoxon rank sum tests and pairwise Wilcoxon tests with post-hoc Bonferroni corrections to test for differences in D/T ratios between closely related species and years. We used paired t-tests to test for differences between years in sea ice concentrations.

Results

1. Are HBI diatom biomarkers transferred across the Antarctic trophic web?

Diene and triene were detected in all Antarctic pelagic organisms, including krill, fish and seabirds (Table 2). The abundance of diene varied from 0.03 to 16.71 ng.g⁻¹ and the abundance of triene varied from 0.53 to 473 ng.g⁻¹ both relative to the internal standard. It is important to notice that HBI analyses were conducted either on whole organisms or biological tissues on dissected organisms. The relative abundances of diene and triene were the highest in krill and in stomach contents/oils, livers, eggs

of seabirds, whereas diene and triene were in very low abundance in muscles of Adélie penguins and fish, kidneys, adipose tissues, and blood of Adélie penguins. Moreover, concentrations of diene varied greatly between tissue type in Adélie penguin chicks (Kruskal-Wallis $\chi^2 = 67.147$, $df = 8$, $p < 0.001$). The highest concentrations of HBIs in Adélie penguin were detected in the stomach content, followed by eggs and liver. The lowest concentrations were found in the quadriceps, pectoralis and gastrocnemius muscles, kidneys, subcutaneous and retroperitoneal white adipose tissues and blood.

2. Do D/T ratios vary among Antarctic consumers?

D/T ratios did not differ between Antarctic and ice krill but differed significantly between sampling sites but eastern sites (i.e. AS607 and AS608) having significantly higher D/T ratios than western sites (AS601 and AS605, pairwise Wilcoxon tests with post-hoc Bonferroni corrections, $p < 0.01$, Figure 3 A). In contrast, no significant difference was observed between neritic (AS601 and AS607) and pelagic (AS605 and AS608) sampling sites (pairwise Wilcoxon tests with post-hoc Bonferroni corrections, $p < 0.210$).

D/T ratios were significantly higher in muscles of bald notothen from under the ice than in muscles of Antarctic silverfish collected in open waters (Wilcoxon rank sum test: $N = 15$, $W = 51$, $p = 0.003$, Figure 3 B).

D/T ratios in chick stomach oil were significantly higher in snow petrels than in cape petrels (Wilcoxon rank sum test: $W = 0$, $p = 0.004$, Figure 3 C). In eggs, D/T ratios were significantly higher in Adélie penguins than in snow petrels (Wilcoxon rank sum test: $N = 9$, $W = 20$, $p = 0.016$, Figure 3 D).

3. Do D/T ratios in Antarctic consumers vary with sea ice conditions?

During the Adélie penguin breeding season (defined as the period from the onset of egg-laying, around the 1st of November until the start of chicks' fledgling, the 31st of January), sea ice concentrations were significantly higher in 2009/2010 (mean \pm SE during these 3 months: 55.8% \pm 2.1%) than in 2010/2011 (46.3% \pm 1.9%, paired t-test, $t = 9.086$, $df = 91$, $p < 0.001$, Figure 4 A). Adélie penguin dead chicks were collected from the 28th of December (just after hatching) to the 21st of January (30-day old). During this sampling period, sea ice concentrations were also significantly higher in 2009/2010 (37.2% \pm 1.2%) than in 2010/2011 (31.5% \pm 1.2%, paired t-test, $t = 5.810$, $df = 24$, $p < 0.001$).

D/T ratios in all analysed tissues (liver, quadriceps, pectoralis and gastrocnemius muscles, subcutaneous and retroperitoneal white adipose tissue) and in stomach contents of Adélie penguin chicks were significantly higher during the 2009/2010 breeding season than during the 2010/2011 breeding season ($p < 0.001$ for all Wilcoxon tests; Figure 4 B).

Prey identified in stomach contents revealed less consumption of ice krill (Wilcoxon test, $W = 6$, $p = 0.015$) and more consumption of Antarctic krill (Wilcoxon test, $W = 44.5$, $p = 0.044$) in 2009/2010 than in 2010/2011 (Figure 4 C). The mean biomass of Antarctic silverfish in stomach contents did not differ significantly between years (Wilcoxon test, $W = 13$, $p = 0.104$, Figure 4 C).

The long term monitoring programme conducted on Petrels Island, where Adélie penguin dead chicks were collected, revealed that the total number of Adélie penguin pairs that bred did not differ significantly between years (15999 breeding pairs in 2009/2010 and 14694 in 2010/2011). However, both the number of chicks in crèche ($N = 15624$ and $N = 9561$, respectively) and the number of fledglings ($N = 15405$ and $N = 9256$, respectively) were higher in 2009/2010 than in 2010/2011. Egg-laying date (peak: 17/11/2009 and 22–23/11/2010) and hatching date (peak: 21/

Table 2. Abundance of diene, triene (mean \pm SE, in $\text{ng}\cdot\text{g}^{-1}$ relative to the internal standard) and the diene/triene ratio in Antarctic pelagic species (whole organism, tissues or eggs) collected during the austral summers 2007/2008, 2009/2010 and 2010/2011.

Species	Diene			Triene			Diene/Triene		
	N	2010/2011		N	2010/2011		N	2010/2011	
Krill									
Ice krill (whole body)	38	11.10 \pm 2.00		38	273.05 \pm 20.88		38	0.03 \pm 0.003	
Antarctic krill (whole body)	27	11.49 \pm 1.17		27	266.13 \pm 22.94		27	0.04 \pm 0.003	
Fish									
Bald notothen (muscle)	9	0.30 \pm 0.10		9	0.53 \pm 0.04		9	0.58 \pm 0.19	
Antarctic silverfish (muscle)	6	0.97 \pm 0.36		6	20.35 \pm 8.88		6	0.06 \pm 0.01	
Adelie penguins									
Egg	4	7.04 \pm 1.41		4	269.60 \pm 42.78		4	0.03 \pm 0.004	
Stomach content	12	7.99 \pm 10.23		12	109.71 \pm 80.76		12	0.24 \pm 0.07	
Subcutaneous adipose tissue	10	1.42 \pm 0.92		10	1.56 \pm 0.39		10	0.3 \pm 0.08	
Retroperitoneal adipose tissue	6	1.47 \pm 1.35		6	2.47 \pm 1.49		6	0.27 \pm 0.13	
Liver	10	14.96 \pm 7.86		10	64.44 \pm 28.11		10	0.17 \pm 0.05	
Gastrocnemius muscle	5	0.33 \pm 0.19		5	1.87 \pm 0.75		5	0.12 \pm 0.05	
Pectoralis	5	0.47 \pm 0.12		5	2.43 \pm 1.13		5	0.28 \pm 0.1	
Quadriceps	8	0.71 \pm 0.29		8	1.74 \pm 0.62		8	0.32 \pm 0.1	
Kidney	11	0.22 \pm 0.07		11	5.73 \pm 0.94		11	0.06 \pm 0.03	
Blood	10	0.45 \pm 0.09		10	17.11 \pm 2.95		10	0.03 \pm 0.01	
Snow petrels									
Egg	14	0.80 \pm 0.16		14	20.16 \pm 3.81		14	0.04 \pm 0.01	
Stomach oil	13	8.67 \pm 0.41		13	77.69 \pm 8.64		13	0.12 \pm 0.01	
Cape petrels									
Stomach oil	4	1.32 \pm 0.51		4	80.62 \pm 23.17		4	0.02 \pm 0.002	

doi:10.1371/journal.pone.0056504.t002




Figure 3. Diene and triene in key marine species from Adélie Land, Antarctica. The D/T ratio (mean \pm SE) were compared between (A) two euphausiid species (ice krill and Antarctic krill) at four sampling sites, (B) two fish species (bald notothen and Antarctic silverfish), (C) eggs of two seabird species (snow petrel and Adélie penguin) and (D) chick stomach oil of two seabird species (snow petrel and cape petrel). doi:10.1371/journal.pone.0056504.g003




Figure 4. Differences in ice concentrations, HBIs biomarker concentrations and prey characteristics of Adélie penguin chicks between 2009/2010 and 2010/2011. (A) Daily difference in sea ice concentration from November to January between the 2009/2010 and 2010/2011 breeding season, (B) D/T ratios (mean \pm SE) measured in tissues and stomach contents of Adélie penguin chicks, (C) prey composition and abundance in chick stomach contents. doi:10.1371/journal.pone.0056504.g004

12/2009 and 23–24/12/2010) were approximately 2.5 days earlier in 2009/2010 than in 2010/2011.

We investigated HBI concentrations in abandoned snow petrel eggs between 2007 and 2010. Female snow petrels forage at sea from mid-November to mid-December while forming their single eggs. Analysis of satellite data revealed that sea ice concentrations were significantly higher in 2007 than in 2010 during this period ($t = 5.647$, $df = 28$, $p < 0.001$, Table 3). D/T ratios of eggs were also significantly higher in 2007 than in 2010 (Wilcoxon rank sum test, $N = 19$, $W = 70$, $p < 0.001$, Table 3). Among the 282 monitored nests, both the number of breeding pairs and the number of fledglings were higher in 2007/2008 than in 2010/2011; however, the mean hatching date was 2 days later in 2007/2008 than in 2010/2011 (Table 3).

Table 3. Variation in sea ice concentration, D/T ratios in snow petrel eggs and breeding parameters (number of breeding pairs, number of fledglings and hatching date) of snow petrels on Ile des Pétreles, Adélie Land.

Breeding Season	2007/2008	2010/2011
Sea ice concentration (%)	65.3 \pm 2.2	54.1 \pm 2.6
D/T ratio in eggs	0.04 \pm 0.01	0.01 \pm 0.0005
Number of breeding pairs	204	193
Number of fledglings	139	100
Hatching date (January)	21.42 \pm 0.25	19.42 \pm 0.20

doi:10.1371/journal.pone.0056504.t003

Discussion

1. Are HBI diatom biomarkers transferred across the Antarctic trophic web?

HBI markers have recently been identified in Arctic benthic macrofauna and pelagic zooplankton [19,20,21], but their applicability to Antarctic marine species has been largely unexplored. The present study provides the first evidence that diatom HBI biomarkers are transferred across an Antarctic pelagic ecosystem up to higher trophic levels. Specifically, the occurrence of the sea ice biomarker diene may provide a new way of assessing the contribution of ice algal-derived organic matter in the diet of consumers, given this HBI lipid was detected in all organisms investigated in this study.

In Adélie penguin chicks, we observed large differences in the concentrations of HBIs between tissue types, suggesting a time-integrated assimilation of HBI lipids within organisms as for fatty acids [e.g. 38]. Our results suggest that the liver preferentially accumulates HBIs relative to adipose tissue, muscle and kidneys which might be useful for assessing the importance of sea ice primary production to vertebrate species over long periods. D/T ratios in stomach contents may permit estimation of the consumption of cryopelagic versus pelagic prey over the preceding days. The presence of HBIs in eggs indicates that these compounds are transferred from females to their eggs. However, little is known regarding the mechanisms of HBI incorporation into tissues, including assimilation, metabolic and elimination rates.

2. Do D/T ratios vary among Antarctic consumers?

Krill use both ice and pelagic food sources throughout the year [39,40]. As ice krill prefer more neritic regions relative to Antarctic krill [22], we expected higher abundances of diene, lower abundances of triene, or both in ice krill. Unexpectedly, the two species did not differ in their D/T ratios, suggesting a similar consumption of cryopelagic and pelagic diatoms in Adélie Land during the summer. However diene and triene abundances differed greatly between sampling sites, with eastern sites having significantly higher D/T ratios than western sites. These results suggest that the incorporation of HBIs in krill swarms may be related to the presence of drifting ice flows. In summer, high densities of Antarctic krill in the 0–2 m surface layer have been documented in the under-ice habitat but not in the open surface layer [24]. In Adélie Land, fast ice broke up in spring, but drift ice was still present during krill sampling. An alternative, non-mutually exclusive hypothesis is that krill swarms vary in the relative proportion of sub-adults to adults between sampling sites [41]. As younger stages are more strongly associated with sea ice,

D/T ratios may increase with increasing occurrence of juveniles and sub-adults in krill swarms.

D/T ratios were higher in muscles of bald notothen collected under the ice than in muscles of Antarctic silverfish collected in open waters. The bald notothen feeds primarily on cryopelagic copepods, pteropods, gammarids and hyperiids in platelet ice and water beneath the ice [26,27], whereas the Antarctic silverfish feeds on euphausiids, pelagic copepods, gastropods and gammarids [25]. Therefore, the D/T ratio in muscle appears to be a good proxy of the contribution of ice algal-derived organic matter in the diet of fish species. We also detected diene in adult Antarctic silverfish. This result may reflect the consumption of partially ice-associated prey or the persistence of diene from juvenile stages, which are associated with sea ice and actively feed on sympagic zooplankton [26,42]. Therefore, even at low levels, the occurrence of diene indicates an interaction with sea ice.

At the top of the trophic web, D/T ratios within chick stomach oil were higher in snow petrels, which are piscivorous seabirds confined to the pack ice zone [29,30,31,32], than in cape petrels, which are krill consumers exploiting open inshore waters [29]. Thus, our findings are consistent with a higher consumption of ice-associated prey in snow petrels than in cape petrels. Moreover, D/T ratios were higher in the eggs of Adélie penguins than in those of snow petrels, suggesting that female Adélie penguins rely more heavily on ice-associated prey during egg formation than snow petrels. As ice has yet to break up during the period of energy accumulation for egg building, it is possible that cryopelagic prey are more readily accessible to divers, such as penguins, than to aerial foragers, such as petrels.

3. Are D/T ratios in Antarctic consumers related to sea ice conditions?

To control for among-tissue differences, we performed comparisons on the same tissue types of Adélie penguins during the two consecutive breeding seasons. D/T ratios in all tissues and stomach contents of Adélie penguin chicks were lower in 2010/2011 than in 2009/2010, coinciding with 10% lower daily sea ice concentrations during the 2010/2011 breeding season than in the 2009/2010 breeding season. Although little is known about HBI production and D/T ratio changes through time in Antarctica, these results suggest a reduced contribution of ice algal-derived

organic matter in the diet of chick-rearing Adélie penguins in 2010/2011. Furthermore, prey items in chick stomachs confirmed a change in diet between years. Although ice krill was the predominant prey (as reported elsewhere; e.g., [43]), Antarctic krill were more abundant during 2009/2010, the period of greater sea ice concentration. Extensive sea ice induces high overall primary production and the dominance of 'large' diatoms over 'small' cryptophytes [44]. As Antarctic krill are more efficient at grazing large particles [23], the survival and recruitment success of Antarctic krill are enhanced during years of heavier ice conditions [45,46]. In turn, Adélie penguins rely more heavily on Antarctic krill, breed earlier and have higher breeding success, as reported here and in previous studies [42,47]. Similarly, lower D/T ratios were reported in snow petrel eggs during a year of reduced sea ice concentration and were accompanied by high breeding failure.

Annual variation in the phenology and ecology of Antarctic top predators with variation in the extent of sea ice has been widely discussed [47]. However, the importance of ice-related primary production has been little studied due to the lack of appropriate tools. HBI lipid biomarkers are a promising tool for tracking carbon flow in ice-covered oceans. As many migratory species feed on the highly productive, ice-covered waters of the Southern Ocean [6], the application of HBI lipid measures to the study of sea ice primary production may be extended to larger spatial scales. Ultimately, they may be useful for understanding the consequences of shrinking sea ice on the structure, functioning and dynamics of polar marine trophic webs.

Acknowledgments

We thank C. Blanc, J. Chevallier, J. Etourneau, C. Gallut, S. de Grissac, and J.-Y. Toullec, for their help in collecting samples in the field. We also thank A. Lorre and F. Cargouet for their excellent technical assistance in lipid purification. The manuscript benefited greatly from the comments of Thomas Brown.

Author Contributions

Conceived and designed the experiments: AG YC COC MR GM. Performed the experiments: AG VK CR GM. Analyzed the data: AG. Contributed reagents/materials/analysis tools: MNH GM. Wrote the paper: AG YC GM.

References

- Clarke A, Harris CM (2003) Polar marine ecosystems: major threats and future change. *Environ Conserv* 30: 1–25.
- Smetacek V, Nicol S (2005) Polar ocean ecosystems in a changing world. *Nature* 437: 362–368.
- Schofield O, Ducklow HW, Martinson DG, Meredith MP, Moline MA, et al. (2010) How do polar marine ecosystems respond to rapid climate change? *Science* 328: 1520–1523.
- Garrison DL (1991) Antarctic sea ice biota. *Am Zool* 31: 17–33.
- Lizotte MP (2001) The contributions of sea ice algae to Antarctic marine primary production. *Amer Zool* 41: 57–73.
- Arrigo KR, Thomas D (2004) Large scale importance of sea ice biology in the Southern Ocean. *Antarct Sci* 16: 471–486.
- Arrigo KR, Lizotte MP, Worthen DL, Dixon P, Dieckmann G (1997) Primary production in Antarctic sea ice. *Science* 276: 394–397.
- Eicken H (1992) The role of sea ice in structuring Antarctic ecosystems. *Polar Biol* 12: 3–13.
- Gosselin M, Levasseur M, Wheeler PA, Horner RA, Booth BC (1997) New measurements of phytoplankton and ice algal production in the Arctic Ocean. *Deep-Sea Res II* 44: 1623–1644.
- Budge SM, Wooller MJ, Springer AM, Iverson SJ, McRoy CP, et al. (2008) Tracing carbon flow in an Arctic marine food web using fatty acid-stable isotope analysis. *Oecologia* 157: 117–129.
- Bec A, Perga ME, Koussoroplis AM, Bardoux G, Desvilettes C, et al. (2011) Assessing the reliability of fatty acids-specific stable isotope analysis for trophic studies. *Methods Ecol Evol* 2: 651–659.
- Sinninghe Damsté JS, Muyzer G, Abbas B, Rampen SW, Massé G, et al. (2004) The rise of the rhizosolenid diatoms. *Science* 304: 584–587.
- Massé G, Belt ST, Crosta X, Schmidt S, Snape I, et al. (2011) Highly branched isoprenoids as proxies for variable sea ice conditions in the Southern Ocean. *Antarct Sci* 23: 487–498.
- Johns L, Wraige EJ, Belt ST, Lewis CA, Massé G, et al. (1999) Identification of a C25 highly branched isoprenoid (HBI) diene in Antarctic sediments, Antarctic sea-ice diatoms and cultured diatoms. *Org Geochem* 30: 1471–1475.
- Belt ST, Massé G, Rowland SJ, Poulin M, Michel C, et al. (2007) A novel chemical fossil of palaeo sea ice: IP₂₅. *Org Geochem* 38: 16–27.
- Massé G, Rowland SJ, Sicre MA, Jacob J, Jansen E, et al. (2008) Abrupt climate changes for Iceland during the last millennium: Evidence from high resolution sea ice reconstructions. *Earth Planet Sci Lett* 269: 564–568.
- Müller J, Massé G, Stein R, Belt ST (2009) Variability of sea ice conditions in the Fram Strait over the past 30,000 years. *Nat Geosci* 2: 772–776.
- Barbara L, Crosta X, Massé G, Ther O (2010) Deglacial environments in eastern Prydz Bay, East Antarctica. *Quat Sci Rev* 29: 2731–2740.
- Brown TA, Belt ST (2012) Identification of the sea ice diatom biomarker IP₂₅ in Arctic benthic macrofauna: direct evidence for a sea ice diatom diet in Arctic heterotrophs. *Polar Biol* 35: 131–137.
- Brown TA, Belt ST (2012) Closely linked sea ice–pelagic coupling in the Amundsen Gulf revealed by the sea ice diatom biomarker IP₂₅. *J Plankton Res* 34: 647–654.
- Brown TA, Belt ST, Piepenburg D (2012) Evidence for a pan-Arctic sea-ice diatom diet in *Strongylocentrotus* spp. *Polar Biol* 35: 1281–1287.

22. Kattner G, Hagen W (1998) Lipid metabolism of the Antarctic euphausiid *Euphausia crystallorophias* and its ecological implications. *Mar Ecol Prog Ser* 170: 203–213.
23. Moline MA, Claustre H, Frazer TK, Schofield O, Vernet M (2004) Alteration of the food web along the Antarctic Peninsula in response to a regional warming trend. *Glob Chang Biol* 10: 1973–1980.
24. Flores H, van Francker JA, Siegel V, Haraldsson M, Strass V, et al. (2012) The Association of Antarctic Krill *Euphausia superba* with the Under-Ice Habitat. *PLoS ONE* 7: e31775.
25. Hubold G (1985) Stomach contents of the Antarctic silver fish *Pleuragramma antarcticum* from the Southern and Eastern Weddell Sea (Antarctica). *Polar Biol* 5: 43–48.
26. La Mesa M, Eastman JT, Vacchi M (2004) The role of notothenioid fish in the food web of the Ross Sea shelf waters: a review. *Polar Biol* 27: 321–338.
27. Hoshiai T, Tanimura A, Fukuchi M, Watanabe K (1989) Feeding by the Nototheniid fish, *Pagothenia borchgrevinkii* on the ice-associated copepod, *Paralabidocera antarctica*. *Proc NIPR Symp Polar Biol* 2: 61–64.
28. Micol T, Jouventin P (2001) Long term population trends in seven Antarctic seabirds at Pointe Géologie Terre Adélie. *Polar Biol* 24: 175–185.
29. Ridoux V, Offredo C (1989) The diets of five summer breeding seabirds in Adélie Land, Antarctica. *Polar Biol* 9: 137–145.
30. Veit RR, Hunt GL (1991) Broadscale density and aggregation of pelagic birds from a circumnavigational survey of the Antarctic ocean. *Auk* 108: 790–800.
31. Croxall JP, Trathan P N, Murphy EJ (2002) Environmental change and Antarctic seabird populations. *Science* 297: 1510–1514.
32. Ainley DG, Wilson PR, Barton KR, Ballard G, Nur N, et al. (1998) Diet and foraging effort of Adélie penguins in relation to pack-ice conditions in the southern Ross Sea. *Polar Biol* 20: 311–319.
33. Cottin M, Raymond B, Kato A, Amélineau F, Le Maho Y, et al. (2012) Foraging strategies of male Adélie penguins during their first incubation trip in relation to environmental conditions. *Mar Biol* 159, 1843–1852.
34. Spreen G, Kaleschke L, Heygster G (2008), Sea ice remote sensing using AMSR-E 89-GHz channels, *J Geophys Res*, 113, C02S03
35. Chastel O, Weimerskirch H, Jouventin P (1993) High annual variability in reproductive success and survival of an Antarctic seabird, the Snow Petrel *Pagodroma nivea* - a 27-year study. *Oecologia* 94: 278–285.
36. Barbraud C, Weimerskirch H (2001) Contrasting effects of the extent of sea-ice on the breeding performance of an Antarctic top predator, the Snow Petrel *Pagodroma nivea*. *J Avian Biol* 32: 297–302.
37. R Development Team 2.12.1 (2010-12-16) The R Foundation for Statistical Computing
38. Williams CT, Buck CL (2010) Using fatty acids as dietary tracers in seabird trophic ecology: theory, application and limitations. *J Ornithol* 151: 531–43.
39. Quetin LB, Ross RM (2001) Environmental variability and its impact on the reproductive cycle of Antarctic krill. *Am Zool* 41: 74–89.
40. Nicol S (2006) Krill, currents, and sea ice: *Euphausia superba* and its changing environment. *BioScience* 56: 111–120.
41. Ono A, Moteki M, Amakasu K, Toda R, Horimoto N, et al. (2011) Euphausiid community structure and population structure of *Euphausia superba* off Adélie Land in the Southern Ocean during austral summer 2003, 2005 and 2008. *Polar Sci* 5: 146–165
42. La Mesa M, Eastman JT (2012) Antarctic silverfish: life strategies of a key species in the high-Antarctic ecosystem. *Fish and Fisheries* 13: 241–266.
43. Chérel Y (2008) Isotopic niches of emperor and Adélie penguins in Adélie Land, Antarctica. *Mar Biol* 154: 813–821.
44. Montes-Hugo MA, Vernet M, Martinson D, Smith R, Iannuzzi R (2008) Variability on phytoplankton size structure in the western Antarctic Peninsula (1997–2006). *Deep Sea Res Part 2 Top Stud Oceanogr* 55: 2106–2117.
45. Siegel V, Loeb V (1995) Recruitment of Antarctic krill (*Euphausia superba*) and possible causes for its variability. *Mar Ecol Prog Ser* 123: 45–56.
46. Atkinson A, Siegel V, Pakhomov E, Rothery P (2004) Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature* 432: 100–103.
47. Barbraud C, Weimerskirch H (2006) Antarctic birds breed later in response to climate change. *P Natl Acad Sci-Biol* 103: 6248–6251.
48. Beaman RJ, O'Brien PE, Post AL, De Santis L (2011) A new bathymetry model for the Terre Adélie and George V continental margin, East Antarctica. *Antarct Sci* 23: 95–103.