



HAL
open science

Cercarial emergence pattern of *Schistosoma haematobium* from Libreville, Gabon

Rodrigue Mintsa Nguema, H. Moné, Moudachirou Ibikounlé, Krystina Mengué-Ngou-Milama, Maryvonne Kombila, Gabriel Mouahid

► **To cite this version:**

Rodrigue Mintsa Nguema, H. Moné, Moudachirou Ibikounlé, Krystina Mengué-Ngou-Milama, Maryvonne Kombila, et al.. Cercarial emergence pattern of *Schistosoma haematobium* from Libreville, Gabon. *Parasite*, 2014, 21 (3), pp.5. 10.1051/parasite/2014004 . halsde-00944713

HAL Id: halsde-00944713

<https://hal.science/halsde-00944713v1>

Submitted on 11 Feb 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cercarial emergence pattern of *Schistosoma haematobium* from Libreville, Gabon

Rodrigue Mintsá-Nguéma¹, H  l  ne Mon  ^{2,3}, Moudachirou Ibikounl  ⁴, Krystina Mengu  -Ngou-Milama⁵, Maryvonne Kombila⁶, and Gabriel Mouahid^{2,3,*}

¹ Institut de Recherche en   cologie Tropicale (IRET), CENAREST-Gabon, BP 13354, Libreville, Gabon

² Universit   Perpignan Via Domitia,   cologie et   volution des Interactions, UMR 5244, 66860 Perpignan, France

³ CNRS,   cologie et   volution des Interactions, UMR 5244, 66860 Perpignan, France

⁴ D  partement de Zoologie et G  n  tique, Facult   des Sciences et Techniques, Universit   d'Abomey-Calavi, 01BP526, Cotonou, B  nin

⁵ Service de Parasitologie-Mycologie-M  decine Tropicale, Facult   de M  decine de Tours, CHRU Bretonneau, 37044 Tours, France

⁶ D  partement de Parasitologie-Mycologie, Universit   des Sciences de la Sant   de Libreville, BP 4009, Libreville, Gabon

Received 7 October 2013, Accepted 30 January 2014, Published online 10 February 2014

Abstract – Although schistosomiasis has been a public health issue in Gabon for nearly a century, little is known about its current transmission dynamics. We analyzed the chronobiology of 137 cercarial emission profiles of *Schistosoma haematobium* from Libreville, the capital of Gabon, located in an open area for schistosomiasis. We found that 88% of the cercariae were shed between 11 a.m. and 3 p.m. and that the average pattern was of circadian type, with the average peak at 1 p.m., and representing 27% of the total number of cercariae of the day. The rhythms of emergence may be associated with environmental pressures on the parasite, especially those related to their definitive hosts.

Key words: Bilharziasis, Gabon, Transmission, *Schistosoma haematobium*, Cercarial emission, Chronobiology.

R  sum   – **Patrons d'  mergence cercarienne de *Schistosoma haematobium* de Libreville, Gabon.** Bien que la schistosomose soit un probl  me de sant   publique au Gabon depuis pr  s d'un si  cle, la dynamique actuelle de sa transmission est peu connue. Nous avons analys   la chronobiologie de 137 profils d'  mission cercarienne de *Schistosoma haematobium* de Libreville, la capitale du Gabon, situ  e dans une zone g  ographique ouverte pour la schistosomose. Nous avons trouv   que 88 % des cercaires   taient   mises entre 11 heures et 15 heures et que le profil moyen   tait de type circadien, avec un pic moyen situ      13 heures et repr  sentant 27 % du nombre total de cercaires journalier. Les rythmes d'  mergence sont associ  s aux pressions environnementales sur le parasite, particuli  rement celles relatives aux h  tes d  finitifs.

Introduction

Schistosomiasis is a public health problem throughout the world. At present, 779 million individuals are at risk of infection and more than 207 million people are infected, with 97% of the latter in Africa [20].

The first Bilharziasis reported in Gabon was a rectal schistosomiasis, called *Schistosomum haematobium*, which occurred in Libreville, the capital of Gabon, in 1923 [2]. It was later demonstrated that the agent responsible for this schistosomiasis was *Schistosoma guineensis*. Bladder schistosomiasis due to *S. haematobium* was described more than 40 years later [5] in the village of Moukoro, in the Nden   district in

the south of Gabon; this schistosomiasis was introduced from N'Kayi (formerly Jacob), located on the Congolese border of Gabon. The prevalence of *S. haematobium* infection in the population at that time was 50%. Its expansion into the rest of the country likely occurred quickly, from the south to the northwest, following the rivers and roads within the country. Indeed, *S. haematobium* was detected in 1967 near the Tchibanga Nyanga area in the southwest of Gabon [5] and in 1969 in the Palmh  v  as region, near Lambar  n   (middle Ogowe) [5]. In 1974, an epidemic was reported in Ekouk, North of Palmh  v  as, near Libreville [9], with populations living along the road connecting the Niari Valley in Congo to Libreville being the most vulnerable to infection. In 1981, *S. haematobium* left the main road and spread in different directions, especially

*Corresponding author: mouahid@univ-perp.fr

Table 1. Variation of the position of the cercarial emission peak for each individual snail.

Snail	NCEP	Hour					
		11 a.m.	12 a.m.	1 p.m.	2 p.m.	3 p.m.	4 p.m.
G1	9	1	6	1	1		
G2	8		5	3			
G3	8		4	4			
G4	9	5	4				
G5	9		2	7			
G6	13	1	4	5	2		1
G7	13		6	3	4		
G8	14		5	9			
G9	5		3	2			
G10	9	1	2	2	2	2	
G11	6	1	1	4			
G12	13	1	3	3	3	3	
G13	7		1	4	2		
G14	13		4	5	4		1
Total	137	10	50	52	18	5	2

NCEP: Number of cercarial emission profiles.

in the west of the country, in the province of Ogooué Maritime (Port-Gentil) and in the southeast, in the province of Upper Ogooué (Moanda). In 1982, seven of the nine provinces of Gabon were affected by urinary schistosomiasis, with the largest foci in Libreville, Koulamoutou in Ogooué Lolo Province, Bibora in Nyanga Province, and Palmhévées in Moyen Ogooué Province. The prevalence of infection was generally higher than that of *S. guineensis* [7, 10]. Thus, at present two schistosome species adapted to humans coexist in Gabon, which has led to research on the possible existence of hybrids [1, 10, 18].

Schistosoma haematobium is a species very sensitive to light [16, 17]. It was shown, in Côte d'Ivoire, that cercariae located in shady transmission sites situated in forest areas were emitted earlier (11 a.m.) than those located in transmission sites receiving direct sunlight, such as savanna areas (1:30 p.m.) [15]. The progression of *S. haematobium* from the South of Gabon to Libreville follows a vegetal axis represented mainly by savannas, degraded forests, and mangroves, resulting in conditions of strong light in potential sites of transmission, whereas the rest of the country (85%) is covered by rainforests [8]. The purpose of this study was to analyze *S. haematobium* cercarial emissions from transmission sites located in Libreville, where the forest has disappeared.

Material and methods

Biological material

The strain of *Schistosoma haematobium* originated from Melen (09°30'59" E, 00°23'46" N) in Libreville city, Gabon [10]. The F₀ generation was extracted from positive urine samples of nine patients and transported to France in experimentally infected *Bulinus globosus* from Toho-Todougba, Benin (2°11'21" E, 6°23'52" N). The strain was then maintained in jirds (*Meriones unguiculatus*) as definitive hosts. We used the

F₁ generation of the parasite for the cercarial emission pattern analyses.

Infection and maintenance of snails

The snails were exposed individually to five miracidia of the F₁ generation, a dose which is usually used in laboratories because it allows one to obtain an optimal percentage of snail infection [12, 13, 14, 19]. The exposed snails were maintained in aquaria containing well water in a temperature-controlled room at 24–25 °C. They were fed ad libitum with fresh lettuce. The photoperiod was balanced, with 12 h of light (from 6:00 a.m. to 6:00 p.m.) and 12 h of darkness (from 6:00 p.m. to 6:00 a.m.), with a simulated dawn (from 5:30 a.m. to 6:00 a.m.) and simulated dusk (from 5:30 p.m. to 6:00 p.m.). At the end of the prepatent period (the time between exposure of the snails to the miracidia and the first cercarial emission), the snails were placed individually in plastic cups containing 150 mL of well water, in a balanced photoperiod light: dark of 12:12.

Cercarial emission patterns

Every hour, starting at 7:00 a.m. (corresponding to the cercariae emitted from 6 a.m. to 7 a.m.), each infected snail was transferred individually to another glass containing the same volume of well water at the same temperature (24–25 °C). The water samples containing the cercariae were filtered through polyamide Nitrile filters (25- μ m mesh sieve). The cercariae retained on the filter were stained with Lugol's solution and counted under a dissecting microscope. To strengthen each profile, we retained only the emission profiles for which the total daily cercarial production was ≥ 50 (range: 53–837 cercariae per day). In total, 14 snails (G1 to G14) were analyzed and followed for 5–14 consecutive days, yielding a total of 137 cercarial emission profiles (Table 1).




Figure 1. Average percentage of cercariae of *Schistosoma haematobium* (Libreville, Gabon) emitted every hour from *B. globosus* (Benin), each exposed to five miracidia. Bars represent standard errors.

Ethics statement

The laboratory has received the permit No. A 66040 for experiments on animals, from both the French Ministère de l'Agriculture et de la Pêche and the French Ministère de l'Education Nationale de la Recherche et de la Technologie (Décret n° 87-848 du 19 octobre 1987). The housing, breeding, and care of the rodents followed the ethical requirements of our country. Animal experimentation followed the guidelines of the French CNRS. The different protocols used in this study have been validated by the French Veterinary Agency.

Results

The histogram representing the profile of the average daily cercarial emissions of Gabonese *S. haematobium* from *B. globosus* snails from Benin is presented in Figure 1. The emergence of cercariae tentatively began at 7 a.m. (i.e. at the beginning of the photophase), increased gradually from 10 a.m. to its peak at 1 p.m., and decreased gradually before stopping completely at 8 p.m. (i.e. 2 h after the beginning of the scotophase). The majority of the cercarial emissions (88%) occurred between 11 a.m. and 3 p.m. The average emission peak occurred at 1 p.m. and represented 26.3% of the total daily cercarial production.

The variation of the position of the cercarial emission peaks is presented in Table 1. The 137 peaks were situated between 11 a.m. and 4 p.m.: 10 peaked at 11 a.m., 50 at 12 a.m., 52 at 1 p.m., 18 at 2 p.m., 5 at 3 p.m., and 2 at 4 p.m. For each individual snail and for each profile, only one peak was observed (circadian rhythm); none of the snails had all its peaks situated at the same hour. Each snail had 2, 3, 4, 5, or 6 peak positions according to its profiles. For example, the snail G1 had four peak positions (one peak at 11 a.m., six at 12 a.m., one at 1 p.m., and one at 2 p.m.) and the snail G4 had two peak positions (five peaks at 11 a.m. and four at 12 a.m.). However, for each snail, two peak positions were over-represented, 12 a.m. and 1 p.m. Five snails (G2, G3, G5, G8, and G9)

had 100% of their peaks situated either at 12 a.m. or at 1 p.m.; the other snails had from 44.4 to 83.3% of their peaks situated either at 12 a.m. or at 1 p.m.

Discussion

Our results on the chronobiology of *S. haematobium* cercarial emission in Libreville showed that the rhythm of emergence was of a circadian type, with 88% of the cercariae emitted between 11 a.m. and 3 p.m. and an average emergence peak at 1 p.m., representing 27% of the cercariae emitted daily. The average emergence peak is similar to that obtained in Ivory Coast for a strain of *S. haematobium* located in the vegetal zone called wooded savanna corresponding to the sub-Saharan climate [15]. Historically, the course of migration of our strain of *S. haematobium* was from the south to northwestern Gabon, corresponding, for the most part, to savanna and mangrove areas [8].

The cercarial emergence behavior is genetically supported [23], with this behavior remaining unchanged when the schistosome is subject to different environmental pressures at the level of its intermediate snail host. For example, the emission profile of *S. haematobium* from the snail *B. truncatus* does not change when the snail is also infected with another species of schistosome, *Schistosoma bovis* [12], and the emission profile of *Schistosoma mansoni* from the snail *Biomphalaria glabrata* does not change when the snail is infected with another fluke, *Ribeiroia marini* [24]. Moreover, the emission patterns of two strains (early and late) of *S. mansoni* remain unchanged when both parasites are present simultaneously in *B. glabrata* [25], and the emission profile of *S. mansoni* remains unchanged when infected *B. glabrata* is subject to the presence of a competitor snail, *Marisa cornuarietis* [4].

From an evolutionary point of view, the diversity of the rhythms of emergence is usually well correlated with the diversity of behavior of the definitive host species [3]. In the region of Libreville, the circadian rhythm of emissions, with a peak at 1 p.m., can be explained by human contact with water, which is essentially related to bathing activities during the hottest hours of the day, between 11 a.m. and 4 p.m. In other areas, the practice of activities other than swimming, such as artisan fishing, can result in a change in the cercarial emission pattern to an ultradian pattern, with a primary peak at midday but secondary peaks at dawn and dusk, when fishermen are in contact with water, as has been shown in *S. mansoni* in Benin [6]. Some strains of *S. mansoni* have adapted to the presence of definitive hosts other than humans, such as the rodent, *Rattus rattus*, with the cercarial emission peak becoming later, as in Guadeloupe [22], or strictly nocturnal, as in Oman [11]. This has also been shown for the species *Schistosoma japonicum*, with cercarial emergence peaks occurring later or becoming nocturnal, due to the reservoir hosts, particularly rodents, used by this species [21]. Indeed, the involvement of murine reservoir hosts in the transmission dynamics has led to intraspecific variability in cercarial emission, with distinct chronobiological races corresponding to sites in which humans and/or reservoir hosts are the definitive hosts. The cercarial emission peaks at 1 p.m. in the open area of Libreville may be related to different behaviors of the definitive

host, man, in different types of landscape, with the shady forest requiring activities be performed earlier in the day to compensate for the lack of light, whereas activities are performed later in open areas such as savannas and mangroves.

Acknowledgements. This work was financially supported by the CNRS, the University of Perpignan, the University of Health Sciences and the National Agency Awards of Gabon. Thanks are due to Lambert Ndong Obiang and Callixte Nkourou Mba from the Service d'Hygiène Publique of Gabon. The French Laboratory is a World Health Organization Collaborating Center for Biological Control and Snail/Parasite relationships. The authors wish to report that there are no competing interests.

References

- Burchard GD, Kern P. 1985. Probable hybridization between *S. intercalatum* and *S. haematobium* in western Gabon. *Tropical and Geographical Medicine*, 37, 119–123.
- Clapier PN. 1923. *Schistosomum haematobium* à Libreville. Bilharziose autochtone, bilharziose importée. *Bulletin de la Société de Pathologie Exotique*, 47, 531–534.
- Combes C, Fournier A, Moné H, Théron A. 1994. Behaviours in trematode cercariae that enhance parasite transmission: patterns and processes. *Parasitology*, 109, S3–S13.
- Ferrer JR, Pointier JP, Théron A, Moné H. 1991. Influence of the nontarget mollusc *Marisa cornuarietis* on the hourly cercarial production of *Schistosoma mansoni* from *Biomphalaria glabrata*. *Journal of Parasitology*, 77, 798–800.
- Gilles JC. 1969. La bilharziose au Gabon, in 4e Conférence technique OCEAC, 484–487.
- Ibikounlé M, Moné H, Abou Y, Kinde-Gazard D, Sakiti NG, Mouahid G, Massougoudji A. 2012. Premier cas d'émission cercarienne de type infradien chez *Schistosoma mansoni* dans deux foyers du sud-Bénin. *International Journal of Biological and Chemical Sciences*, 6, 1081–1089.
- Mabika-Mamfoumbi M, Moussavou-Boussougou MN, Nzenze-Afene S, Owono-Medang M, Bouyou-Okotet M, Kendzo E, Kombila M. 2009. Prevalence evaluation of intestinal parasites in rural and sub-urban area in Gabon. *Bulletin Médical d'Owendo*, 12, 85–88.
- Maloba Makanga JD. 2011. Mosaique forêt-savane et exploitation des ressources forestières du Gabon. *Geo-Eco-Trop*, 35, 41–50.
- Mc-Cullough FS. 1975. Les Schistosomiasés en République du Gabon, in Rapport de mission 4–24 juin et 11 octobre – 27 novembre 1974. O.M.S: Brazzaville. p. 65.
- Mintsa-Nguéma R, Mengue-Ngou-Milama K, Kombila M, Richard-Lenoble D, Tisseyre P, Ibikounlé M, Moné H, Mouahid G. 2010. Morphometric and molecular characterizations of schistosome populations in Estuaire province Gabon. *Journal of Helminthology*, 84, 81–85.
- Mouahid G, Idris MA, Verneau O, Théron A, Shaban MA, Moné H. 2012. A new chronotype of *Schistosoma mansoni*. Adaptive significance. *Tropical Medicine and International Health*, 17, 727–732.
- Mouahid A, Moné H, Chaib A, Théron A. 1991. Cercarial shedding patterns of *Schistosoma bovis* and *S. haematobium* from single and mixed infections of *Bulinus truncatus*. *Journal of Helminthology*, 65, 8–14.
- Mouahid A, Théron A. 1986. *Schistosoma bovis* : patterns of cercarial emergence from snails of the genera *Bulinus* and *Planorbarius*. *Experimental Parasitology*, 62, 389–393.
- Mubila L, Rollinson D. 2002. Snail-parasite compatibility and prevalence of *Schistosoma haematobium* on the shores of Lake Kariba, Zambia. *Annals of Tropical Medicine & Parasitology*, 96, 165–173.
- N'goran E, Brémond P, Sellin E, Sellin B, Théron A. 1997. Intraspecific diversity of *Schistosoma haematobium* in West Africa: chronobiology of cercarial emergence. *Acta Tropica*, 66, 35–44.
- Noda S, Sato K, Katsumata T, Nojima H, Muhoho ND. 1986. The influence of shadowing on emergence of cercarial emergence of *Schistosoma haematobium* during day time. *Japanese Journal of Parasitology*, 35, 249–251.
- Raymond KM, Probert AJ. 1987. The effect of light and darkness on the production of cercariae of *Schistosoma haematobium* from *Bulinus globosus*. *Journal of Helminthology*, 61, 291–296.
- Richard-Lenoble D, Kombila M, Duong TH, Gendrel D. 1993. Bilharziose à *Schistosoma intercalatum*, bilharziose récente et oubliée. *Revue du Praticien*, 43, 432–438.
- Sène M, Southgate VR, Vercruysse J. 2004. *Bulinus truncatus*, hôte intermédiaire de *Schistosoma haematobium* dans le bassin du fleuve Sénégal. *Bulletin de la Société de Pathologie Exotique*, 97, 29–32.
- Steinmann P, Keiser J, Tanner M, Utzinger J. 2006. Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infectious Diseases*, 6, 411–425.
- Su J, Zhou F, Lu DB. 2013. A circular analysis of chronobiology of *Schistosoma japonicum* cercarial emergence from hilly areas of Anhui, China. *Experimental Parasitology*, 135, 421–425.
- Théron A. 1984. Early and late shedding patterns of *Schistosoma mansoni* cercariae: ecological significance in transmission to human and murine hosts. *Journal of Parasitology*, 70, 652–655.
- Théron A, Combes C. 1988. Genetic analysis of cercarial emergence rhythms of *Schistosoma mansoni*. *Behavior Genetics*, 18, 201–209.
- Théron A, Moné H. 1986. Shedding patterns of *Schistosoma mansoni* and *Ribeiroia marini* cercariae from a mixed infection of *Biomphalaria glabrata*. *Journal of Helminthology*, 60, 255–259.
- Théron A, Mouahid G, Moné H. 1997. *Schistosoma mansoni* : cercarial shedding patterns from a mixed infection of *Biomphalaria glabrata* with two (early and late) chronobiological variants. *Parasitology Research*, 83, 356–358.

Cite this article as: Mintsa-Nguéma R, Moné H, Ibikounlé M, Mengué-Ngou-Milama K, Kombila M & Mouahid G: Cercarial emergence pattern of *Schistosoma haematobium* from Libreville, Gabon. *Parasite*, 2014, 21, 3.



An international open-access, peer-reviewed, online journal publishing high quality papers on all aspects of human and animal parasitology

Reviews, articles and short notes may be submitted. Fields include, but are not limited to: general, medical and veterinary parasitology; morphology, including ultrastructure; parasite systematics, including entomology, acarology, helminthology and protistology, and molecular analyses; molecular biology and biochemistry; immunology of parasitic diseases; host-parasite relationships; ecology and life history of parasites; epidemiology; therapeutics; new diagnostic tools.

All papers in Parasite are published in English. Manuscripts should have a broad interest and must not have been published or submitted elsewhere. No limit is imposed on the length of manuscripts.

Parasite (open-access) continues **Parasite** (print and online editions, 1994-2012) and **Annales de Parasitologie Humaine et Comparée** (1923-1993) and is the official journal of the Société Française de Parasitologie.

Editor-in-Chief:
Jean-Lou Justine, Paris

Submit your manuscript at
<http://parasite.edmgr.com/>