



# Genetic diversity, clonality and connectivity in the scleractinian coral *Pocillopora damicornis*: a multi-scale analysis in an insular, fragmented reef system

Mehdi Adjeroūd, Aurélie Guérécheau, Jeremie Vidal-Dupiol, Jean-François Flot, Sophie Arnaud-Haond, François Bonhomme

## ► To cite this version:

Mehdi Adjeroūd, Aurélie Guérécheau, Jeremie Vidal-Dupiol, Jean-François Flot, Sophie Arnaud-Haond, et al.. Genetic diversity, clonality and connectivity in the scleractinian coral *Pocillopora damicornis*: a multi-scale analysis in an insular, fragmented reef system. *Marine Biology*, 2013, 161 (3), pp.531-541. 10.1007/s00227-013-2355-9 . halsde-01007685

**HAL Id: halsde-01007685**

**<https://hal.science/halsde-01007685>**

Submitted on 3 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Genetic diversity, clonality and connectivity in the scleractinian coral *Pocillopora damicornis*: a multi-scale analysis in an insular, fragmented reef system

Mehdi Adjeroud<sup>1,2</sup> · Aurélie Guérécheau<sup>1</sup> · Jeremie Vidal-Dupiol<sup>3,4</sup> · Jean-François Flot<sup>5</sup> · Sophie Arnaud-Haond<sup>6</sup> · François Bonhomme<sup>7</sup>

1 Institut de Recherche pour le Développement, Unité 227 CoRéUs2, BP A5, 98848 Nouméa Cedex, New Caledonia

2 Laboratoire d'Excellence "CORAIL", 58 Avenue Paul Alduy, 66860 Perpignan Cedex, France

3 CNRS, Ecologie et Evolution des Interactions, UMR 5244, Université de Perpignan Via Domitia, 58 Avenue Paul Alduy, 66860 Perpignan, France

4 Université de Perpignan Via Domitia, Ecologie et Evolution des Interactions, UMR 5244, 58 Avenue Paul Alduy, 66860 Perpignan, France

5 Max Planck Institute for Dynamics and Self-Organization, Biological Physics and Evolutionary Dynamics, Bunsenstraße 10, 37073 Göttingen, Germany

6 Ifremer, UMR 212 Ecosystèmes Marins Exploités, Institut Français de Recherche pour l'Exploitation de la Mer, BP 171, 34203 Sète Cedex, France

7 CNRS, Institut des Sciences de l'Evolution, UMR 5554, Université de Montpellier 2, cc 63, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

**Abstract** Clonality and genetic structure of the coral *Pocillopora damicornis* sensu lato were assessed using five microsatellites in 12 populations from four islands of the Society Archipelago (French Polynesia) sampled in June 2008. The 427 analysed specimens fell into 132 multilocus genotypes (MLGs), suggesting that asexual reproduction plays an important role in the maintenance of these populations. A haploweb analysis of ITS2 sequences of each MLG was consistent with all of them being conspecific. Genetic differentiation was detected both between and within islands, but when a single sample per MLG was included in the analyses, the populations turned out to be nearly panmictic. These observations provide further evidence of the marked variability in reproductive strategies and genetic structure of *P. damicornis* throughout its geographic range; comparison with results previously obtained for the congeneric species *Pocillopora meandrina* underlines the importance of life history traits in shaping the genetic structure of coral populations.

## Introduction

Coral reefs are one of the most diverse ecosystems on Earth and provide goods and services to ~500 million people (Moberg and Folke 1999; Wilkinson 2008). However, in recent decades, coral reefs have been confronted with various types of natural and anthropogenic disturbances and are expected to be highly vulnerable to future climatic changes (Hughes et al. 2003; Bellwood et al. 2004; Hoegh-Guldberg et al. 2007; Adjeroud et al. 2009; Hoegh-Guldberg 2009). Coral reefs are mostly supported by colonial calcifying organisms, the scleractinian corals, which are also a key component of coral reef health and diversity as they provide a variety of shelter and food for thousands of reef organisms (Lesser 2004; Bellwood et al. 2006). Thus, the recent widespread mortality of coral species in response to large-scale disturbances and chronic stressors has caused extensive habitat degradation and, in a number of cases, phase shifts in community structure (Gardner et al. 2003; Pandolfi et al. 2003; Bellwood et al. 2004, 2006). Classically, these phase shifts have involved the replacement of scleractinian corals by macroalgae or other non-reef-building benthic organisms, with

important consequences for the structure and functioning of coral reef ecosystems (McManus and Polsenberg 2004).

In this context of an ongoing “coral reef crisis” (Bellwood et al. 2004), considerable efforts have been recently made to conserve and manage coral reefs with, for example, the implementation of marine protected area (MPA) networks (Mora et al. 2006; Almany et al. 2009). Such conservation actions largely depend on the knowledge of the key biological and physical factors that govern the structure and dynamics of reef communities. Among these factors, the level of gene flow and the scale of genetic connectivity have been shown to have a large influence on the vulnerability and resilience of coral populations to perturbations and are thus key parameters for the location, size and spacing of MPAs (Sale et al. 2005; Miller and Ayre 2008; Almany et al. 2009; Botsford et al. 2009; Jones et al. 2009).

In scleractinian corals, as in most marine species, the level of connectivity is often impossible to infer from direct observations. Except for some species of brooding corals that release relatively large larvae, the small size of coral propagules and the absence of available structure for tagging them make it impossible to follow dispersing individuals for long distances (Harrison 2011). The pelagic larval duration (PLD) has sometimes been used to estimate the maximum connectivity. However, correlation between PLD and level of gene flow remains elusive for several biological and methodological reasons (Riginos et al. 2011; Selkoe and Toonen 2011; Tremblay et al. 2012; Faurby and Barber 2012). First, PLDs can be extremely difficult to assess, as it is a plastic life history trait (Nozawa and Harrison 2008) under strong influence of environmental conditions such as temperature (Duarte 2007; O’Connor et al. 2007). Moreover, most larvae are able to change their vertical positioning in the water column and, together with the complexity of oceanic currents, this behaviour makes it highly difficult to correlate PLD and maximum distance of larval migration (Palumbi 1994; Parsons 1996). Therefore, connectivity among coral populations has been mainly assessed by the examination of its indirect effects on their genetic structure and differentiation (Ayre and Hughes 2000; Elmhirst et al. 2009).

Scleractinian corals exhibit a variety of modes of reproductive systems that may directly influence their potential for dispersal and the level of clonality within local populations (Stoddart 1984a, b; Ayre et al. 1997; Ayre and Hughes 2000; Nishikawa et al. 2003; Heyward and Negri 2012; among others). Corals can be hermaphroditic or gonochoric, and brooders or broadcasters (Harrison 2011). Broadcasters are thought to be great dispersers, whereas brooders are considered to be poor dispersers and have the potential for self-fertilization (Carlson 1999; Ayre and Hughes 2000; Nishikawa et al. 2003; Miller and Ayre 2008; Underwood et al. 2009). These two modes coexist in certain species and may vary across their geographic range (Harrison 2011). Larvae that contain symbiotic dinoflagellates are also thought to disperse farther than aposymbiotic ones as they can obtain energy from their symbionts (Richmond 1987; Hariri et al. 2010; Figueiredo et al. 2012). Moreover, asexual reproduction exists in corals in a variety of forms, including fragmentation, budding or polyp expulsion, asexually produced planulae and embryo fragmentation (Richmond 1987; Harrison 2011; Heyward and Negri 2012). The relative contributions of sexual and asexual reproduction largely determine the level of clonality within local populations and the pattern of genetic structure and connectivity among metapopulations (Stoddart 1984a, b; Ayre et al. 1997; Ayre and Hughes 2000; Ayre and Miller 2004; Becheler et al. 2010).

The scleractinian coral *Pocillopora damicornis* sensu lato (following Veron 2000) has one of the largest spatial repartitions of Indo-Pacific corals and is commonly found in tropical and subtropical regions from the Red Sea and East African coast to the Eastern Pacific (Veron 2000). The taxonomy of *P. damicornis* is presently in a state of flux, with several molecular studies and morphological species suggesting that it is actually composed of cryptic species, some of which sympatric (Richmond and Jokiel 1984; Flot 2007a; Flot et al. 2008; Souter 2010; Pinzón and Lajeunesse 2011; Schmidt-Roach et al. 2013). *P. damicornis* has been described as a brooder in most studies (Stoddart 1983; Richmond 1987) but as a broadcast spawner in some populations (Glynn et al. 1991;

Combosch and Vollmer 2011; Schmidt-Roach et al. 2012). This coral can also release asexually produced planulae (Stoddart 1983; Ayre and Miller 2004; Sherman et al. 2006) and may rely on fragmentation, thus facilitating the persistence of local populations (Richmond 1987; Adjeroud and Tsuchiya 1999). As expected from such a diversity of reproductive strategies, contrasting patterns of genetic differentiation and connectivity have been observed (see Table 3 in Pinzón et al. 2012), ranging from fine-scale genetic subdivision along East African (Souter et al. 2009), Southwest Australian (Stoddart 1984a, b; Whitaker 2006) and Panaman (Combosch and Vollmer 2011) coasts to large-scale panmixia on the Great Barrier Reef (Ayre et al. 1997; Ayre and Hughes 2000). Clonality levels in *P. damicornis* populations seem also highly variable, with some populations considered highly clonal (Stoddart 1984a, b; Souter et al. 2009; Gorospe and Karl 2013), whereas others exhibit low (Adjeroud and Tsuchiya 1999) or almost no clonality (Benzie et al. 1995; Ayre et al. 1997; Ayre and Miller 2004; Sherman et al. 2006; Combosch and Vollmer 2011).

The aim of the present study was to examine the genetic diversity, clonality and connectivity of *P. damicornis* sensu lato in an insular, fragmented reef system. We chose to focus on populations from the Society Archipelago (French Polynesia) in order to compare our results with those of a similar study conducted previously on *P. meandrina* in the same region (Magalon et al. 2005). As in this previous study, we used a hierarchical sampling design integrating both the regional scale (among four islands) and the local scale (among three sites on each island). *P. meandrina* is abundant on exposed reef fronts and mainly reproduces sexually through broadcast spawning gametes, resulting in a high gene flow and reduced genetic differentiation among regional populations (Magalon et al. 2005). In contrast, *P. damicornis* sensu lato is mainly found in sheltered back reef habitats (Veron 2000) and is able to reproduce asexually, two factors that should result in more clonality and less gene flow compared with *P. meandrina*. Both species are abundant in the Society Islands, which is therefore an ideal place to test the hypothesis of an influence of life history traits and ecology on the genetic structure of coral populations and on their connectivity.

## Materials and methods

### Sampling

Populations identified morphologically as *P. damicornis* following Veron and Pichon (1976) were sampled in four islands of the Society Archipelago (Moorea, Raiatea, Taha'a and Tahiti), French Polynesia (CITES export: FR089800007E, CITES import: FR089800007-I). In order to integrate the regional and the local scales, a hierarchical sampling design was established, with samples collected from three sites on each of these four islands (Moorea: MH, MP, MT; Raiatea: RA, RT, RU; Taha'a: TaF, TaT, TaV; Tahiti: TP, TT, TV; Fig. 1) in June 2008. Between 25 and 43 coral colonies were sampled at each site (Table 1), for a total of 427 colonies. The collecting sites were located on the fringing reef, at approximately 2 m depth. Colonies were sampled along transect lines with a minimum distance of 2 m between two sampled colonies to minimize the collection of several ramets per genet (Adjeroud and Tsuchiya 1999; Magalon et al. 2005; Paz-García et al. 2012). Each sample was obtained by cutting off the tip of a coral branch (1–2 cm). Fragments were placed in zip-lock plastic bags and held in seawater before being placed in CHAOS buffer (4 M guanidium thiocyanate, 0.5 % *N*-lauroyl sarcosine sodium, 25 mM Tris–HCl pH 8, 0.1 M 2-mercaptoethanol) modified from Fukami et al. (2004) following Flot (2007a). Samples were then conserved at ambient temperature and protected from light. DNA was purified following a conventional phenol/chloroform (1:1) protocol and precipitated with isopropanol.

### Genotyping

Samples were genotyped using five microsatellite markers: PV2, PV7 (Magalon et al. 2004), Pd3-002, Pd3-004 and Pd3-005 (Starger et al. 2008), previously shown to be coral-specific and polymorphic in *P. damicornis*. The polymorphism of PV2 and PV7 had previously been tested on populations from French Polynesia, whereas Pd3-002, Pd3004 and Pd3-005 had been tested on populations from Indonesia and Taiwan (Starger et al. 2008). In the present study, intra-population and inter-population polymorphisms were both checked for each marker using Spreadex® gel electrophoresis (Elchrom Scientific). Two multiplex polymerase chain reactions (PCR) were then performed on each sample using fluorescently labelled primers. The QIAGEN multiplex PCR kit was used with a final volume of 10 µL, following the microsatellite protocols. The PCR programme included 15 min at 95 °C followed by 25 cycles of 30 s at 94 °C, 90 s at 57 °C and 60 s at 72 °C. A final elongation step consisted of 30 min at 60 °C. Amplified fragments were visualized on an ABI Prism 3130 XL sequencer (Applied Biosystems) using a GeneScan 500-Rox ladder. Genotype calling was performed using GENEMAPPER 4.0 (Applied Biosystems), and each genotype was visually checked.

#### Molecular delimitation of species

Several studies have reported cryptic diversity within *P. damicornis* sensu lato (Richmond and Jokiel 1984; Flot 2007a; Flot et al. 2008; Souter 2010; Pinzón and Lajeunesse 2011; Schmidt-Roach et al. 2013). To rule out a possibly confounding effect of cryptic species on our population genetic analyses, we used the published coral-specific PCR primers ITSc2-5 and R28S1 (Flot and Tillier 2006; Flot et al. 2008) to amplify and sequence the nuclear ITS2 for one representative individual of each MLG. The haplotypes of length-variant heterozygotes were inferred directly from the patterns of double peaks in the forward and reverse chromatograms using the program Champuru (Flot et al. 2006; Flot 2007b); heterozygotes having haplotypes of equal lengths were subsequently phased using the programs SeqPHASE (Flot 2010) and PHASE (Stephens et al. 2001). All haplotype sequences were deposited in GenBank (accession numbers: KF846307– KF846519); the processed chromatograms (in a Sequencher file) and the haplotype alignment were deposited in Dryad (doi:10.5061/dryad.b5r86). A median-joining haplotype network was constructed using the program Network (Fluxus Technology) and exported as PDF using Network Publisher (from the same company). This haplonet was subsequently imported into Inkscape and turned into a haploweb (Flot et al. 2010) by adding connections between haplotypes found co-occurring in heterozygotes individuals.

#### Data set preparation

In order to associate each ramet (i.e., each physical individual colony that was sampled) to its genet (i.e., clonal lineage encompassing all ramets deriving from the same zygote), the genotypic resolution of the five microsatellites analysed here was checked by visualizing in GENCLONE the number of distinct multilocus genotypes (MLGs) detected for all possible subsets of loci (Arnaud-Haond and Belkhir 2007). Then, for each repeated MLG, the probability of obtaining the same MLG more than once from distinct random reproductive events was estimated as  $p_{\text{sex}}(F_{\text{IS}})$  (Arnaud-Haond et al. 2007).

#### Population genetics analyses

The magnitude and direction of departure from the Hardy–Weinberg equilibrium (HWE) at each locus were estimated using GENETIX software version 4.05.2 (Belkhir et al. 1996) with 10,000 permutations. These parameters were expressed as Wright's  $F_{\text{IS}}$ , which is the percentage deviation from expected heterozygosity in each population (Wright 1950). In order to estimate clonal richness and heterogeneity in each population, the clonal richness  $R$  was estimated as :

$$R = \frac{(G - 1)}{(N - 1)}$$

where  $G$  is the number of MLGs and  $N$  the number of ramets sampled (following Arnaud-Haond et al. 2007). This index ranges from 0 (for a monoclonal stand) to 1 (when all the different samples analysed correspond to a single MLG) and provides a lower bound estimate of the clonal versus sexual input (Arnaud-Haond et al. 2007). In order to test for significant genetic variation among sites, the genetic differentiation between populations was estimated using Weir and Cockerham's  $F_{ST}$  estimator  $\vartheta$  (Weir and Cockerham 1984) with 10,000 permutations between samples. However, the differentiation captured by  $F_{ST}$  among populations in which both sexual and asexual reproduction occurs can be the result of different clones being dominant in each population. A second data set was obtained by keeping only one sample for each MLG in each population.  $F_{ST}$  was then estimated for this data set restricted to colonies of different sexual origins. In order to test for genetic variation within populations, Wright's  $F_{IS}$  at each locus was also estimated for this data set. GENETIX was then used to construct a correspondence analysis (CA) to visualize the genetic distance among populations in a multidimensional space. The output of the CA was simplified by plotting only the centroids of all samples within a population.

## Results

### Species delimitation

All 23 ITS2 haplotypes in the network were connected through heterozygous individuals, with no obvious departure from panmixia: the most frequent haplotypes (represented by the large circles on Fig. 2) were found co-occurring in many heterozygotes (represented by thick red curves connecting these circles), whereas rare haplotypes were found co-occurring with frequent ones but not with other rare haplotypes. Hence, the 132 individuals sequenced belonged to a single field for recombination sensu Doyle (1995) and therefore were conspecific following the criterion of mutual allelic exclusivity (Flot et al. 2010). Our studied populations shared several haplotypes with *P. damicornis* from Oahu (Flot et al. 2008, analysis not shown), suggesting that the same species is found both in Hawaii and in the Society Archipelago.

### Genotypic variation

Table 1 presents the values of genotypic diversity indices calculated for the entire data set. Six of the 12 populations significantly deviated from HWE expectations, as estimated by Wright's  $F_{IS}$  (Table 1). When analysing singlelocus genotypic variation, 43 Hardy–Weinberg tests (of 72) showed significant deviation from HWE (Online Resource 1). Of these, 32 showed heterozygote excesses, and the remaining 11 exhibited deficits. Allelic frequencies and allelic richness recorded at the 12 populations are presented in Online Resource 2.

The hypothesis of multiple reproduction events being responsible for the presence of several individuals per MLG was rejected (with  $p_{sex}(F_{IS}) < 0.05$ ) for about half of the repeated MLGs, but not for the other half that included mostly the predominant genotypes in the populations. This could be either due to the lack of power associated with the  $p_{sex}(F_{IS})$  estimates in cases of low allelic and low clonal richness or due to the presence of genotypic variation that remained undetected using the set of five loci available. In every population, some MLGs were repeated, indicating that asexual reproduction or fragmentation occurred at all sampling sites. Sixteen MLGs were shared by different samples, of which only 11 were shared by populations from different islands. Six MLGs were shared by Moorea and Tahiti (ca. 30 km distance), three by Taha'a and Raiatea (ca. 10 km distance), one by Moorea and Taha'a (ca. 200 km distance) and one by Moorea, Taha'a and Raiatea. No

MLGs were common to Taha'a and Tahiti (ca. 230 km distance) or to Raiatea and Tahiti (ca. 220 km distance). However, after removal of all duplicated MLGs, the newly estimated  $F_{IS}$  showed no significant departure from HWE. Clonal richness ( $R$ ) estimated only on the basis of unique MLGs was therefore an underestimate of the genotypic richness potentially present in the sampling sets. Nevertheless,  $R$  was highly variable among locations (Table 1), reaching 0.600 in Moorea (MT) and 0.097 in a population from Raiatea (RU) that was entirely dominated by four MLGs.

#### Genetic differentiation among sampling sites

Population differentiation was assessed using pairwise  $F_{ST}$  comparisons (Table 2). On the basis of the entire data set (including replicates),  $F_{ST}$  was always significant ( $p < 0.01$ ) except for two pairs of samples from Moorea (MH/MP and MP/MT). It was highly variable, ranging from 0.019 between Moorea-MP and Moorea-MT to 0.392 between Raiatea-RT and Tahiti-TP. When each MLG was considered only once, a different pattern was obtained. Taha'a-TaF showed significant  $F_{ST}$  in comparison with two Moorea (MH/MT), two Raiatea (RA/RU) and two Tahiti (TP/TT) samples. Another pair of samples exhibited a significant  $F_{ST}$ : Raiatea-RT/Tahiti-TP.  $F_{ST}$  ranged between 0.000 (MH/MT) and 0.120 for (TaF/RU).

Correspondence analyses (CA) were performed in order to illustrate the genetic differences among populations in a multidimensional space (Fig. 3). The three populations from Taha'a were largely dispersed along both axes 1 and 2 (Fig. 3a) and axes 1 and 3 (Fig. 3b), indicating a higher genetic distance compared to populations from the three other islands. Genetic distance among the Raiatea populations was also much higher than the variability among Moorea or Tahiti populations, for which populations were grouped together in the CA plots (Fig. 3).

## Discussion

#### Asexual reproduction and clonality levels

Negative  $F_{IS}$  values were recorded at each locus in 10 of the 12 populations of *Pocillopora damicornis* analysed, indicating an excess of heterozygosity compared to the HWE expectation. Such an excess of heterozygotes has previously been interpreted as an effect of clonality (De Meeus and Balloux 2005; Stoeckel et al. 2006): if heterozygotes have better fitness (size, clonal growth rate, survival, etc.), then their clonal propagation will result in their dominance in natural populations. In line with this interpretation,  $F_{IS}$  values no longer differed from HWE once the repeated MLGs were excluded from the analysis, confirming that these departures from HWE were probably explained by the frequent occurrence of asexual reproduction and the higher heterozygosity level of highly represented (and therefore probably more widespread) MLGs. Notwithstanding this, the low number of MLGs retained in some populations, with  $p_{sex}(F_{IS})$  values meaning that it was not possible to determine the clonal identity of all of them, may not provide enough statistical power to detect departures from HWE among sexually generated colonies. The occurrence of several significantly positive  $F_{IS}$  values, indicating a deficit of heterozygotes for some populations and loci, may result from a combination of several other factors such as null alleles and inbreeding (Gaffney et al. 1990; Ayre et al. 1997), or the occurrence of micro-scale Wahlund effects due to genetic patchiness (i.e. the admixture of several differentiated cohorts; Johnson and Black 1984; David et al. 1997; Hedgecock et al. 2007; ArnaudHaond et al. 2008; Selkoe et al. 2010) potentially accentuated by clonal growth after settlement (Becheler et al. 2010).

Restricted levels of genotypic diversity were observed in our study despite a collection strategy designed to minimize the sampling of replicates of the same clonal lineage. Although the set of markers we used showed a limited power for ascertaining the clonal membership of some of the MLGs, the satisfying level of discrimination

obtained for many other MLGs suggests that the low level of genotypic diversity detected here also reflects an important role of asexual reproduction in the effective recruitment of colonies and the maintenance of populations of *P. damicornis* from the Society Islands. These outcomes are comparable with those obtained for *P. damicornis* sensu lato in south-western Australia (Stoddart 1984a, b; Whitaker 2006), East Africa (Souter et al. 2009), Hawai'i (Gorospe and Karl 2013) and in some islands of the Ryukyu Archipelago (Adjeroud and Tsuchiya 1999). However, they contrast with the predominance of sexual reproduction and high levels of genotypic richness observed in populations of *P. damicornis* from the Great Barrier Reef (Benzie et al. 1995; Ayre et al. 1997), West Papua New-Guinea, Indonesia (Starger et al. 2008), and the Tropical Eastern Pacific (Combosch and Vollmer 2011). These outcomes highlight the marked geographic variability in the extent of asexual reproduction and clonality levels in *P. damicornis*.

The prevalence of asexual reproduction sensu lato (i.e. including fragmentation) and the resulting clonality levels that we recorded for *P. damicornis* in the Society Islands ( $0.097 < \text{clonal richness } R < 0.600$ ) also contrast with the higher levels of genotypic richness found in the same region for *P. meandrina* ( $0.847 < R < 1.000$ ; Magalon et al. 2005). These differences could be partly explained by the respective reproductive modes of these corals and the environmental conditions of their preferred habitats. In French Polynesia, the brooding *P. damicornis* relies more on asexual reproduction than does the broadcast spawning *P. meandrina* (Magalon et al. 2005). Given its shape, clonality in *P. damicornis* may be enhanced by a higher level of fragmentation, even if production of asexual (parthenogenetic) larvae could also explain such a high level of clonality. In the Society Islands, *P. damicornis* is mainly restricted to back reef habitats, with a preference for fringing reefs where it forms abundant populations of delicately branching colonies. This skeletal fragility, in association with the chronic occurrence of perturbations such as trampling, gleaning, fishing and anchoring, probably results in high fragmentation rates associated with sizeable survival of some fragments, possibly leading to high levels of clonality within local populations (Pinzón et al. 2012). The frequent occurrence of *P. damicornis* fragments at our sampling sites (M. Adjeroud, pers. obs.) reinforces the hypothesis of fragmentation as an important mechanism underlying the clonal spread and persistence of local populations. In contrast, *P. meandrina* is mainly found on outer reef slopes, where it forms robust branching colonies. Fragmentation of these colonies is probably less common, occurring mainly during major storms. In any case, the chronically high hydrodynamic forces resulting from the currents, swell and wave actions in this habitat probably do not allow the successful settlement and survival of coral fragments of *P. meandrina*, a combination of factors that may explain the contrasting level of clonal richness with its sister species *P. damicornis*.

#### Genetic differentiation and gene flow

A significant genetic differentiation was detected at the regional scale among populations of the four islands of the Society Archipelago, and at the island scale among the three populations of each island. Some populations sampled on Raiatea and its sister island Taha'a (these two islands share the same lagoon) were clearly differentiated from all other populations. While it is difficult to discern between the various hypotheses that may explain this pattern, the observed genetic differentiation could partly result from two consecutive cyclones that have affected Raiatea, Taha'a and Bora Bora in October (Martin) and November (Osea) 1997 (Adjeroud et al. 2005). The post-disturbance recolonization by coral fragments and by asexually generated propagules from surviving colonies, together with the sexual propagules from nearby, less-disturbed coral reefs, may explain the high levels of clonality within these populations and their genetic distance compared to other populations in the Society Islands. This mechanism has been suggested for several coral species from other Indo-Pacific coral reefs (Hunter 1993; Sherman et al. 2006; Underwood et al. 2007; Souter et al. 2010; Starger et al. 2010).



The regional-scale genetic differentiation found for *P. damicornis* is quite different from the panmixia observed in the Society Islands for the broadcast spawner *P. meandrina* (Magalon et al. 2005). The population structure of *P. damicornis* appears to be largely shaped by the sizeable clonality observed within most of the studied populations. The values of  $F_{ST}$  obtained when each multilocus genotype (MLG) was considered only once (e.g. restricting our analysis to colonies of putative sexual origin) reveal that the pattern observed in the Society Islands for *P. damicornis* is not far from panmixia, although for some samples the low number of genotypes may not offer sufficient statistical power to detect a moderate departure from panmixia. As all samples were differentiated based on the entire data sets, and the reduced sample was based on a potentially incomplete set of genetically distinct individuals, it is likely that  $F_{ST}$  estimates based on the reduced data set would lack statistical power and underestimate some of the genetic differentiation occurring at the local scale among the French Polynesian locations.

In conclusion, this study provides further evidence of marked variability in the reproductive strategies and genetic structure of *P. damicornis* throughout its geographic range. Our results clearly demonstrate that asexual reproduction plays a primary role in the maintenance of local populations of *P. damicornis* in the Society Islands and results in a relatively high level of clonality within populations and significant genetic differentiation among populations. However, when considering only colonies of distinct sexual origin, our results suggest a potentially substantial gene flow among some populations from the Society Islands. These outcomes are in accordance with previous models of dispersal and population maintenance in species with mixed modes of reproduction (Williams 1975; Bell 1982), which postulate that sexually derived larvae provide long-distance colonists and gene flow between distant populations, whereas asexual reproduction plays a primary role in the maintenance of local populations (Ayre and Willis 1988; Adjeroud and Tsuchiya 1999). However, our results (sharing of MLGs by distant populations) suggest that regional-scale dispersal of asexually produced larvae may also occur, as proposed by Schmidt-Roach et al. (2013) on the Great Barrier Reef. This contrasts with the lower level of clonality previously found for the congeneric species *P. meandrina* at the same study area; hence, our results clearly underline the importance of life history traits and ecology in shaping the genetic structure of coral populations. However, our hypothesis that *P. damicornis* should exhibit a lower genetic connectivity than *P. meandrina* was not clearly supported by our results. This suggests that, at the spatial scales examined here, high level of clonality within populations is compatible with high genetic connectivity among distant populations.

## Acknowledgments

Some samples were collected with the help of Pauline Bosserelle from the CRIOBE USR 3278 CNRS-EPHE. The samples were genotyped at the IFR 119 "Montpellier Environnement Biodiversité". We would particularly like to thank Marie-Thérèse Augé, Érick Desmarais, Frédérique Cerqueira and Nicolas Bierre for their help in data acquisition and analysis. This paper is a contribution to the ANR "ADACNI" (ANR-12-ADAP-0016).

## References


- Adjeroud M, Tsuchiya M (1999) Genetic variation and clonal structure in the scleractinian coral *Pocillopora damicornis* in the Ryukyu Archipelago, southern Japan. *Mar Biol* 134:753–760
- Adjeroud M, Chancerelle Y, Schrimm M, Perez T, Lecchini D, Galzin R, Salvat B (2005) Detecting the effects of natural disturbances on coral assemblages in French Polynesia: a decade survey at multiple scales. *Aquat Living Resour* 18:111–123
- Adjeroud M, Michonneau F, Edmunds PJ, Chancerelle Y, Penin L, Vidal-Dupiol J, Salvat B, Galzin R (2009) Recurrent disturbances, recovery trajectories, and resilience of coral assemblages on a south Central Pacific reef. *Coral Reefs* 28:775–780
- Almany GR, Connolly SR, Heath DD, Hogan JD, Jones GP, McCook LJ, Mills M, Pressey RL, Williamson DH (2009) Connectivity, biodiversity conservation and the design of marine reserve networks for coral reefs. *Coral Reefs* 28:339–351
- Arnaud-Haond S, Belkhir K (2007) GENCLONE: a computer program to analyse genotypic data, test for clonality and describe spatial clonal organization. *Mol Ecol Notes* 7:15–17
- Arnaud-Haond S, Duarte CM, Alberto F, Serrão EA (2007) Standardizing methods to address clonality in population studies. *Mol Ecol* 16:5115–5139
- Arnaud-Haond S, Vonau V, Rouxel C, Bonhomme F, Prou J, Goyard E, Boudry P (2008) Genetic structure at different spatial scales in the pearl oyster (*Pinctada margaritifera cumingii*) in French Polynesian lagoons: beware of sampling strategy and genetic patchiness. *Mar Biol* 155:147–157
- Ayre DJ, Hughes TP (2000) Genotypic diversity and gene flow in brooding and spawning corals along the Great Barrier Reef, Australia. *Evolution* 54:1590–1605
- Ayre DJ, Miller KJ (2004) Where do clonal coral larvae go? Adult genotypic diversity conflicts with reproductive effort in the brooding coral *Pocillopora damicornis*. *Mar Ecol Prog Ser* 277:95–105
- Ayre DJ, Willis BL (1988) Population structure in the coral *Pavona cactus*: clonal genotypes show little phenotypic plasticity. *Mar Biol* 99:495–505
- Ayre DJ, Hughes TP, Standish RJ (1997) Genetic differentiation, reproductive mode, and gene flow in the brooding coral *Pocillopora damicornis* along the Great Barrier Reef, Australia. *Mar Ecol Prog Ser* 159:175–187
- Becheler R, Diekmann O, Hily C, Moalic Y, Arnaud-Haond S (2010) The concept of population in clonal organisms: mosaics of temporally colonized patches are forming highly diverse meadows of *Zostera marina* in Brittany. *Mol Ecol* 19:2394–2407
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (1996) Genetix 4.05, Logiciel sous Windows pour la Génétique des Populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II, Montpellier, France
- Bell G (1982) The masterpiece of nature: the evolution and genetics of sexuality. Croon Helm, London
- Bellwood DR, Hughes TP, Folke C, Nyström M (2004) Confronting the coral reef crisis. *Nature* 429:827–833
- Bellwood DR, Hoey AS, Ackerman JL, Depczynski M (2006) Coral bleaching, reef fish community phase shifts and the resilience of coral reefs. *Global Chang Biol* 12:1587–1594
- Benzie JAH, Haskell A, Lehman H (1995) Variation in the genetic composition of coral (*Pocillopora damicornis* and *Acropora palifera*) populations from different reef habitats. *Mar Biol* 121:731–739
- Botsford LW, White JW, Coffroth MA, Paris CB, Planes S, Shearer TL, Thorrold SR, Jones GP (2009) Connectivity and resilience of coral reef metapopulations in marine protected areas: matching empirical efforts to predictive needs. *Coral Reefs* 28:327–337
- Carlson DB (1999) The evolution of mating systems in tropical reef corals. *Trends Ecol Evol* 14:491–495

- Combosch DJ, Vollmer SV (2011) Population genetics of an ecosystem-defining reef coral *Pocillopora damicornis* in the Tropical Eastern Pacific. *PLoS One* 6:e21200
- David P, Berthou P, Noel P, Jarne P (1997) Patchy recruitment patterns in marine invertebrates: a spatial test of the density-dependent hypothesis in the bivalve *Spisula ovalis*. *Oecologia* 111: 331–340
- De Meeus T, Balloux F (2005) F-statistics of clonal diploids structured in numerous demes. *Mol Ecol* 14:2695–2702
- Doyle JJ (1995) The irrelevance of allele tree topologies for species delimitation, and a non-topological alternative. *Syst Bot* 20:574–588
- Duarte CM (2007) Marine ecology warms up to theory. *Trends Ecol Evol* 22:332–334
- Elmhirst T, Connolly SR, Hughes TP (2009) Connectivity, regime shifts and resilience of coral reefs. *Coral Reefs* 28:949–957
- Faurby S, Barber PH (2012) Theoretical limits to the correlation between pelagic larval duration and population genetic structure. *Mol Ecol* 21:3419–3432
- Figueiredo J, Baird A, Cohen M, Flot J-F, Kamiki T, Meziane T, Tsuchiya M, Yamasaki H (2012) Ontogenetic change in the lipid and fatty acid composition of scleractinian coral larvae. *Coral Reefs* 31:613–619
- Flot J-F (2007a) Towards a molecular taxonomy of corals of the genus *Pocillopora*. PhD Thesis, Muséum National d'Histoire Naturelle, Paris
- Flot J-F (2007b) Champuru 1.0: a computer software for unraveling mixtures of two DNA sequences of unequal lengths. *Mol Ecol Notes* 7:974–977
- Flot J-F (2010) SeqPHASE: a web tool for interconverting PHASE input/output files and FASTA sequence alignments. *Mol Ecol Resour* 10:162–166
- Flot J-F, Tillier S (2006) Molecular phylogeny and systematics of the scleractinian coral genus *Pocillopora* in Hawaii. In: *Proceedings of the 10th International Coral Reef Symposium*, vol 1, p 24–29
- Flot J-F, Tillier A, Samadi S, Tillier S (2006) Phase determination from direct sequencing of length-variable DNA regions. *Mol Ecol Notes* 6:627–630
- Flot J-F, Magalon H, Cruaud C, Couloux A, Tillier S (2008) Patterns of genetic structure among Hawaiian corals of the genus *Pocillopora* yield clusters of individuals that are compatible with morphology. *C R Biol* 331:239–247
- Flot J-F, Couloux A, Tillier S (2010) Haplowebs as a graphical tool for delimiting species: a revival of Doyle's "field for recombination" approach and its application to the coral genus *Pocillopora* in Clipperton. *BMC Evol Biol* 10:372
- Fukami H, Budd AF, Levitan DR, Jara J, Kersanach R, Knowlton N (2004) Geographic differences in species boundaries among members of the *Montastraea annularis* complex based on molecular and morphological markers. *Evolution* 58:324–337
- Gaffney PM, Scott TM, Koehn RK, Diehl WJ (1990) Interrelationships of heterozygosity, growth rate and heterozygote deficiencies in the coot clam, *Mulina lateralis*. *Genetics* 124:687–699
- Gardner TA, Côté IM, Gill JA, Grant A, Watkinson AR (2003) Long-term region-wide declines in Caribbean corals. *Science* 301:958–960
- Glynn PW, Gassman NJ, Eakin CM, Cortes J, Smith DB, Guzman HM (1991) Reef coral reproduction in the eastern Pacific: Costa Rica, Panama, and Galapagos Islands (Ecuador). I. Pocilloporidae. *Mar Biol* 109:355–368
- Gorospe KD, Karl SA (2013) Genetic relatedness does not retain spatial pattern across multiple spatial scales: dispersal and colonization in the coral, *Pocillopora damicornis*. *Mol Ecol*. doi:10.1111/mec.12335
- Harii S, Yamamoto M, Hoegh-Guldberg O (2010) The relative contribution of dinoflagellate photosynthesis and stored lipids to the survivorship of symbiotic larvae of the reef-building corals. *Mar Biol* 157:1215–1224
- Harrison PL (2011) Sexual reproduction of scleractinian corals. In: Dubinsky Z, Stambler N (eds) *Coral reefs: an ecosystem in transition*, Part 3. Springer, Berlin, pp 59–85


- Hedgecock D, Barber PH, Edmands S (2007) Genetic approaches to measuring connectivity. *Oceanography* 20:70–79
- Heyward AJ, Negri AP (2012) Turbulence, cleavage, and the naked embryo: a case for coral clones. *Science* 335:1064
- Hoegh-Guldberg O (2009) Climate change and coral reefs: Trojan horse or false prophecy? *Coral Reefs* 28:569–575
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatzioi ME (2007) Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737–1742
- Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nyström M, Palumbi SR, Pandolfi JM, Rosen B, Roughgarden J (2003) Climate change, human impacts, and the resilience of coral reefs. *Science* 301:929–933
- Hunter CL (1993) Genotypic variation and clonal structure in coral populations with different disturbance histories. *Evolution* 47:1213–1228
- Johnson MS, Black R (1984) Pattern beneath the chaos: the effect of recruitment on genetic patchiness in an intertidal limpet. *Evolution* 38:1371–1383
- Jones GP, Almany GR, Russ GR, Sale PF, Steneck RS, van Oppen MJH, Willis BL (2009) Larval retention and connectivity among populations of corals and reef fishes: history, advances and challenges. *Coral Reefs* 28:307–325
- Lesser MP (2004) Experimental biology of coral reef ecosystems. *J Exp Mar Biol Ecol* 300:217–252
- Magalon H, Samadi S, Richard M, Adjerdou M, Veuille M (2004) Development of coral and zooxanthella-specific microsatellites in three species of *Pocillopora* (Cnidaria, Scleractinia) from French Polynesia. *Mol Ecol Notes* 4:206–208
- Magalon H, Adjerdou M, Veuille M (2005) Patterns of genetic variation do not correlate with geographical distance in the reef-building coral *Pocillopora meandrina* in the South Pacific. *Mol Ecol* 14:1861–1868
- McManus JW, Polsenberg JF (2004) Coral-algal phase shifts on coral reefs: ecological and environmental aspects. *Prog Oceanogr* 60:263–279
- Miller KJ, Ayre DJ (2008) Protection of genetic diversity and maintenance of connectivity among reef corals within marine protected areas. *Conserv Biol* 22:1245–1254
- Moberg F, Folke C (1999) Ecological goods and services of coral reef ecosystems. *Ecol Econ* 29:215–233
- Mora C, Andréfouët S, Costello MJ, Kranenburg C, Rollo A, Veron J, Gaston KJ, Myers RA (2006) Coral reefs and the global network of marine protected areas. *Science* 312:1750–1751
- Nishikawa A, Katoh M, Sakai K (2003) Larval settlement rates and gene flow of broadcast-spawning (*Acropora tenuis*) and planula-brooding (*Stylophora pistillata*) corals. *Mar Ecol Prog Ser* 256:87–97
- Nozawa Y, Harrison PL (2008) Temporal patterns of larval settlement and survivorship of two broadcast-spawning acroporid corals. *Mar Biol* 155:347–351
- O'Connor MI, Bruno JF, Gaines SD, Halpern BS, Lester SE, Kinlan BP, Weiss JM (2007) Temperature control of larval dispersal and the implications for marine ecology, evolution, and conservation. *Proc Natl Acad Sci USA* 104:1266–1271
- Palumbi SR (1994) Genetic divergence, reproductive isolation, and marine speciation. *Annu Rev Ecol Syst* 25:547–572
- Pandolfi JM, Bradbury RH, Sala E, Hughes TH, Bjorndal KA, Cooke RG, McArdle D, McClenachan L, Newman MJH, Paredes G, Warner RR, Jackson JBC (2003) Global trajectories of the longterm decline of coral reef ecosystems. *Science* 301:955–958
- Parsons KE (1996) The genetic effects of larval dispersal depend on spatial scale and habitat characteristics. *Mar Biol* 126:403–414

- Paz-García DA, Chávez-Romo HE, Correa-Sandoval F, Reyes-Bonilla H, López-Pérez A, Medina-Rosas P, Hernández-Cortés MP (2012) Genetic connectivity patterns of corals *Pocillopora damicornis* and *Porites panamensis* (Anthozoa: Scleractinia) along the west coast of Mexico. *Pac Sci* 66:43–61
- Pinzón JH, Lajeunesse TC (2011) Species delimitation of common reef corals in the genus *Pocillopora* using nucleotide sequence phylogenies, population genetics and symbiosis ecology. *Mol Ecol* 20:311–325
- Pinzón JH, Reyes-Bonilla H, Baums IB, Lajeunesse TC (2012) Contrasting clonal structure among *Pocillopora* (Scleractinia) communities at two environmentally distinct sites in the Gulf of California. *Coral Reefs* 31:765–777
- Richmond RH (1987) Energetics, competency, and long-distance dispersal of planula larvae of the coral *Pocillopora damicornis*. *Mar Biol* 93:527–533
- Richmond RH, Jokiel PL (1984) Lunar periodicity in larva release in the reef coral *Pocillopora damicornis* at Enewetak and Hawaii. *Bull Mar Sci* 34:280–287
- Riginos C, Douglas KE, Jin Y, Shanahan DF, Trembl EA (2011) Effects of geography and life history traits on genetic differentiation in benthic marine fishes. *Ecography* 34:566–575
- Sale PF, Cowen RK, Danilowicz BS, Jones GP, Kritzer JP, Lindeman KC, Planes S, Polunin NVC, Russ GR, Sadovy YJ, Steneck RS (2005) Critical science gaps impede use of no-take fishery reserves. *Trends Ecol Evol* 20:74–80
- Schmidt-Roach S, Miller KJ, Woolsey E, Gerlach G, Baird AH (2012) Broadcast spawning by *Pocillopora* species on the Great Barrier Reef. *PLoS One* 7(12):e50847
- Schmidt-Roach S, Lundgren P, Miller K, Gerlach G, Noreen A, Andreakis N (2013) Assessing hidden species diversity in the coral *Pocillopora damicornis* from Eastern Australia. *Coral Reefs* 32:161–172
- Selkoe KA, Toonen RJ (2011) Marine connectivity: a new look at pelagic larval duration and genetic metrics of dispersal. *Mar Ecol Prog Ser* 436:291–305
- Selkoe KA, Watson JR, White C, Ben Horin T, Iachhei M, Mitarai S, Siegel DA, Gaines SD, Toonen RJ (2010) Taking the chaos out of genetic patchiness: seascape genetics reveals ecological and oceanographic drivers of genetic patterns in three temperate reef species. *Mol Ecol* 19:708–726
- Sherman CDH, Ayre DJ, Miller KJ (2006) Asexual reproduction does not produce clonal populations of the brooding coral *Pocillopora damicornis* on the Great Barrier Reef, Australia. *Coral Reefs* 25:7–18
- Souter P (2010) Hidden genetic diversity in a key model species of coral. *Mar Biol* 157:875–885
- Souter P, Henriksson O, Olsson N, Grahm M (2009) Patterns of genetic structuring in the coral *Pocillopora damicornis* on reefs in East Africa. *BMC Ecol* 9:19
- Souter P, Willis BL, Bray LK, Caley MJ, Muirhead A, van Oppen MJH (2010) Location and disturbance affect population genetic structure in four coral species of the genus *Acropora* on the Great Barrier Reef. *Mar Ecol Prog Ser* 416:35–45
- Starger CJ, Yeoh SSR, Dai CF, Baker AC, Desalle R (2008) Ten polymorphic STR loci in the cosmopolitan reef coral, *Pocillopora damicornis*. *Mol Ecol Resour* 8:619–621
- Starger CJ, Barber PH, Ambariyanto, Baker AC (2010) The recovery of coral genetic diversity in the Sunda Strait following the 1883 eruption of Krakatau. *Coral Reefs* 29:547–565
- Stephens M, Smith NJ, Donnelly P (2001) A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet* 68:978–989
- Stoddart JA (1983) Asexual production of planulae in the coral *Pocillopora damicornis*. *Mar Biol* 76:279–284
- Stoddart JA (1984a) Genetic structure within populations of the coral *Pocillopora damicornis*. *Mar Biol* 81:19–30
- Stoddart JA (1984b) Genetic differentiation amongst populations of the coral *Pocillopora damicornis* of southwestern Australia. *Coral Reefs* 3:149–156

- Stoeckel S, Grange J, Fernández-Manjarres JF, Bilger I, Frascaria-Lacoste N, Mariette S (2006) Heterozygote excess in a self-incompatible and partially clonal forest tree species—*Prunus avium* L. *Mol Ecol* 15:2109–2118
- Trembl EA, Roberts JJ, Chao Y, Halpin PN, Possingham HP, Riginos C (2012) Reproductive output and duration of the pelagic larval stage determine seascape-wide connectivity of marine populations. *Integr Comp Biol* 52:525–537
- Underwood JN, Smith LD, van Oppen MJH, Gilmour JP (2007) Multiple scales of genetic connectivity in a brooding coral on isolated reefs following catastrophic bleaching. *Mol Ecol* 16:771–784
- Underwood JN, Smith LD, van Oppen MJH, Gilmour JP (2009) Ecologically relevant dispersal of corals on isolated reefs: implications for managing resilience. *Ecol Appl* 19:18–29
- Veron JEN (2000) Corals of the world. The Australian Institute of Marine Science, Townsville
- Veron JEN, Pichon M (1976) Scleractinia of Eastern Australia Part I. Australian Institute of Marine Science, Canberra
- Weir B, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358–1370
- Whitaker K (2006) Genetic evidence for mixed modes of reproduction in the coral *Pocillopora damicornis* and its effect on population structure. *Mar Ecol Prog Ser* 306:115–124
- Wilkinson C (2008) Status of coral reefs of the world: 2008. Global Coral Reef Monitoring Network and Reef and Rainforest Research Center, Townsville
- Williams GC (1975) Sex and evolution. Princeton University Press, Princeton
- Wright S (1950) Genetical structure of populations. *Nature* 166:247–249




**Fig. 1** Map of the Society Archipelago in French Polynesia showing the location of the four studied islands (a), and of the 12 sampling sites, in Taha'a and Raiatea (b) and in Moorea and Tahiti (c)



**Fig. 2** Haplotype web (haploweb) of ITS2 sequences. Each pie represents one ITS2 type (with its area proportional to the number of individuals in which it was detected, see scale on the *left*), with the following colour code: Tahaa (Ta) in *light blue*, Raiatea (R) in *dark blue*, Moorea (M) in *orange* and Tahiti (T) in *red*. The lengths of the *grey lines* connecting the ITS2 types are proportional to the number of mutations separating them, whereas ITS2 types found co-occurring in heterozygous individuals are connected with *red curves* (the thickness of which is proportional to the number of heterozygotes in which a given pair of ITS2 types co-occur). Since all ITS2 types are connected by heterozygotes, they belong to a single gene pool sensu Doyle (1995), and therefore, all individuals sequenced are conspecific following the criterion of mutual allelic exclusivity (Flot et al. 2010)





**Fig. 3** Genetic distance between populations. Correspondence analysis conducted on the 12 populations sampled on the four islands in the Society Archipelago. Plots of stations along axes 1 and 2 (a) and along axes 1 and 3 (b). The percentage of variance (inertia) of each of the first three axes is given

**Table 1** Indices of genotypic diversity

		<i>N</i>	<i>G</i>	<i>R</i>	<i>Ho</i>	<i>He</i>	<i>F<sub>IS</sub></i>
<i>N</i> sample size, <i>G</i> number of MLGs, <i>R</i> clonal richness, <i>Ho</i> observed heterozygosity ( $\pm$ SD), <i>He</i> expected heterozygosity ( $\pm$ SD); departures from Hardy–Weinberg equilibrium are expressed as <i>F<sub>IS</sub></i> . Negative values indicate an excess of heterozygotes, and positive values a deficit of heterozygotes  * <i>p</i> < 0.1; ** <i>p</i> < 0.01	MH	30	12	0.379	0.473 (0.296)	0.495 (0.128)	0.045
	MP	30	17	0.552	0.547 (0.122)	0.531 (0.082)	−0.030
	MT	31	19	0.600	0.652 (0.143)	0.534 (0.042)	−0.225 **
	RA	31	10	0.300	0.594 (0.205)	0.568 (0.073)	−0.047
	RT	25	5	0.167	0.256 (0.207)	0.253 (0.146)	−0.013
	RU	32	4	0.097	0.663 (0.262)	0.459 (0.121)	−0.453 **
	TaF	42	16	0.366	0.610 (0.311)	0.477 (0.171)	−0.281 **
	TaT	43	6	0.119	0.651 (0.307)	0.492 (0.168)	−0.327 **
	TaV	43	17	0.381	0.465 (0.187)	0.503 (0.152)	0.076
	TP	41	17	0.400	0.556 (0.293)	0.506 (0.185)	−0.100
	TT	40	9	0.205	0.690 (0.275)	0.533 (0.066)	−0.299 **
	TV	39	20	0.500	0.646 (0.078)	0.578 (0.067)	−0.120 *

**Table 2** Genetic differentiation between populations

The estimator is Weir and Cockerham's  $F_{ST}$ . Values above the diagonal were calculated on the complete data set (i.e. colonies of both sexual and asexual origin). Values under the diagonal were calculated on a data set where only one of each MLG had been retained (i.e. colonies of sexual origin)

\*  $p < 0.01$

	MH	MP	MT	RA	RT	RU	TaF	TaT	TaV	TP	TT	TV
MH		0.036	0.041*	0.030*	0.335*	0.082*	0.197*	0.126*	0.078*	0.028*	0.082*	0.081*
MP	0.028		0.019	0.048*	0.209*	0.052*	0.069*	0.050*	0.121*	0.085*	0.111*	0.069*
MT	0.000	0.013		0.038*	0.213*	0.068*	0.122*	0.070*	0.082*	0.077*	0.059*	0.041*
RA	-0.001	0.023	-0.004		0.289*	0.262*	0.189*	0.183*	0.309*	0.049*	0.326*	0.058*
RT	0.112	0.026	0.063	0.079		0.274*	0.201*	0.199*	0.324*	0.392*	0.343*	0.273*
RU	-0.022	-0.007	-0.007	-0.045	0.053		0.145*	0.115*	0.167*	0.139*	0.160*	0.132*
TaF	0.101*	0.033	0.052*	0.080*	0.047	0.120*		0.092*	0.276*	0.206*	0.214*	0.167*
TaT	0.048	0.006	-0.004	-0.007	0.056	0.032	0.006		0.217*	0.157*	0.177*	0.163*
TaV	-0.001	0.018	-0.017	-0.014	0.064	-0.016	0.052	0.004		0.123*	0.074*	0.062*
TP	0.019	0.057	0.017	0.07	0.115*	0.056	0.067*	0.008	0.010		0.059*	0.100*
TT	0.011	0.072	0.020	0.001	0.090	0.038	0.050*	0.050	-0.002	0.010		0.031*
TV	-0.010	0.016	-0.009	-0.012	0.068	-0.016	0.064	0.025	-0.021	0.012	-0.004	