



HAL
open science

Schistosomiasis Haematobium, Corsica, France

Antoine Berry, H el ene Mon e, Xavier Iriart, Gabriel Mouahid, Olivier Abbo, J er ome Boissier, Judith Fillaux, Sophie Cassaing, Sophie Debuisson, Alexis Valentin, et al.

► **To cite this version:**

Antoine Berry, H el ene Mon e, Xavier Iriart, Gabriel Mouahid, Olivier Abbo, et al.. Schistosomiasis Haematobium, Corsica, France. *Emerging Infectious Diseases*, 2014, 20 (9), pp.1595-1597. 10.3201/eid2009.140928 . halsde-01007702

HAL Id: halsde-01007702

<https://hal.science/halsde-01007702>

Submitted on 7 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

A phylogenetic tree constructed by using the entire sequence showed that all PorCoV HKU15 strains from the United States clustered together in 1 clade of the genus *Deltacoronavirus* with HKU15-155 and HKU15-44 (online Technical Appendix Figure, panel A, <http://wwwnc.cdc.gov/EID/article/20/9/14-0756-Techapp1.pdf>). This finding indicates that 1 genotype of PorCoV HKU15 is currently circulating in multiple US states. This result was further supported by phylogenetic trees constructed by using the full-length amino acids of spike and nucleocapsid proteins (online Technical Appendix Figure, panels B, C). Because of limitation of the samples received, only a partial genome sequence was determined for strain MN3092 from Minnesota. However, on the basis of the spike and nucleocapsid protein sequence analyses, it is highly likely that the entire genome of the Minnesota strain is genetically identical to that of the other 8 strains (online Technical Appendix Figure, panels B, C).

In addition to the 9 states reported in this study, Iowa has also had a recent detection of PorCoV HKU15 (3). Thus, PorCoV HKU15 has been detected in 10 of the 50 US states, and those 10 states mainly cluster in the midwestern United States (Figure). As with data collected for porcine epidemic diarrhea outbreaks by the US National Animal Health Laboratory Network (<http://www.nahln.org/default/>), data collected on the geographic location and numbers of PorCoV HKU15 cases is also required to be reported weekly.

Earlier reports support avian coronaviruses as the gene source for *Deltacoronavirus* spp. (1,4). To confirm this, surveillance for PorCoV should be carried out among birds. Moreover, effective control strategies, including vaccine development, should be in place for prevention and control of infections caused by PorCoV HKU15.

Acknowledgments

We acknowledge and appreciate the excellent technical help provided by Jason Herr and Kerri Lawrence.

Leyi Wang,¹ Beverly Byrum,
and Yan Zhang¹

Author affiliation: Ohio Department of Agriculture, Reynoldsburg, Ohio, USA

DOI: <http://dx.doi.org/10.3209/eid2009.140756>

References

1. Woo PC, Lau SK, Lam CS, Lau CC, Tsang AK, Lau JH, et al. Discovery of seven novel mammalian and avian coronaviruses in the genus *Deltacoronavirus* supports bat coronaviruses as the gene source of *Alphacoronavirus* and *Betacoronavirus* and avian coronaviruses as the gene source of *Gammacoronavirus* and *Deltacoronavirus*. *J Virol*. 2012;86:3995–4008. <http://dx.doi.org/10.1128/JVI.06540-11>
2. Wang L, Byrum B, Zhang Y. Detection and genetic characterization of a delta-coronavirus in pigs in the United States. *Emerg Infect Dis*. 2014;
3. Li G, Chen Q, Harmon KM, Yoon KJ, Schwartz KJ, Hoogland MJ, et al. Full-length genome sequence of porcine deltacoronavirus strain USA/IA/2014/8734. *Genome Announc*. 2014;2:e00278-14. <http://dx.doi.org/10.1128/genomeA.00278-14>
4. Chu DK, Leung CY, Gilbert M, Joyner PH, Ng EM, Tse TM, et al. Avian coronavirus in wild aquatic birds. *J Virol*. 2011;85:12815–20. <http://dx.doi.org/10.1128/JVI.05838-11>

Address for correspondence: Yan Zhang, Animal Disease Diagnostic Laboratory, Ohio Department of Agriculture, 8995 East Main St, Bldg. 6, Reynoldsburg, OH 43068, USA; email: y Zhang@agri.ohio.gov

¹These authors were co-principal investigators.

Schistosomiasis Haematobium, Corsica, France

To the Editor: In Europe, urinary schistosomiasis (*I*) has previously been detected only in Portugal, where this focus disappeared during the 1950s (2). However, freshwater snails of the species *Bulinus contortus*, *B. truncatus*, and *Planorbarius metidjensis*, which are recognized intermediate hosts for *Schistosoma haematobium* trematodes, have been found in Portugal (3), Spain (4), and Corsica (5,6). This finding suggested that autochthonous schistosomiasis could re-emerge in southern Europe if these mollusks become infected. We report a probable focus for transmission of schistosomiasis haematobium in Corsica, France.

In March 2014, a 4-year-old girl (index case-patient) from France was referred to the Toulouse University Hospital (Toulouse, France), with gross hematuria. Ultrasonography and cystoscopic examination of the bladder detected a polyp. Examination of the polyp for parasites identified bodies that were consistent with schistosome eggs. Parasitologic examination of urine confirmed schistosomiasis by detecting viable *S. haematobium* eggs.

The parents of the girl (family A) did not report any stay or travel in an area to which urinary schistosomiasis was endemic; they reported summer holidays only in Mallorca in the Balearic Islands (Spain) and Corsica. However, her father reported that since 2012, he had experienced gross hematuria that had been evaluated by standard urologic investigations but not by cystoscopy; no etiology was determined. Parasitologic urinalysis in our hospital department showed numerous viable *S. haematobium* eggs in the father's urine.

The parents of the index case-patient also reported that an 8-year-old boy in a friend's family (family B), who shared summer vacations with

**EMERGING
INFECTIOUS DISEASES**
On the web
www.cdc.gov/eid

them had exhibited gross hematuria since February 2013. A third family (family C) was also investigated because they also spent holidays in Corsica with families A and B. Families B and C had also spent a summer in Mallorca, but they denied any contact with freshwater. Of 11 French native-born members of the 3 families, 6 had positive results for *S. haematobium* by urine examination. All case-patients had specific positive immunodiagnostic results by an ELISA that used *S. mansoni* extracts and by indirect hemagglutination. In addition, 2 family members who had a negative result by urine examination had a positive serologic result.

Spending summer vacations in the same village in Corsica (Sainte-Lucie de Porto-Vecchio), where members of the 3 families had bathed at least once per holiday period in the Cavu River, was the epidemiologically prominent feature that linked these persons. Families A and C were in Sainte-Lucie de Porto-Vecchio in August 2011, and families A, B, and C were in the same location in August 2013.

During these investigations, we were contacted by the Department of Tropical Medicine, Dusseldorf University Hospital (Dusseldorf, Germany), because a 10-year-old boy and his father had been given diagnoses of schistosomiasis haematobium on the basis of positive urinalysis results for *S. haematobium* eggs. Two other members of this family (5 persons) had a positive immunodiagnostic result. Locations of previous vacations for this family outside Germany included Spain (not the Balearic Islands) and Corsica, where they bathed frequently in the Cavu River. These epidemiologic findings provide strong circumstantial evidence supporting the presence of a previously unrecognized focus of urinary schistosomiasis in Corsica.

We performed molecular analysis of schistosomal miracidia DNA. The second internal transcribed spacer region of the ribosomal gene complex

(7,8) was amplified and sequenced. Viable eggs obtained from the patients in France were those of *S. haematobium*. Additional molecular investigations are being conducted to assess genetic diversity of this isolate from Corsica and the geographic origin of the introduced parasite.

The malacologic situation in Sainte Lucie de Porto-Vecchio was investigated during May 12–19, 2014; three rivers (Figure) were included in the survey. Four sites were sampled in the Cavu River, and *B. truncatus* snails were found in 3 sites that corresponded to bathing areas (site 1: 41°43'53.57"N, 9°17'36.70"E; site 2: 41°43'22.13"N, 9°17'59.87"E; site 3: 41°42'8.40"N, 9°21'5.82"E). Snails were also found in the nearby Tarcu River (site 5) and Osu River (site 6). These findings confirmed previous data for the presence of *B. truncatus*

snails in Corsica (5,6). Water temperature was recorded at 11:00 AM at these 3 sites (range 15°C–16°C). This temperature range is not optimal for the snail intermediate host stage of the parasite life cycle (9,10). Of 148 live snails that were obtained in the Cavu River, none were infected with schistosome cercariae.

Data from the field survey and epidemiologic information for the cases in France and Germany, indicated transmission of schistosomiasis haematobium in the Cavu River in southeastern Corsica in 2011 and 2013. Additional supportive evidence is the fact that the father of the index case-patient had gross hematuria in 2012 and 2013.

Two hypotheses are proposed to account for this situation. The first hypothesis is that the parasite (i.e., schistosome eggs) was transmitted




Figure. Corsica, France, showing malacologic survey sampling sites (oval) in 3 rivers (Tarcu, Cavu, and Osu). *Bulinus truncatus* snails were found at sites 1, 2, 3, 5, and 6.

by an infected person into the Cava River in June or July 2011, when environmental conditions were favorable for snail infection. However, questions arise about survival of infected snails during the winter and their ability to reinfect the area during the following summers in 2012 and 2013. The second hypothesis is that schistosome eggs were spread by infected persons at the beginning of summer and caused a permanent transmission cycle in this focus. This situation would be difficult to control. Additional information should be obtained by a long-term malacologic survey to detect infected mollusks in this region.

**Antoine Berry, Hélène Moné,
Xavier Iriart, Gabriel Mouahid,
Olivier Aboo, Jérôme Boissier,
Judith Fillaux,
Sophie Cassaing,
Cécile Debuissou,
Alexis Valentin,
Guillaume Mita, André Théron,
and Jean-François Magnaval**

Author affiliations: Centre Hospitalier Universitaire, Toulouse, France (A. Berry, X. Iriart, O. Aboo, J. Fillaux, S. Cassaing, C. Debuissou, A. Valentin); Institut National

de la Santé et de la Recherche Médicale, Toulouse (A. Berry, X. Iriart); Centre National de la Recherche Scientifique, Perpignan, France (H. Moné, J. Boissier, G. Mouahid, G. Mita, A. Théron); and Centre National de la Recherche Scientifique, Toulouse (J.-F. Magnaval)

DOI: <http://dx.doi.org/10.3201/eid2009.140928>

References

- Gryseels B. Schistosomiasis. *Infect Dis Clin North Am*. 2012;26:383–97. <http://dx.doi.org/10.1016/j.idc.2012.03.004>
- Fraga de Azevedo J, da Silva JB, Coito AM, Coelho MF, Colaco AT. Portuguese focus of schistosomiasis [in Portuguese]. *Anais do Instituto de Medicina Tropical (Lisb)*. 1948;5:175–222.
- de Azevedo JF. Biological relationships among the different geographical strains of the *Schistosoma haematobium* complex [in French]. *Bulletin de la Société Pathologie Exotique*. 1969;62:348–75.
- Perez-Quintero JC, Bech Taberner M, Huerta Dionisio L. Molluscs of the inland waters of the province of Huelva (SO Spain) [in Spanish]. *Iberus*. 2004;22:19–31.
- Gretillat S. Epidemiology of certain trematode diseases of domestic animals in Corsica (bovine bilharzosis and bovine and ovine distomiasis). Observations conducted during a mission accomplished during the autumn of 1962 [in French]. *Ann Parasitol Hum Comp*. 1963;38:471–81.
- Doby JM, Rault B, Deblock S, Chabaud A. Snails and bilharziasis in Corsica.

Distribution, frequency and biology of *Bulinus truncatus* [in French]. *Ann Parasitol Hum Comp*. 1966;41:337–49.

- Webster BL, Emery AM, Webster JP, Gouvras A, Garba A, Diaw O, et al. Genetic diversity within *Schistosoma haematobium*: DNA barcoding reveals two distinct groups. *PLoS Negl Trop Dis*. 2012;6:e1882. Epub 2012 Oct 25. <http://dx.doi.org/10.1371/journal.pntd0001881>
- Glenn TC, Lance SL, McKee AM, Webster BL, Emery AM, Zerlotini A, et al. Significant variance in genetic diversity among populations of *Schistosoma haematobium* detected using microsatellite DNA loci from a genome-wide database. *Parasit Vectors*. 2013;6:300. <http://dx.doi.org/10.1186/1756-3305-6-300>
- Capron A, Deblock S, Biguet J, Clay A, Adenis L, Vernes A. Contribution to the experimental study of bilharziasis caused by *Schistosoma haematobium* [in French]. *Bull World Health Organ*. 1965;32:755–78.
- Pflüger W, Roushdy MZ, El Emam M. The prepatent period and cercarial production of *Schistosoma haematobium* in *Bulinus truncatus* (Egyptian field strains) at different constant temperatures. *Z Parasitenkd*. 1984;70:95–103. <http://dx.doi.org/10.1007/BF00929579>

Address for correspondence: Antoine Berry, Service de Parasitologie-Mycologie, Institut Fédératif de Biologie, Hôpital Purpan, Centre Hospitalier Universitaire de Toulouse, TSA 40031, 31059 Toulouse CEDEX 9, France; email: berry.a@chu-toulouse.fr

EMERGING INFECTIOUS DISEASES®

Please discontinue my print subscription.

Return:
Fax: 404 639-1954

or mail to:

EID Editor
CDC/NCID/MS D61
1600 Clifton Rd, NE
Atlanta, GA 30333

or email request to:
eideditor@cdc.gov

Number on mailing label:(required) _____

Name: _____

Full mailing address: (BLOCK LETTERS)

Help us do more by
reducing our printing
costs. Read or download
the journal online.

UNSUBSCRIBE

Full text free online at www.cdc.gov/eid