



# Heat shock factor 1 is a potent therapeutic target for enhancing the efficacy of treatments for multiple myeloma with adverse prognosis

Sophie Bustany, Julie Cahu, Géraldine Descamps, Catherine Pellat-Deceunynck, Brigitte Sola

## ► To cite this version:

Sophie Bustany, Julie Cahu, Géraldine Descamps, Catherine Pellat-Deceunynck, Brigitte Sola. Heat shock factor 1 is a potent therapeutic target for enhancing the efficacy of treatments for multiple myeloma with adverse prognosis. *Journal of Hematology and Oncology*, 2015, 8 (1), pp.40. 10.1186/s13045-015-0135-3 . inserm-01237704


**HAL Id: inserm-01237704**

**<https://inserm.hal.science/inserm-01237704>**

Submitted on 3 Dec 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



LETTER TO THE EDITOR

Open Access

# Heat shock factor 1 is a potent therapeutic target for enhancing the efficacy of treatments for multiple myeloma with adverse prognosis

Sophie Bustany<sup>1</sup>, Julie Cahu<sup>1</sup>, Géraldine Descamps<sup>2</sup>, Catherine Pellat-Deceunynck<sup>2</sup> and Brigitte Sola<sup>1\*</sup>

## Abstract

Deregulated expression of heat shock proteins (HSPs) encoding genes is frequent in multiple myeloma. HSPs, which are molecular chaperones involved in protein homeostasis pathways, have emerged recently as promising therapeutic targets. Using human myeloma cell lines and primary myeloma cells belonging to various molecular groups, we tested the efficacy of HSP90, HSP70, and heat shock factor 1 (HSF1) inhibitors alone or associated with current antimyeloma drugs. We report here that KNK-437 (an inhibitor of HSF1) and bortezomib have additive effects on apoptosis induction in cells belonging to groups with bad prognosis.

**Keywords:** Myeloma, Heat shock proteins, HSP inhibitor, Heat shock factor 1, Combined therapy, Lenalidomide, Dexamethasone, Bortezomib, Combination index, Apoptosis

## Findings

Deregulated expression of heat shock proteins (HSPs) and heat shock transcription factor 1 (HSF1) plays a major role in the pathogenesis of multiple myeloma (MM) [1,2]. In turn, several HSP/HSF1 inhibitors are currently undergoing preclinical and/or clinical investigations [3,4].

We used human myeloma cell lines (HMCLs) belonging to several molecular groups [5,6] to analyze HSP expression (Figure 1A). HSP90 and its co-chaperone HSP70 were constitutively expressed in all HMCLs. HSP27 expression was more heterogeneous. Using the Little Rock public database [6], we investigated whether the expression of *HSPB1*, *HSPA4*, and *HSP90AA1* genes varied according to the MM molecular classification. Compared to normal bone marrow plasma cells, *HSP* genes were constantly overexpressed (Figure 1B). *HSPB1* and *HSP90AA1* expressions were higher in the groups with bad prognosis, PR/MS/MF, and *HSPA4* expression in the HY/MF/PR groups. The material and methods used in the study are detailed in Additional file 1.

We studied the sensitivity of HMCLs towards 17-AAG that targets HSP90 or KNK-437 (an inhibitor of HSF1

and, in turn, of both HSP70 and HSP27). HMCLs were constantly sensitive to both inhibitors although heterogeneously responding (Figure 1C, Additional files 2 and 3). This suggests that inhibiting HSPs might potentiate drug treatments for MM patients.


HSPs contribute to MM survival by impairing the mitochondria- and endoplasmic reticulum (ER)-mediated apoptotic pathways [7,8]. In L363 cells (MF group), HSP70 expression decreased following KNK-437 treatment while increased after 17-AAG (Figure 1D). As confirmed by the activation of procaspases 9 and 3 and the cleavage of PARP, a mitochondrial-mediated apoptosis was triggered. The expression of anti-apoptotic BCL2 and MCL1 proteins decreased after KNK-437 treatment. Last, both inhibitors induced a decrease of the procaspase 4, thus favoring an ER stress.

We investigated the capacity of HSP90/HSF1 inhibitors to co-operate with common antimyeloma drugs (bortezomib, dexamethasone, or lenalidomide). We calculated the combination index using the method of Chou [9]. Both inhibitors antagonized lenalidomide effects, suggesting that those associations could be harmful (Additional file 4). The combination of KNK-437 with bortezomib or dexamethasone was highly potent in all cell lines tested but not the association 17-AAG/

\* Correspondence: [brigitte.sola@unicaen.fr](mailto:brigitte.sola@unicaen.fr)

<sup>1</sup>Normandie Univ, UNICAEN, EA4652, Caen, France

Full list of author information is available at the end of the article




dexamethasone. The activation of procaspases 9/3 and the decrease of MCL1 and BCL2 levels were enhanced by the association KNK-437/bortezomib but not the association 17-AAG/bortezomib (Figure 2A). VER-155008, a strict HSP70 inhibitor, combined with bortezomib was no more potent for inducing apoptosis (Figure 2B).

We tested the response of HMCLs co-cultured with human bone marrow stromal cells (HS-5 cells). The percentage of apoptotic cells was enhanced by the co-treatment KNK-437/bortezomib (Figure 2B). This indicates that KNK-437/bortezomib combined therapy could overcome cell adhesion-mediated drug resistance.

We finally analyzed the response of primary cells isolated from four MM or plasma cell leukemia (PCL) patients (Additional file 5) towards KNK-437 and bortezomib after CD138 staining [10]. For patient #3, the fraction of CD138<sup>+</sup> cells decreased in the presence of both drugs, revealing an additive effect in primary cells (Figure 2C). Similar results were obtained for other MM primary samples (Additional file 6).

Our results strongly suggest that HSF1 inhibitors might be promising agents in association with bortezomib-based therapeutic protocols to treat MM patients with adverse prognosis or in relapse.



**Additional files**

**Additional file 1: Material and methods used in the study.** The file contains data on cell line cultures, treatments, and proliferation measurement; primary samples, treatments, and CD138 expression analyses; and gene expression profiling.

**Additional file 2: Response towards HSP90 and HSF1 inhibitors in a panel of MM cell lines.** Cells ( $0.5 \times 10^6$  cells/ml) were seeded for 48 h in 96-well plates and treated with increasing concentrations of 17-AAG (0.3

to 20  $\mu$ M) or KNK-347 (3.1 to 200  $\mu$ M). Cell death was measured by APO2.7 staining and cytometry sorting. LD50 values were defined as the dose that killed 50% of cells. Data represent the mean and SD of three experiments.

**Additional file 3: Inhibitors of HSP90 and HSF1 co-operate differently with antimyeloma drugs in various HMCLs.** Cells were treated for 24 h with HSP inhibitors and then with dexamethasone (Dex), bortezomib (Bort) or lenalidomide (Len) for additional 24 h at the concentrations indicated or with vehicle (DMSO). The absorbance (OD at 490 nm) of

each clone treated with the drug is expressed relative to that of the corresponding clone treated with vehicle (ratio defined as 1 arbitrary unit, AU). For each set of culture conditions, the mean of triplicate ratios is indicated on the graph, together with the SD.

**Additional file 4: Interactions between the drugs analyzed by the combination index (CI) method.** HMCLs were treated for 24 h with HSP90 or HSF1 inhibitors and with dexamethasone, bortezomib, or lenalidomide for additional 24 h at the indicated concentrations. Cell viability was then determined by MTT assay. CIs were calculated according to Chou [9].  $CI < 1$  indicates synergy;  $CI = 1$ , additive effect;  $CI > 1$ , antagonism (gray boxes).

**Additional file 5: Clinical characteristics of MM/PCL patients.** Samples were obtained from patients at diagnosis (D) or relapse (R); patients had multiple myeloma (MM), primary plasma cell leukemia (pPCL), or secondary plasma cell leukemia (sPCL).

**Additional file 6: Additivity of bortezomib and KNK-347 co-treatment on MM primary samples.** Primary cells were obtained from patients with MM or PCL. Purified CD138+ cells were cultured for 24 h and then treated with 5 nM bortezomib alone or in combination with 10  $\mu$ M KNK-437. Cell death was determined as the percentage of CD138+ cells that have lost CD138. The percentage of dead cells directly measured (observed) and the percentage of dead cells calculated for an additive effect (expected) were not significantly different ( $p = 0.37$ , Wilcoxon matched-pairs signed-rank test). This signifies that the effect of the combination of both drugs was indeed additive.

## Abbreviations

Ab: Antibody; BCL2: B-cell lymphoma 2; BMPC: Normal bone marrow plasma cells; CI: Combination index; HMCL: Human myeloma cell line; HSF1: Heat shock transcription factor 1; HSP: Heat shock protein; MCL1: Myeloid cell leukemia 1; MM: Multiple myeloma; PARP: Poly (ADP-ribose) polymerase; VER-155008: 5'-O-[(4-Cyanophenyl)methyl]-8-[[[(3,4-dichlorophenyl)methyl]amino]-adenosine; 17-AAG: 17-Allylamino-17-demethoxy-geldanamycin.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

SB and BS designed the research; SB, JC, GD, CPD, and BS acquired the data; SB, JC, GD, CPD, and BS analyzed the data; and SB and BS wrote the paper. All authors approved the final version of the paper.

## Acknowledgements

The authors thank A Barbaras and Y Zozulya for the technical assistance with cell cultures, the tumorothèque of IRCNA (CHU and ICO, Nantes, France) for providing us with the purified myeloma cells and the technical platforms of flow cytometry (SFR ICORE, Université de Caen Basse-Normandie, PT Cytocell, SFR Bonamy, Nantes). Celgene Corporation (Summit, NJ) provided the lenalidomide. This work was supported by the Fondation de France (Engt n°201200029144) and Comité de la Manche de la Ligue contre le Cancer (to BS). SB was supported by the Ministère de l'Enseignement Supérieur et de la Recherche and JC by the Conseil Régional de Basse-Normandie.

## Author details

<sup>1</sup>Normandie Univ, UNICAEN, EA4652, Caen, France. <sup>2</sup>CRCNA, INSERM U892, CNRS UMR6299, Université de Nantes, Nantes, France.

Received: 25 February 2015 Accepted: 1 April 2015

Published online: 23 April 2015

## References

- Davenport EL, Moore HE, Dunlop AS, Sharp SY, Workman P, Morgan GJ. Heat shock protein inhibition is associated with activation of the unfolded protein response pathway in myeloma plasma cells. *Blood*. 2007;110:2641–9.
- Heimberger T, Andrusis M, Riedel S, Stühmer T, Schraud H, Beilhack A, et al. The heat shock transcription factor 1 as a potential new therapeutic target in multiple myeloma. *Br J Haematol*. 2013;160:465–76.
- Mahindra A, Laubach J, Raju N, Munshi N, Richardson PG, Anderson KC. Latest advances and current challenges in the treatment of multiple myeloma. *Nat Rev Clin Oncol*. 2012;9:135–43.

- Podar K. Novel targets and derived small molecule inhibitors in multiple myeloma. *Curr Cancer Drug Targets*. 2012;12:797–813.
- Moreaux J, Klein B, Bataille R, Descamps G, Maiga S, Hose D, et al. A high-risk signature for patients with multiple myeloma established from the molecular classification of human myeloma cell lines. *Haematologica*. 2011;96:574–82.
- Zhan F, Huang Y, Colla S, Stewart J, Hanamura I, Gupta S, et al. The molecular classification of multiple myeloma. *Blood*. 2006;108:2020–8.
- Mitsiades CS, Mitsiades NS, McMullan CJ, Poulaki V, Kung AL, Davies FE, et al. Antimyeloma activity of heat shock protein-90 inhibition. *Blood*. 2006;107:1092–100.
- Guo F, Rocha K, Bali P, Pranpat M, Fiskus W, Boyapalle S, et al. Abrogation of heat shock protein 70 induction as a strategy to increase antileukemia activity of heat shock protein 90 inhibitor 17-allylamino-demethoxy geldanamycin. *Cancer Res*. 2005;65:10536–44.
- Chou TC. Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. *Pharmacol Rev*. 2006;58:621–81.
- Surget S, Chiron D, Gomez-Bougie P, Descamps G, Ménoret E, Bataille R, et al. Cell death via DR5, but not DR4, is regulated by p53 in myeloma cells. *Cancer Res*. 2012;72:4562–73.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

