



An efficient Bayesian method for inverting physical models on massive planetary data

Benoit Kugler, Florence Forbes, Sylvain Douté

► To cite this version:

Benoit Kugler, Florence Forbes, Sylvain Douté. An efficient Bayesian method for inverting physical models on massive planetary data. EPSC 2020 - Europlanet Science Congress, Sep 2020, Granada (Virtual meeting), Spain. pp.1-4. insu-03705153v1

HAL Id: insu-03705153

<https://hal.science/insu-03705153v1>

Submitted on 28 Sep 2020 (v1), last revised 2 Jul 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AN EFFICIENT BAYESIAN METHOD FOR INVERTING PHYSICAL MODELS ON MASSIVE PLANETARY DATA

Benoit KUGLER¹, Florence FORBES¹ and Sylvain DOUTÉ²

¹ *Univ. Grenoble Alpes, CNRS, Inria, Grenoble INP, LJK, 38000 Grenoble, France
(firstname.lastname@inria.fr)*

² *Univ. Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France (sylvain.doute@univ-grenoble-alpes.fr)*

Introduction

When studying the Martian surface, the composition of the materials is established on the basis of spectral detection, unmixing, and physical modelling using images produced by hyperspectral cameras. Information on the microtexture of surface materials such as grain size, shape, roughness and internal structure can also be used as tracers of geological processes [1]. This information is accessible under certain conditions thanks to hyperspectral image sequences acquired over a site of interest from eleven different angles by the Compact Reconnaissance Imaging Spectrometer for Mars [2] (CRISM@MRO). Similar multi-angular observations can also be realized in the laboratory with spectro-photo-goniometers, on planetary analog materials or real extraterrestrial matter [3]. In both cases, the interpretation of the surface Bidirectional Reflectance Distribution Factor (BRDF) extracted from these observations, in terms of composition and microtexture, is based on the inversion of the Hapke model (see [4] for details), a semi-empirical photometric model that relates physically meaningful parameters \mathbf{x} to the reflectivity \mathbf{y} of a granular material for a given geometry of illumination and viewing.

This work presents an efficient method, based on a learning approach in a Bayesian framework, able to invert the Hapke model on large experimental or remote sensing datasets, as illustrated on a challenging CRISM multi-angular sequence of images.

1 Inversion characteristics

In the CRISM case, the dataset provides both spatial and spectral dimensions. More specifically, each of the $N_{spatial}$ spatial point of the scene is observed at $N_{spectral}$ wavelengths, yielding $N_{spatial} \times N_{spectral}$ \mathbf{y}_{obs} vectors of reflectance values (typically of order $3000 \times 100 = 300000$). This **high number of observations to be inverted** usually makes approaches like Markov Chain Monte-Carlo simulations unacceptably slow. The approach we propose starts with a computer intensive learning phase that builds a statistical model for the Hapke model; it is then applied for all observables \mathbf{y}_{obs} .

Each observable is a vector of D reflectances, one for each geometry of measure. In the case of CRISM data, $D = 11$ is moderate, but in laboratory, D may be one order of magnitude higher. The statistical model we propose has the advantage to be able to handle **high dimensional settings** while remaining computationally tractable.

The Bayesian framework offers a natural solution to **propagate uncertainties** on measures, which are formalized as variance on the prior distribution, while the variance of the posterior distribution assesses the uncertainty induced by the inversion.

The complexity of the forward radiative transfer model often makes our inverse problem ill-posed. In practice, **multiple solutions might be acceptable**. More than a point estimator, our statistical model provides a full posterior density, whose multi-modality can be assessed and interpreted.

2 Bayesian inversion via a learning approach

From a statistical point of view, we formalize our model as couple of random variables (X, Y) , where $Y = F(X) + \varepsilon$. X has a prior distribution on the physical parameters space, F is the forward model and ε is a centered Gaussian noise accounting for measure uncertainties. The main idea is to use a two steps approach: first, approximate this model by a parametric surrogate model, second use the surrogate model to invert the observations.

We chose the family of the so-called Gaussian Locally-Linear Mapping (GLLiM) [5]. Such models are expressive enough to approximate highly non-linear, complex, forward models, while remaining tractable.

The learning step is performed by generating a training dictionary composed of samples from (X, Y) , on which the likelihood is maximized by a standard EM algorithm. Regarding the dimensionality of the problem, the number of parameters is proportional to D , making this model suitable even for high dimensional observables. Moreover, the computational cost of the learning phase does not scale with the number of observations to be inverted.

The second step is performed by computing the posterior distribution of the random variable X given \mathbf{y}_{obs} . The inverse of the surrogate GLLiM has an explicit formulation in the form of a Gaussian Mixture. From this posterior density, the mean and variance are straightforward to compute. To handle a multi-solution scenario, the mixture can be further exploited.

3 Massive inversion of spectro-images

The dataset comes from a multi-angular observation of the South Pole of Mars by CRISM [2]. The targeted scene presents spatially segregated CO₂ ice, H₂O frost, and mineral dust [6]. After fusion and atmospheric correction of eleven hyperspectral images [7], the dataset provides both spatial and spectral dimensions, totaling $N_{obs} = 154650 = 3093 \times 50$ measurements vectors \mathbf{y}_{obs}^n , which makes MCMC approach unacceptably slow. Maps of Hapke's parameter values are generated from the results of our inversion and superposed with transparency onto the full resolution CRISM nadir image, which serves as a geological control background image.

The results are satisfying from the application point of view. The colour composition of Figure 1 reflects the variation of ω at three wavelengths and corresponds well with the spatial distribution of the three previous materials and their known spectral optical properties. The map of the roughness parameter $\bar{\theta}$ averaged over the spectral dimension (Figure 2) is color coded by intervals of values whose spatial variations are correlated with the composition and the structures of the terrains. In general, we find that all predicted parameters preserve some spatial regularity and show meaningful correlations with the composition and the geology.

Conclusion

We propose an efficient statistical and learning method for inverting physical models on massive remote sensing or experimental data. In the case of the Hapke model, the method has proved to produce satisfactory results for CRISM multi-angular acquisitions as illustrated in this paper and for laboratory spectrophotometric measurements as illustrated in Potin et al., this conference.

References

- [1] Fernando, J., Schmidt, F., and Douté, S. (September, 2016) Martian Surface Microtexture from Orbital CRISM Multi-Angular Observations: A New Perspective for the Characterization of the Geological Processes. *Planetary and Space Science*, **128**, 30–51.
- [2] Murchie, S. L., Seelos, F. P., Hash, C. D., Humm, D. C., Malaret, E., McGovern, J. A., Choo, T. H., Seelos, K. D., Buczkowski, D. L., Morgan, M. F., Barnouin-Jha, O. S., Nair, H., Taylor, H. W., Patterson,




Figure 1: Mars South Pole dataset. Parameter ω in synthetic colors coding 3 wavelengths.




Figure 2: Mars South Pole dataset. Parameter $\bar{\theta}$ averaged over spectral dimension.

- G. W., Harvel, C. A., Mustard, J. F., Arvidson, R. E., McGuire, P., Smith, M. D., Wolff, M. J., Titus, T. N., Bibring, J.-P., and Poulet, F. (2009) Compact Reconnaissance Imaging Spectrometer for Mars Investigation and Data Set from the Mars Reconnaissance Orbiter’s Primary Science Phase. *Journal of Geophysical Research: Planets*, **114**(E2).
- [3] Potin, S., Beck, P., Schmitt, B., and Moynier, F. (November, 2019) Some Things Special about NEAs: Geometric and Environmental Effects on the Optical Signatures of Hydration. *Icarus*, **333**, 415–428.
 - [4] Schmidt, F. and Fernando, J. (2015) Realistic Uncertainties on Hapke Model Parameters from Photometric Measurements. *Icarus*, **260**, 73–93 (IF 2,84).
 - [5] Deleforge, A., Forbes, F., and Horaud, R. (September, 2015) High-Dimensional Regression with Gaussian Mixtures and Partially-Latent Response Variables. *Statistics and Computing*, **25**(5), 893–911.
 - [6] Douté, S. and Pilorget, C. (September, 2017) Physical State and Temporal Evolution of Icy Surfaces in the Mars South Pole. *European Planetary Science Congress*, **11**, EPSC2017–491.
 - [7] Ceamanos, X., Douté, S., Fernando, J., Schmidt, F., Pinet, P., and Lyapustin, A. (2013) Surface Reflectance of Mars Observed by CRISM/MRO: 1. Multi-Angle Approach for Retrieval of Surface Reflectance from CRISM Observations (MARS-ReCO). *Journal of Geophysical Research: Planets*, **118**(3), 514–533.