



HAL
open science

THE USE OF THE ECM METHOD AS AN INDICATOR OF HIGH DUST CONCENTRATIONS IN POLAR ICE CORES

C. Hammer

► **To cite this version:**

C. Hammer. THE USE OF THE ECM METHOD AS AN INDICATOR OF HIGH DUST CONCENTRATIONS IN POLAR ICE CORES. *Journal de Physique Colloques*, 1987, 48 (C1), pp.C1-697-C1-697. 10.1051/jphyscol:19871113 . jpa-00226255

HAL Id: jpa-00226255

<https://hal.science/jpa-00226255>

Submitted on 4 Feb 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**THE USE OF THE ECM METHOD AS AN INDICATOR OF HIGH DUST CONCENTRATIONS
IN POLAR ICE CORES**

C.H. HAMMER

*Department of Glaciology, Geophysical Institute, University of
Copenhagen, DK-2200 Copenhagen, Denmark*

Abstract : ECM*, pH and dust concentration data obtained on the Dye 3, Camp Century and Byrd ice cores suggest, that the ECM data may be used to indicate high dust concentrations in polar ice cores i.e. higher than approx. 300 $\mu\text{g}/\text{kg}$ ice. The dust concentration can only be inferred in a semi-quantitative way as the calibration curve shows a high scatter, especially for the internal 300-500 $\mu\text{g}/\text{kg}$. Various data from the 3 cores will be presented in order to illustrate the potential and limitations.

*ECM : Electrical Conductivity Method.