



**HAL**  
open science

## Crossover scaling in surface growth with truncated power-law noise

Jacques Amar, Fereydoon Family

► **To cite this version:**

Jacques Amar, Fereydoon Family. Crossover scaling in surface growth with truncated power-law noise. Journal de Physique I, 1991, 1 (2), pp.175-179. 10.1051/jp1:1991123 . jpa-00246312

**HAL Id: jpa-00246312**

**<https://hal.science/jpa-00246312>**

Submitted on 4 Feb 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Classification  
 Physics Abstracts  
 47.55M — 05.40 — 64.60A

### Short Communication

## Crossover scaling in surface growth with truncated power-law noise

Jacques G. Amar and Fereydoon Family

Department of Physics, Emory University, Atlanta, GA 30322, U.S.A.

(Received 14 November 1990, accepted 21 November 1990)

**Abstract.** — We demonstrate the existence of a simple scaling form which describes the crossover from anomalous to Gaussian exponents as a function of cutoff in the Zhang model of surface growth for  $\mu > 2$ , where  $\mu$  is the exponent characterising the power-law noise distribution. For  $\mu = 2$  we find a novel scaling form with logarithmic corrections. Our results for cutoff scaling at  $\mu = 2$  are supported by the scaling of the saturation growth velocity.

Zhang [1] has recently proposed a model of surface growth in which the noise  $\eta$  has a power-law distribution of the form,

$$P(\eta) \sim 1/\eta^{1+\mu} \quad (1)$$

for large  $\eta$ . From simulations of this model as well as of the related directed polymer model [1, 2] in  $d = 2$ , he obtained anomalously large values for the surface scaling exponents, which were found to depend on  $\mu$ . In addition, he suggested that the presence of power-law noise may explain the results of recent experiments on porous media [3-6] and bacterial growth [7] in  $d = 2$ , for which anomalously high values of the growth exponents have also been reported. Recently, we have carried out extensive simulations [8] of several different surface growth models with power-law noise which strongly support the existence of anomalous exponents. However, there still exists some controversy over whether the observed anomalous exponents represent a new universality class for the Kardar-Parisi-Zhang (KPZ) equation [9]. In order to resolve some of these questions we have conducted a study of the Zhang model [1] with truncated noise.

In our simulations we used the power-law noise distribution given by equation (1), with a cutoff  $\eta_c$  such that  $P(\eta) = 0$  for  $\eta > \eta_c$ . The noise  $\eta$  was determined at each odd (even) site at each odd (even) time step  $t$ , by generating an independent random number  $r$  at each site such that  $\eta_c^{-\mu} < r < 1$  and calculating the quantity  $\eta = r^{-1/\mu}$ . Simulations were performed for a range of values of the cutoff, with  $\eta_c^{-\mu}$  ranging from  $2^{-31}$  to  $10^{-3}$  in powers of 10 for each value of  $\mu$ . The saturation value of the surface width, as measured by the r.m.s. fluctuation of the interface height, was determined for each value of the cutoff  $\eta_c$ , for system sizes  $L = 16-1024$  in powers of 2. Averages were taken over times of the order of several million time steps.

The surface width  $w(L, t)$  on length scale  $L$  at time  $t$  is expected to satisfy the scaling form [10],  $w(L, t) = L^\alpha f(t/L^z)$ , where  $z = \alpha/\beta$  is the dynamic exponent,  $f(x) \sim x^\beta$  for  $x \ll 1$  and  $f(x) \rightarrow \text{const}$  for  $x \gg 1$ , and  $\alpha$  has the value of  $1/2$  [9] for Gaussian noise. In order to scale our results for the saturation width  $w(L, t = \infty, \eta_c)$  as a function of cutoff  $\eta_c$ , we propose the following crossover form for  $\mu > 2$ ,

$$w(L, \infty, \eta_c) \sim L^\alpha f(L^{-x_c} \eta_c) \quad (2)$$

where  $f(u) \sim u^{(\alpha-1/2)/x_c}$  for  $u \ll 1$  and  $f(u) \rightarrow \text{const}$  for  $u \gg 1$  and  $\alpha$  and  $x_c$  depend on  $\mu$ . This is equivalent to the scaling form,

$$w(L, \infty, \eta_c) \sim L^{1/2} \eta_c^{(\alpha-1/2)/x_c} g(L^{x_c} \eta_c^{-1}) \quad (3)$$

with  $g(u) \sim u^{(\alpha-1/2)/x_c}$  for  $u \ll 1$  and  $g(u) \rightarrow \text{const}$  for  $u \gg 1$ . Figure 1 shows scaling plots of the form of (2) for  $\mu = 3-5$ . For  $\alpha$  we have used the values ( $\alpha = 0.75, 0.62, 0.56$ ) obtained from a previous study [8], while the crossover exponents  $x_c$  were obtained from best fits ( $x_c \simeq 0.62, 0.375$  and  $0.2$  respectively). We find good scaling for all three values of  $\mu$ , except at small  $L$  for which the asymptotic region has not yet been reached. We note that with increasing  $\mu$  the crossover scaling exponent  $x_c$  decreases towards zero. However, the quantity  $(\alpha - 1/2)/x_c \simeq 0.33-0.4$  appears to remain approximately constant for  $\mu > 2$ .




Fig. 1. — Scaling plots for  $\mu = 3$  ( $L = 32-1024$ ), 4 and 5 ( $L = 64-1024$ ). Highest curve for  $\mu = 5$  is for  $L = 64$ .

We now consider cutoff scaling for  $\mu = 2$ . This is the borderline for the existence of self-invariant noise, because for  $\mu = 2$  the second moment of the noise is logarithmically divergent. Because of this, the surface is fractal and  $\alpha$  has been assumed to be exactly 1 for  $\mu = 2$  [1, 11]. The inset of figure 2 shows the best scaling of the form (2) obtained with  $\alpha = 1$ , which occurred for  $x_c = 1$ . The scaling is relatively poor even though for  $\mu = 2$  the data for the saturation width

appears to show asymptotic behavior for very small  $L$  [1, 8]. This led us to consider an alternate form with logarithmic corrections at  $\mu = 2$ ,

$$w(L, \infty, \eta_c) \sim L(\ln L)^p F\left([L(\ln L)^{2p}]^{-x_c} \eta_c\right) \quad (4)$$




Fig. 2. — Scaling plot of form of equation (4) for  $\mu = 2$ , with  $p = 0.20$  and  $x_c = 1$ .  $L$  ranges from 16 to 1024 and  $\eta_c$  varies from 31.6 to 46340. Inset shows scaling plot of form of equation (3).

where  $F(u) \sim u^{1/2x_c}$  for  $u \ll 1$  and  $F(u) \rightarrow \text{const}$  for  $u \gg 1$ . Using this scaling form with  $x_c = 1$  and  $p \simeq 0.20-0.25$  we find very good scaling for the entire range of system sizes  $L$ . Thus, there appears to exist a logarithmic correction at  $\mu = 2$ . The existence of such a correction may explain why values of  $\alpha$  slightly higher than 1 ( $\alpha \simeq 1.04$ ) were obtained for several different growth models with power-law noise at  $\mu = 2$  in reference [8].

In order to obtain further support for our results, we also studied the scaling of the saturation velocity  $V(L)$  which is expected to have a finite-size correction of the form [12],

$$V(L) - V(\infty) \sim \langle \lambda |\nabla h|^2 \rangle \sim A(\lambda) L^{2\alpha-2} \quad (5)$$

where  $A(\lambda)$  is proportional to the nonlinearity parameter  $\lambda$  in the KPZ equation. Very good fits of the form of equation (5) were obtained for  $\mu = 3-5$ , using the values of  $\alpha$  previously obtained from the scaling of the interface width. For  $\mu = 2$ , however, the saturation velocity does not approach a constant value as expected for  $\alpha = 1$  and in fact appears to be diverging with  $L$ . We conjecture a logarithmic correction of the form  $V(L) \sim (\ln L)^{2p}$ . Figure 3 shows a plot of  $\ln(V(L))$  versus  $\ln(\ln(L))$ . The slope of the straight-line fit is  $2p \simeq 0.58$ . This supports the existence of a logarithmic correction at  $\mu = 2$ , which is most likely due to the fact that  $\mu = 2$  is the critical value for the transition from non-invariant to self-invariant noise.

Our results for cutoff scaling have implications for experiments as well as for simulations of the Zhang model with large system sizes. The reason is that typically 32-bit or 46-bit integer




Fig. 3. — Plot of  $\ln V(L)$  versus  $\ln \ln L$  showing logarithmic divergence of  $V(L)$  at  $\mu = 2$ . Dashed line fit has slope  $2p = 0.58$ .

random number generators, for which the cutoff  $r_c$  is  $2^{-31}$  ( $4.65 \times 10^{-10}$ ) and  $2^{-45}$  ( $2.8 \times 10^{-14}$ ) respectively, are used. Equation (2) implies that the scaling region for which the surface width is not affected by the cutoff occurs for  $u = \eta_c/L^{x_c} = r_c^{-1/\mu}/L^{x_c} > u_c$  where  $u_c \gg 1$ . For a 32-bit random number generator, using the value  $u_c \simeq 40$  ( $\log(u_c) = 1.6$ , see Fig. 1), we find for  $\mu = 3, 4, 5$  that  $L_c = 257, 88, 21$  respectively. For a 46-bit random number generator for which  $r_c = 2.8 \times 10^{-14}$ , we obtain substantially larger values for  $L_c$  (of the order of  $10^4$ ). However, on the order of  $10^{14}$  random numbers must be generated to sample the full range of the random number generator, while a reasonable estimate of the number of random numbers accessible in current computer simulations is only of the order of  $10^9$ - $10^{11}$ . Thus, the effects of a cutoff should begin to show up for systems of order  $10^3$  even in large-scale simulations. Crossover to Gaussian exponents, however, ( $u < 1$ ) should only occur for much larger system sizes ( $10^5$  or larger).

### Summary.

We have demonstrated the existence of a simple scaling form which describes the crossover to Gaussian exponents as a function of cutoff in the Zhang model of surface growth for  $\mu > 2$ . This crossover form strongly supports the existence of anomalous exponents in this model. In addition, for  $\mu = 3, 4$  and  $5$  we have obtained approximate values for the crossover exponents  $x_c(\mu)$ . With increasing  $\mu$ ,  $x_c$  decreases towards zero while the quantity  $(\alpha - 1/2)/x_c$  remains approximately constant.

For  $\mu = 2$  we find a novel scaling form with logarithmic corrections. Our results for cutoff scaling at  $\mu = 2$  are further supported by the scaling of the saturation growth velocity  $V(L)$  which scales as  $(\ln L)^{2p}$  with  $p \simeq 0.29$ . For  $\mu > 2$ , our results for the finite-size correction to the growth velocity agree with previously obtained estimates for the saturation exponent  $\alpha$  as a function of  $\mu$ , thus verifying the validity of the nonlinear KPZ equation as a description of the Zhang model.

Due to the natural existence of cutoffs, our results may be useful in the analysis of experiments with power-law noise.

### Acknowledgements.

This work was supported by the Office of Naval Research and the Petroleum Research Fund Administered by the American Chemical Society.

### References

- [1] ZHANG Y.C., *J. Phys. France* **51** (1990) 2129.
- [2] MARINI-BETTOLO-MARCONI U. and ZHANG Y.C., to be published.
- [3] RUBIO M.A., EDWARDS C.A., DOUGHERTY A. and GOLLUB J.P., *Phys. Rev. Lett.* **63** (1989) 1685.
- [4] HORVÁTH V.K., FAMILY F. and VICSEK T., *Phys. Rev. Lett.* **65** (1990) 1388.
- [5] RUBIO M.A., DOUGHERTY A. and GOLLUB J.P., *Phys. Rev. Lett.* **65** (1990) 1389.
- [6] HORVÁTH V.K., FAMILY F. and VICSEK T., *J. Phys. A : Math. Gen.* **24** (1991) L-25.
- [7] VICSEK T., CSERZŐ M. and HORVÁTH V.K., *Physica A* **167** (1990) 315.
- [8] AMAR J.G. and FAMILY F., *J. Phys. A : Math. Gen.* **24** (1991) L-79.
- [9] KARDAR M., PARISI G. and ZHANG Y.-C., *Phys. Rev. Lett.* **56** (1990) 889.
- [10] FAMILY F. and VICSEK T., *J. Phys. A : Math. Gen.* **18** (1985) L75.
- [11] ZHANG Y.-C., *Physica A* **170** (1990) 1.
- [12] KRUG J. and MEAKIN P., *J. Phys. A : Math. Gen.* **23** (1990) L987.

Cet article a été imprimé avec le Macro Package "Editions de Physique Avril 1990".