

Toxicity, transfer and depuration of anatoxin-a (cyanobacterial neurotoxin) in medaka fish exposed by single-dose gavage

Simon Colas, Charlotte Duval, Benjamin Marie

▶ To cite this version:

Simon Colas, Charlotte Duval, Benjamin Marie. Toxicity, transfer and depuration of anatoxin-a (cyanobacterial neurotoxin) in medaka fish exposed by single-dose gavage. Aquatic Toxicology, 2020, 222, pp.105422. 10.1016/j.aquatox.2020.105422. mnhn-02465304v1

HAL Id: mnhn-02465304 https://hal.science/mnhn-02465304v1

Submitted on 15 Dec 2020 (v1), last revised 3 Feb 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Toxicity, transfer and depuration of anatoxin-a (cyanobacterial neurotoxin) in medaka fish exposed by single-dose gavage

3

4 Simon COLAS¹, Charlotte DUVAL¹ and Benjamin MARIE¹

¹UMR 7245 CNRS/MNHN "Molécules de Communications et Adaptations des Micro-organismes, Muséum
 national d'Histoire naturelle, Paris – France

7 8

9 Abstract

10 The proliferations of cyanobacteria are increasingly prevalent in warm and nutrient-enriched 11 waters and occur in many rivers and water bodies due especially to eutrophication. The aim of 12 this work is to study in female medaka fish the toxicity, the transfer and the depuration of the 13 anatoxin-a, a neurotoxin produced by benthic cyanobacterial biofilms. This work will provide 14 answers regarding acute toxicity induced by single gavage by anatoxin-a and to the risks of 15 exposure by ingestion of contaminated fish flesh, considering that data on these aspects remain 16 particularly limited.

17

The oral LD₅₀ of a single dose of (\pm)-anatoxin-a was determined at 11.50 µg.g⁻¹. First of all, lethal dose (100% from 20 µg.g⁻¹) provokes rapid respiratory paralysis (in 1-2 min) of the fish inducing the death by asphyxia. Noticeably, no death nor apparent neurotoxicologic effect occurred during the experimentation period for the 45 fish exposed to a single sub-acute dose of (\pm)-anatoxin-a corresponding to the no observable adverse effect level (NOAEL = 6.67 µg.g⁻¹). Subsequently, the toxico-kinetics of the (\pm)-anatoxin-a was observed in the guts, the livers and the muscles of female medaka fish for 10 days.

25 In parallel, a protocol for extraction of anatoxin-a has been optimized beforehand by testing 3 different solvents on several matrices, the extraction with 75% methanol + 0.1% formic acid 26 27 appearing to be the most efficient. Anatoxin-a was quantified by high-resolution qTOF mass 28 spectrometry coupled upstream to a UHPLC chromatographic chain. The toxin could not be 29 detected in the liver after 12 h, and in the gut and muscle after 3 days. The mean clearance rates 30 of (±)-anatoxin-a calculated after 12 h are above 58%, 100% and 90% for the guts, the livers 31 and the muscles, respectively. Non-targeted metabolomics investigations performed on the fish 32 liver indicates that the single sub-acute exposure by gavage induces noticeable metabolome 33 dysregulations, including important phospholipid decreases, with an organism recovery period 34 of above 12-24h. Overall, the medaka fish do not appear to accumulate (\pm) -anatoxin-a and to 35 largely recover after 24h following a single sub-acute oral liquid exposure at the NOAEL. 36

37 Keywords: cyanobacteria; neurotoxin; fish; toxicology; gavage; toxico-kinetics.

39 Introduction

40 Worldwide proliferation of cyanobacterial blooms constitutes a serious environmental and 41 economic problem that menaces wildlife and human health. Moreover, many cyanobacteria are able to produce potent hepatotoxins such as microcystin, cylindrospermopsin and nodularin, 42 43 and/or neurotoxins such as anatoxins-a, homo-anatoxin-a, anatoxin-a(s) and saxitoxin (Sivonen 44 & Jones, 1999). Harmful cyanobacterial blooms in lakes have been described for decades; in 45 rivers, however, first reports of animal deaths from toxic benthic cyanobacteria in many regions 46 occurred only in the last 20 years (Wood et al., 2007). Since toxic benthic cyanobacteria in 47 rivers have been documented in many countries and benthic taxa have been found to produce a 48 large spectrum of those cyanotoxins, though most animal fatalities reported in relation to 49 benthic cyanobacteria were associated with the presence of anatoxin-a (ANTX) and/or 50 homoanatoxin-a (HANTX) (for review Quiblier et al., 2013).

51 Anatoxin-a and homoanatoxin-a are potent neurotoxins produced by some planktonic and 52 benthic strains of the genera Anabaena, Oscillatoria, Aphanizomenon and Cylindrospermum 53 (Bouma-Gerson et al., 2018). Indeed, numerous fatal intoxications of animals, by anatoxin-a 54 and homoanatoxin-a, have been reported all over the world (Gugger et al., 2005; Puschner et al., 2010; Wood et al., 2007). These alkaloids bind tightly to the nicotinic acetylcholine 55 56 receptor, in the sub-nanomolar range, and thus provoke the death of animals almost 57 immediately after ingestion (Wonnacott and Gallagher, 2006). Anatoxin-a, as a competitive 58 agonist of acetylcholine can bind it specific membrane receptors, notably at neuromuscular 59 junctions, affecting signal transmission between neurons and muscles, causing muscle cell 60 overstimulation (Carmichael, 1994; Aráoz et al. 2010). Acute effects in vertebrates include a 61 rapid loss of coordination, a decreased of locomotor activity, a paralysis of the peripheral 62 skeletal and respiratory muscles, causing symptoms such as loss of coordination, twitching, irregular breathing, tremors, altered gait and convulsions before death by acute asphyxia 63 64 induced by respiratory arrest (Dittmann and Wiegand, 2006).

65

66 It is maybe because anatoxin-a is very unstable and labile in the water (Stevens and Krieger, 1991) and because no chronic effects have been described in mammals so far (Fawell et al., 67 1999), that this toxin has been considered of less environmental concern comparing to other 68 69 cvanotoxins. Despite the acute neurotoxic effects of anatoxin-a, the consequence of the 70 proliferation of anatoxins-a producing cyanobacterial biofilms on ecosystem health and aquatic 71 organisms remains largely unknown. Despite, toxic benthic cvanobacterial mats have been 72 associated with decreased macro-invertebrate diversity (Aboal et al., 2002), few studies have 73 investigated the genuine toxicological effects of these compounds on aquatic organisms (Carneiro et al., 2015; Osswald et al., 2007b; Anderson et al., 2018). In some experiments 74 75 performed on fishes, such as carps and goldfish, behaviourial defects such as rapid opercular 76 movement, abnormal swimming (Osswald et al., 2007a) and muscle rigidity (Carmichael et al., 77 1975) were observed. Oberemm et al. (1999) described also the alterations in heart rates in 78 zebrafish embryos after an exposure to anatoxin-a. Thus a scarcity of information regarding its 79 capability of aquatic species to bio-concentrate and bioaccumulation anatoxin-a when 80 administrated by natural oral pathways, and its subsequent potential toxicological impacts on 81 organisms still exists. In a previous study, Osswald and co-workers (2007b) found that 82 anatoxin-a may be bioaccumulated by carps in significant levels (0.768 μ g.g⁻¹ of carp weight).

- 83 Whether this may have an impact in aquatic food webs is not yet known.
- 84

85 In this study, we wanted to study the toxico-kinetics of the anatoxin-a and its consecutive 86 possibility of accumulation of in fish tissues. We use medaka model fish in order to be able to 87 administrate, under controlled conditions by a single gavage, predetermined dose of anatoxin-88 a in order to determine the dose-response toxicity parameters and to follow the 89 assimilation/depuration processes of fish gavaged with a no observable adverse effect level 90 (NOAEL) dose during the 10 following days. As there is still a lack of reference and 91 standardized protocol for anatoxins-a extraction from biological matrix, such as fish tissues, 92 and for quantification analysis, we have also tested three different extraction procedures 93 inspired by previously published works (Triantis et al., 2016) and describe a high accuracy 94 detection method developed on UHPLC coupled HR-gTOF mass spectrometer using TASQ 95 software. This present work provides significant outcomes for the investigation of fish 96 contamination by anatoxin-a.

97

98 Material and methods

99 Chemicals

In solution certified (+)-Anatoxin-a and dry (±)-Anatoxin-a were purchased from CRM-NRC (Canada) and Abcam (UK), respectively. The purity and the concentration of the daily reconstituted (±)-Anatoxin-a in ultra-pure water was initially checked by LC–MS/MS as described in the following protocol. UHPLC-MS grade methanol and acetonitrile were purchased from Bio TechnoFix (France). Proteomics grade formic acid was purchased from Sigma-Aldrich (Germany).

106

107 Cyanobacteria cultures and Anatoxin-a extraction procedure test

Three monoclonal non-axenic cultures of Phomidium sp. (PMC 1001.17, 1007.17 and 108 109 1008.17) maintained at 25°C in 15-mL vessels with Z8 media in the PMC (Paris Museum 110 Collection) of living cyanobacteria. Larger volume of all strains was simultaneously cultivated during one month in triplicates in 250 mL Erlenmeyer vessels at 25°C using a Z8 medium with 111 a 16h:8h light/dark cycle (60 μ mol.m⁻².s⁻¹). Cyanobacterial cells were centrifuged (at 4,000 g 112 for 10 min), then freeze-dried and stored at -80°C prior to anatoxin-a extraction. The 113 114 lyophilized cells were weighted, sonicated 2 min in a constant ratio of 100 µL of solvent for 1 115 mg of dried biomass, centrifuged at 4°C (12,000 g; 10 min), then the supernatant was collected 116 and directly analysed by mass spectrometry. We have presently tested in triplicates the 117 extraction efficiency of three different solvent mixtures already propositioned for anatoxin-a extraction (Bogialli et al., 2006; Rellan et al., 2007; Triantis et al., 2016; Haddad et al., 2019), 118 119 comprising: a pure water solution acidified with 0.1% formic acid ("Water" extraction), a 25-120 % acetonitrile solution acidified with 0.1% formic acid ("Acetonitrile" extraction), and a 75-% methanol solution acidified with 0.1% formic acid ("Methanol" extraction). The efficiency of 121 122 the extraction procedure was significantly determined according to Dunn's post-hoc test 123 performed after Krustal-Wallis non-parametric tests process on R software.

- 124
- 125 Fish experimentation

126 Experiments on medaka fish was conducted according to European community good 127 practices. validated by the ethical "comité Cuvier" (Author. APAFiS#19417-128 2019022711043436 v4) and under the supervision of accredited personnel (B.M.). Adult female 129 medaka fishes (*Oryzias latipes*) of the inbred cab strain and of above 1 ± 0.1 g of wet weight were used in all experiments. They were raised in 20-L glass aquaria filled with a continuously 130 131 aerated mixture of tap water and reverse osmosis filtered water (1/3-2/3, respectively), which 132 was changed once a week. Fish were maintained at 25 ± 1 °C, with a 12 h:12 h light:dark 133 standard cycle.

134 Fish were individually anesthetized in 0.1% tricaine methane sulfonate (MS-222; Sigma, St. Louis, MO), and then, briefly, 2 μ L of a (±)-anatoxin-a mixture containing from 0.2 to 20 μ g 135 of (±)-anatoxin-a in water saturated with phenol red dyes was administrated by gavage 136 137 performed with a smooth plastic needle. Control fishes were gavaged with 2 µL of water 138 saturated with phenol red dyes. For all individual, the efficiency of the gavage uptakes was 139 carefully checked according to the total lack of phenol red release from the mouth or the gill opercula of the fish, otherwise the individual was immediately sacrificed. The fish was then 140 141 instantaneously placed in fresh water tanks and individually observed during 30 min in order 142 to detect any behavioural sign of anatoxin-a neurotoxicity, that comprises: paralysis, decrease of breathing activity through opercular movements, locomotors activity or buoyancy default. 143 144 Indeed, when gavaged with none toxic dose or control mixture, fish promptly recovers from 145 anaesthesia within a minute and rapidly normal swimming activity (in less than 3-5 minutes). 146 But, alternatively, the toxicological effects of toxic doses of (\pm) -anatoxin-a induce immediate 147 neuro-muscular pathology and provoke a full breathing stop. After 30 min of observation, the 148 individual was declared as "dead" as no recover was observed and the experiment was then 149 concluded, then all fishes were anesthetized in 0.1% tricaine methane sulfonate (MS-222) and 150 euthanized. The medium lethal dose (LD_{50}) and the no observable adverse effect limit (NOAEL) were calculated from toxicological results by logistic regression after log 151 152 transformation of the concentration values using R software.

For toxico-kinetics investigations, adult female medaka fish were similarly gavaged individually by a single NOAEL dose, then placed in fresh water and collected after 1h, 3h, 6h, 12h, 24h, 3d, 7d or 10d of maintaining under classical conditions. Accordingly, all fishes were anesthetized in 0.1% tricaine methane sulfonate, sacrificed, dissected, and the whole gut, the liver and the muscles were sampled and flash-frozen in liquid nitrogen, and kept frozen at -80°C prior to analysis.

159

160 Anatoxin-a and metabolite extraction from fish tissues

The fish tissues were weighted then sonicated 2 min in a constant ratio of 10 µL of 75-% 161 162 methanol solution acidified with 0.1% formic acid for 1 mg of wet biomass for guts and livers, 163 and of lyophilised biomass for muscles, grinded into a fine powder on Tissue-lyser (with 5 mm steel beads, Oiagen), centrifuged at 4°C (12,000 g; 10 min); then the supernatant was collected 164 and directly analysed by mass spectrometry. The efficiency of the extraction was estimated 165 166 according to the recovery rate determined in triplicats by injecting known amount of (\pm) -167 anatoxin-a to negative samples before extraction (method A) or just before the mass 168 spectrometry analysis (method B).

170 Anatoxin-a detection and quantification

171 Ultra high performance liquid chromatography (UHPLC) was performed on 2 µL of each of the metabolite extracts using a Polar Advances II 2.5 pore C₁₈ column (Thermo) at a 300 172 173 μ L.min⁻¹ flow rate with a linear gradient of acetonitrile in 0.1% formic acid (5 to 90 % in 21 174 min). The metabolite contents were analyzed in triplicate for each strain using an electrospray 175 ionization hybrid quadrupole time-of-flight (ESI-QqTOF) high resolution mass spectrometer (Maxis II ETD, Bruker) at 2 Hz speed on simple MS mode and subsequently on broad-band 176 177 Collision Ion Dissociation (bbCID) MS/MS mode on the 50-1500 m/z rang. Calibrants, 178 composed by serial dilutions of (+)-anatoxin-a, and negative control (phenylalanine) were 179 analysed similarly. The raw data were automatically process with the TASQ 1.4 software for 180 internal recalibration (< 0.5 ppm for each sample, as an internal calibrant of Na formate was 181 injected at the beginning of each analysis) for global screening and quantification, and 182 molecular featuring, respectively. Then, the automatic screening and quantification of anatoxin-183 a was performed with threshold parameters set to recommended default value for the Maxis II mass spectrometer (Δ RT < 0.4 s, Δ m/z < 3 ppm, mSigma < 50 and S/N < 5). Quantification 184 was performed according to the integration of the area under the peaks and calibration curve 185 186 was performed with certified standards.

187

188 Analysis of liver metabolomes

189 Metabolites composition of the fish livers were analysed by injection of 2 µL of the 75% 190 methanol extracts on an UHPLC (ELUTE, Bruker) coupled with a high-resolution mass 191 spectrometer (ESI-Qq-TOF Compact, Bruker) at 2 Hz speed on simple MS mode and subsequently on broad-band Collision Ion Dissociation (bbCID) or autoMS/MS mode on the 192 193 50-1,500 m/z rang. The analyte annotations were performed according to precise mass and 194 isotopic and fragmentation MS/MS patterns, as previously described (Kim Tiam et al., 2019). 195 The feature peak list was generated from recalibrated MS spectra (< 0.5 ppm for each sample, 196 as an internal calibrant of Na formate was injected at the beginning of each analysis) within a 197 1-15 min window of the LC gradient, with a filtering of 5,000 count of minimal intensity, a 198 minimal occurrence in at least 50% of all samples, and combining all charge states and related 199 isotopic forms using MetaboScape 4.0 software (Bruker). The intensity data table of the 591 200 extracted analytes was further treated using MetaboAnalyst 4 tool (Chong et al., 2019) for 201 Pareto's normalization, ANOVA, PCA and PLS-DA, and data representation by heatmap with 202 hierarchical clustering, loading plots and box plots.

203 Unsupervised PCA models were first used to evaluate the divide between experimental 204 groups, while supervised PLS-DA models allowed us to increase the separation between sample 205 classes and to extract information on discriminating metabolites. The PLS-DA allowed the 206 determination of discriminating metabolites using the analytes score values of the variable 207 importance on projection (VIP) indicating the respective contribution of a variable to the 208 discrimination between all of the experimental classes of samples. The higher score being in 209 agreement with a strongest discriminatory ability and thus constitutes a criterion for the 210 selection of the analytes as discriminative components. The PLS models were tested for over fitting with methods of permutation tests. The descriptive, predictive and consistency 211 performance of the models was determined by R^2 , Q^2 values and permutation test results (n = 212 213 100), respectively.

215 **Results**

216 Anatoxin-a extraction and analysis

217 Anatoxin-a specific mass spectrometry detection and quantification was automatically 218 determined with high accuracy using TASO (Bruker, Germany) from raw data generated by 219 LC-MS/MS system according to the observation of analytes signal exhibiting targeted precise 220 molecular mass, retention time, isotopic pattern and fragmentation ions (Fig. 1A-B). This 221 approach allows to discriminate both (+)-anatoxin-a and (-)-anatoxin-a isomers according to 222 their respective retention times (Fig. 1C), as well as phenylalanine (Fig. 1D) used as negative 223 control, that does not exhibit any signal interaction with the anatoxin-a quantification. Standard 224 solutions containing certified quantity of (+)-anatoxin-a was diluted in ultra-pure water in the range of 5 μ g.mL⁻¹ - 2 ng.mL⁻¹ and used for calibration with good linearity of the calibration 225 curve exhibiting a correlation coefficient with $R^2 = 0.99326$ (Fig. 1E). Ouantification of the 226 227 (+)-anatoxin-a was performed according to the area-under-the-curve signal that was 228 automatically integrated and process by the software that provide calculation details comprising 229 all diagnostic elements in the report table generate for each analysis (Fig 1F).

230 Anatoxin-a extraction efficiency was assayed in triplicates with 3 different Phormidium 231 strains using rather acetonitrile, methanol of water solvent solutions. In our hands, the 75% 232 methanol acidified with 0.1% formic acid present significantly higher efficiency for anatoxin-233 a extraction (Table 1) and was further performed for the fish tissue extraction. Instrument and 234 method detection and quantitation limits (LOD and LOQ) and recovery rate were determined 235 by using triplicate injections of the control different fish tissues spiked with of determined doses 236 of (+)-anatoxin-a that were administrated before the extraction (method A), or just before the 237 sample analysis by mass spectrometry (method B) (table 2). The LOD and LOQ of the technics 238 was estimated to be in the same rang as those previously described (Triantis et al., 2016). These 239 investigations also show that anatoxin-a extraction and detection recovery rate vary from 25-78% and 51-126%, respectively, according to the tissue analysis with the guts and the muscles 240 241 presenting the best and the worth recovery scores, respectively.

242

243 Anatoxin-a toxicology

244 All individuals, including negative controls, gavaged with (\pm) -anatoxin-a doses up to 6.67 $\mu g.g^{-1}$ survive without presenting any apparent symptoms of toxicosis and were able to recover 245 246 from the tricaine sedation within less than 3 minutes when placed in fresh water. On contrary, all individual gavaved with 20 μ g.g⁻¹ (±)-anatoxin-a rapidly present (in the first 5 minutes) 247 obvious signs of neurotoxic effects, comprising a complete stop or a rapid diminution of 248 249 opercular movement, abnormal swimming with hemi- or complete paresis of the fins, 250 accompanied with a global musculature rigidity. After 10 min, only one individual even 251 presents few sporadic breathing activity, that completely stop after 15 min, when all other organisms already present complete paresis and ventilation cease. After 30 min, these 252 253 individuals, presenting no sign of recovery, were considered as dying, if not dead, and euthanized, then were considered for further toxicological dose calculation. When LD₁₀₀ and 254 the NOAEL were observed at 20 μ g.g⁻¹ and 6.67 μ g.g⁻¹(±)-anatoxin-a, respectively, the LD 50 255 was calculated as 11.5 μ g.g⁻¹ of (±)-anatoxin-a (Fig. 2A; table 3). 256

258 Anatoxin-a toxico-kinetics

259 Following gavage experiment of adult female medaka fish to NOAEL, we have quantified 260 (±)-anatoxin-a in the guts, the livers and the muscles after 1h, 3h, 6h, 12h, 24h, 3d, 6d and 10d, 261 in order to monitor the dynamic of the anatoxin-a assimilation/depuration efficiency in these 262 various compartments (Fig. 1B-D). The highest (±)-anatoxin-a amount was observed just 1h after the exposure, representing up to $15,789 \,\mu g.g^{-1}$ in some individual livers, representing more 263 than 100 times more anatoxin-a than in the livers of other organisms similarly gavaged, 264 265 illustrating the relative individual variability of the anatoxin-a uptake and 266 assimilation/elimination during our experimentation. Although, fish tissues present a global individual variability, the larger amount of anatoxin-a were observed in livers, the guts, and in 267 a lesser amount in the muscles after 1h, then the tissues present a rapidly decrease of anatoxin-268 269 a contents. Almost no more anatoxin-a was detectable after 24h in all tissues. The depuration 270 rate was calculated for the 12 first hours of depuration as being of 57, 100 and 90%, in guts, 271 livers and muscles, respectively, leading to a rapid elimination of the anatoxin-a that does no 272 seems to bio-accumulate in any of the investigated fish tissues.

273

274 Anatoxin-a effects on the liver metabolome

275 In order to investigate the molecular effects induced by (\pm) -anatoxin-a NOAEL exposure, 276 the liver metabolite composition was compared between ungavaged fish (control) and fish 277 collected 1h, 3h, 6h, 12h or 24h after gavage. The same livers extracts than those extracted with 278 75% methanol and analysed for anatoxin-a quantification were investigated by LC-MS/MS for 279 untargeted metabolomics. A total of 591 different analytes were then extracted by the optimized 280 pipeline, and their respective quantification (determined from the area-under-the-peak signal) 281 compared between the different time-course groups using multivariate statistical methods, 282 including unsupervised principal component analysis (PCA) and supervised partial leastsquares discriminate analysis (PLS-DA), together with univariate groups variance analyses 283 284 (ANOVA), in order to evaluate the potential of the method to discriminate among the 285 experimental groups according to the time-course of anatoxin-a exposure and elimination.

286 Although the analysed fish livers present a global metabolome variability (Fig. 3A), the (\pm) -287 anatoxin-a gavage seems to rapidly modify the specific amount of various metabolites which progressively retrieved their initial state, as observed on components 1-3 projection of the PCA 288 289 (Fig. 3B). The 29 analytes that present significant variation between the different groups 290 according to ANOVA (P < 0.05) indicate a clear difference between the groups, as observed on 291 heatmap with hierarchical clustering (Fig. 3D), with a rapid increase or decrease of the 292 metabolite quantity that diminishes after few hour and almost recover control levels after 24h post-gavage (Fig. 3E). The supervised multivariate analysis (PLS-DA) model shows consistent 293 R^2 cumulative, Q^2 cumulative and permutation scores (Fig. 3F). The most discriminating m/z294 295 features in the PLS-DA model (Figure 3C) were selected based on their respective VIP score, 296 which resulted in 25 compounds with VIP value higher than 2 (Table 4), on component 1 and/or 297 2 (both contributing to the experimental group discrimination). The molecular formulas of each VIP were proposed based on accurate mass measurement, true isotopic pattern, and their 298 299 putative identification were attempted with Metfrag and GNPS according to additional 300 respective MS/MS fragmentation patterns. Interestingly, the tricaine (MW 165.0794 Da) belong to this VIP list and presents, as one could expect, a clear increase between the Control
 and 1h post-gavage fish livers (in relation with pre-gavage anesthesia procedure by balneation
 in 0.1% tricaine), then a complete disappearance between 1 and 3h.

304 The list of other putatively annotated compounds comprises then various phospholipids (n=9) belonging to the glycerol-phosphocholine group, all presenting comparable variation 305 306 patterns among the experimental groups, with an initial drop of these metabolite quantity 307 between the control and the 1h group, then a progressive re-increase until an almost complete 308 recovery of the initial metabolite amount within less than 24h. Those metabolites are directly 309 related to lipid metabolism process and their successive decrease and increase within the liver may denote an important lipid consumption by the organism and a progressive recover of the 310 311 liver to an unstressed condition. On the contrary, the sole VIPs which relative quantities present 312 an initial increase were an undetermined analyte (MW 1064.600 Da) and the adenosine, both 313 retrieving their initial levels within less than 24h. Such adenosine transient increase may denote 314 an intensification of hepatic blood circulation, as it presents direct effect on vasodilation of liver 315 arteria (Robson & Schuppan 2010). Although our experimental design does allow to 316 discriminate the specific effects of the anatoxin-a from those of the tricaine anesthesia alone, it 317 overall shows that the organisms present a complete recovery 24h after being gavaged with 318 anatoxin-a single NOAEL dose.

320 Discussion

319

321 In the present study, the neurotoxicological response of medaka fish subjected to anatoxin-322 a appears comparable to that found in carp (Osswald et al., 2007a) and zebrafish (Carneiro et 323 al., 2015) supporting the evidences for the existence of a similar mechanism of action. Whereas 324 no apparent precursor effect appears when fish are gavaged with sub-acute doses of anatoxin-325 a, the symptoms observed in the fish exposed to a higher dose denote an all-or-nothing effect, 326 comprising rapid and intense neurotoxic signs, that are compatible with the mechanism of toxic 327 action of anatoxin-a in the nervous system that has been described so far for other vertebrates (Fawell et al., 1999). These symptoms comprise different manifestations of muscular paralysis 328 329 that are likely due to the primary, and nearly irreversible, binding of the anatoxin-a, being an 330 acetylcholine antagonist, to the acetylcholine receptors of the cholinergic synapses of the 331 neuromuscular junctions, leading to a continuous muscular contraction. Previous works 332 indicate that below a certain levels the effects appear to be transient, the animals being able to 333 make a complete and rapid recovery, although the data available in the literature remains limited 334 (Dittman and Wiegand 2006).

335 Our dose-response toxicological analysis indicates that anatoxin-a exhibits toxicological reference dose of 5.75 and 3.33 µg.g⁻¹ of medaka body weight for LD₅₀ and NOAEL, 336 respectively. This results are in remarkable agreement with precedent data obtained on other 337 338 organisms. Indeed, previous investigations of anatoxin-a toxicity determined on mouse that, 339 when administrated by single dose gavage, (+)-anatoxin-a exhibits a LD₅₀ and a NOAEL of above 5 and 3 µg.g⁻¹ of body weight, respectively (Fawell et al., 1999). In addition, Stevens and 340 co-workers (1991) have initially shown that (+)-anatoxin-a presents LD₅₀ value of 16.2 and 6.7 341 µg.g⁻¹ of body weight, for pure (+)-anatoxin-a and complex extract containing anatoxin-a 342 administrated by single gavage to mouse. Taken together, those results suggest that medaka 343 344 fish and mouse apparently present very similar toxicological dose-response to anatoxin-a.

Indeed, as the unnatural (-)-anatoxin-a isomer was shown to insignificantly contribute to the global toxicity to the 1/1 racemic mixture of (\pm)-anatoxin-a used in this study (MacPhail et al., 2007), one could estimate that toxicity values of (+)-anatoxin-a might theoretically be very similar for medaka fish than for mouse.

349

350 In our experiments, the promptitude of the neuromuscular effects of anatoxin-a when 351 administrated by gavage are in agreement with previous investigation performed on mice 352 (Fawell et al., 1999), suggesting that anatoxin-a might be rapidly assimilated by the organisms 353 by crossing the intestine barrier and being widespread to the whole musculature within less than 354 2 minutes. When anatoxin-a producing cyanobacterial biofilm are accidently ingested by dogs, 355 the animals present first appearance of neurotoxic symptoms within less than 5 minutes (Wood 356 et al., 2007), testifying for a very rapid assimilation of a toxinogenous dose of anatoxin-a 357 released from the cyanobacterial biomass within the stomach. To date, the genuine mechanism 358 of the anatoxin-a transfer through the intestinal epithelia remains undetermined but the 359 promptitude of the effects suggests that this small molecule (MW = 165 da) may remain 360 uncharged in order to be able to sharply cross the intestinal barrier, potentially through passive 361 paracellular diffusion (Dahlgren and Lennernäs 2019).

362 Our toxico-kinetics investigation has shown that no detectable amount of anatoxin-a was 363 still observed in guts 12h after the fish having been gavaged to a single NOAEL dose. Then, 364 the medaka fish presents also a rapid elimination of the anatoxin-a that transiently have been addressed to the liver, and in a lesser extent to the muscle, indicating that it may not being 365 366 apparently accumulating anatoxin-a under those conditions. Interestingly, Osswald and co-367 workers (2011) have shown no significant bioaccumulation of anatoxin-a in the trout tissues when administrated by balneation up to 5 mg. L^{-1} of anatoxin-a. Previously, Osswald et al. 368 (2008) have also similarly shown no bioaccumulation of anatoxin-a by the Mediterranean 369 370 mussel *M. galloprovincialis*, that has been otherwise shown to present high accumulation 371 capability for various contaminants, such as microcystins (Vasconcelos 1995). Although they observed some anatoxin-a uptake from the surrounding water filtered by the animals (observed 372 373 maximum accumulation efficiency = 11%), this seemed rapidly reduced rather by the 374 depuration process of phase II detoxification enzymes or by passive elimination processes. These data show that, as anatoxin-a is capable to rapidly penetrate the whole organisms body. 375 376 and induces obvious and acute neurotoxicological effects within a minute, it may also be rapidly 377 eliminated by classical excretion or depuration mechanisms.

378

379 Conclusion

In our gavage experiment of medaka fish to a single NOAEL dose of anatoxin-a, the toxin appears to have been rapidly eliminated and the molecular effects were no more perceptible within the fish liver metabolome after 24h.

These observations suggest that when the dose remains below the acute toxicological limit producing neurotoxicosis that can lethal consequences, the organism are able to make a completely and rapidly recovery, that seems not to induce obvious effects even if the exposure is repeated several times (Fawell et al. 1999). However, one could still suspect that chronic anatoxin-a exposure may induce more insidious long-term pathologies on the central nervous system (Lombardo and Maskos 2015). But this hypothesis still remains to be explored.

- More over, the accurate investigations of the fish flesh contamination by anatoxin-a under highest toxinogenous cyanobacterial proliferations, especially when, in nature, certain fish are potentially actively feeding on those biofilms (Ledreux et al., 2014), still remain to be
- 392 performed in order to provide convincing data supporting the evaluation of the associated risks.
- 393
- 394

395 Acknowledgements

This work was supported by grant CRD from ANSES attributed to the Cyanariv project, leadby Catherine Quiblier.

- 398
- 399 The authors declare not conflict of interest
- 400 401

402 References

- Aboal, M., Puig, M. A., Mateo, P., & Perona, E. (2002). Implications of cyanophyte toxicity on biological
 monitoring of calcareous streams in north-east Spain. *Journal of Applied Phycology* 14(1), 49-56.
- Anderson, B., Voorhees, J., Phillips, B., Fadness, R., Stancheva, R., Nichols, J., ... & Wood, S. A. (2018).
 Extracts from benthic anatoxin-producing *Phormidium* are toxic to 3 macroinvertebrate taxa at
 environmentally relevant concentrations. *Environmental toxicology and chemistry* 37(11), 2851-2859.
- 408 Aráoz, R., Molgó, J., & De Marsac, N. T. (2010) Neurotoxic cyanobacterial toxins. *Toxicon* 56(5), 813-828.
- Bogialli, S., Bruno, M., Curini, R., Di Corcia, A., & Laganà, A. (2006). Simple and rapid determination of
 anatoxin-a in lake water and fish muscle tissue by liquid-chromatography–tandem mass spectrometry. *Journal of Chromatography A* 1122(1-2), 180-185.
- Bouma-Gregson, K., Kudela, R. M., & Power, M. E. (2018). Widespread anatoxin-a detection in benthic
 cyanobacterial mats throughout a river network. *PloS One* 13(5), e0197669.
- 414 Carmichael, W. W., Biggs, D. F., & Gorham, P. R. (1975). Toxicology and pharmacological action of 415 *Anabaena flos-aquae* toxin. *Science* 187(4176), 542-544.
- 416 Carmichael, W. W. (1994). The toxins of cyanobacteria. *Scientific American* 270(1), 78-86.
- 417 Carneiro, M., Gutierrez-Praena, D., Osorio, H., Vasconcelos, V., Carvalho, A. P., & Campos, A. (2015).
 418 Proteomic analysis of anatoxin-a acute toxicity in zebrafish reveals gender specific responses and
 419 additional mechanisms of cell stress. *Ecotoxicology and environmental safety* 120, 93-101.
- Chong, J., Wishart, D. S., & Xia, J. (2019). Using metaboanalyst 4.0 for comprehensive and integrative
 metabolomics data analysis. *Current protocols in bioinformatics* 68(1).
- 422 Dahlgren, D., & Lennernäs, H. (2019). Intestinal Permeability and Drug Absorption: Predictive
 423 Experimental, Computational and *In Vivo* Approaches. *Pharmaceutics* 11(8), 411.
- Dittmann, E., & Wiegand, C. (2006). Cyanobacterial toxins–occurrence, biosynthesis and impact on human
 affairs. *Molecular nutrition & food research* 50(1), 7-17.
- Fawell, J. K., Mitchell, R. E., Hill, R. E., & Everett, D. J. (1999). The toxicity of cyanobacterial toxins in the
 mouse: II anatoxin-a. *Human & experimental toxicology* 18(3), 168-173.
- Gugger, M., Lenoir, S., Berger, C., Ledreux, A., Druart, J. C., Humbert, J. F., ... & Bernard, C. (2005). First
 report in a river in France of the benthic cyanobacterium Phormidium favosum producing anatoxin-a
 associated with dog neurotoxicosis. *Toxicon* 45(7), 919-928.
- Greer, B., Maul, R., Campbell, K., & Elliott, C. T. (2017) Detection of freshwater cyanotoxins and
 measurement of masked microcystins in tilapia from Southeast Asian aquaculture farms. *Analytical and Bioanalytical Chemistry* 409(16), 4057-4069.
- 434 Haddad, S. P., Bobbitt, J. M., Taylor, R. B., Lovin, L. M., Conkle, J. L., Chambliss, C. K., & Brooks, B. W.
- 435 (2019). Determination of microcystins, nodularin, anatoxin-a, cylindrospermopsin, and saxitoxin in water

- and fish tissue using isotope dilution liquid chromatography tandem mass spectrometry. *Journal of chromatography*. A 1599, 66.
- Kim Tiam, S., Gugger, M., Demay, J., Le Manach, S., Duval, C., Bernard, C., & Marie, B. (2019). Insights
 into the Diversity of Secondary Metabolites of *Planktothrix* Using a Biphasic Approach Combining
 Global Genomics and Metabolomics. *Toxins* 11(9), 498.
- Ledreux, A., Brand, H., Chinain, M., Bottein, M. Y. D., & Ramsdell, J. S. (2014). Dynamics of ciguatoxins
 from Gambierdiscus polynesiensis in the benthic herbivore Mugil cephalus: trophic transfer implications. *Harmful Algae* 39, 165-174.
- Lombardo, S., & Maskos, U. (2015). Role of the nicotinic acetylcholine receptor in Alzheimer's disease
 pathology and treatment. *Neuropharmacology* 96, 255-262.
- MacPhail, R. C., Farmer, J. D., & Jarema, K. A. (2007). Effects of acute and weekly episodic exposures to
 anatoxin-a on the motor activity of rats: comparison with nicotine. *Toxicology* 234(1-2), 83-89.
- Oberemm, A., Becker, J., Codd, G. A., & Steinberg, C. (1999). Effects of cyanobacterial toxins and aqueous
 crude extracts of cyanobacteria on the development of fish and amphibians. *Environmental Toxicology: An International Journal* 14(1), 77-88.
- Osswald, J., Rellan, S., Gago, A., & Vasconcelos, V. (2007a). Toxicology and detection methods of the
 alkaloid neurotoxin produced by cyanobacteria, anatoxin-a. *Environment international* 33(8), 1070-1089.
- Osswald, J., Rellán, S., Carvalho, A. P., Gago, A., & Vasconcelos, V. (2007b). Acute effects of an anatoxin a producing cyanobacterium on juvenile fish—Cyprinus carpio L. *Toxicon* 49(5), 693-698.
- Osswald, J., Rellán, S., Gago, A., & Vasconcelos, V. (2008). Uptake and depuration of anatoxin-a by the
 mussel *Mytilus galloprovincialis* (Lamarck, 1819) under laboratory conditions. *Chemosphere* 72(9),
 1235-1241.
- Osswald, J., Azevedo, J., Vasconcelos, V., & Guilhermino, L. (2011). Experimental determination of the
 bioconcentration factors for anatoxin-a in juvenile rainbow trout (*Oncorhynchus mykiss*). Proceedings of
 the International Academy of Ecology and Environmental Sciences 1(2), 77.
- Puschner, B., Pratt, C., & Tor, E. R. (2010). Treatment and diagnosis of a dog with fulminant neurological
 deterioration due to anatoxin-a intoxication. *Journal of veterinary emergency and critical care* 20(5), 518522.
- Quiblier, C., Wood, S. A., Echenique-Subiabre, I., Heath, M., Villeneuve, A., & Humbert, J. F. (2013). A
 review of current knowledge on toxic benthic freshwater cyanobacteria–ecology, toxin production and
 risk management. *Water Research* 47(15), 5464-5479.
- Rellán, S., Osswald, J., Vasconcelos, V., & Gago-Martinez, A. (2007). Analysis of anatoxin-a in biological
 samples using liquid chromatography with fluorescence detection after solid phase extraction and solid
 phase microextraction. *Journal of Chromatography A* 1156(1-2), 134-140.
- 470 Robson, S. C., & Schuppan, D. (2010). Adenosine: tipping the balance towards hepatic steatosis and fibrosis.
 471 *Journal of hepatology* 52(6), 941-943.
- Sivonen, K., Jones, G. (1999) Cyanobacterial toxins. In: Chorus, I., Bartram, J. (Eds.), Toxic Cyanobacteria
 in Water: a Guide to Their Public Health Consequences, Monitoring and Management. E &FN Spon,
 London, pp. 41–111.
- 475 Stevens, D. K., & Krieger, R. I. (1991). Stability studies on the cyanobacterial nicotinic alkaloid snatoxin-A.
 476 *Toxicon* 29(2), 167-179.
- Triantis, T. M., Kaloudis, T., & Hiskia, A. (2016). Determination of Anatoxin-a in Filtered and Drinking
 Water by LC-MS/MS. *Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis*, 408-412.
- 479 Vasconcelos, V. M. (1995). Uptake and depuration of the heptapeptide toxin microcystin-LR in Mytilus
 480 galloprovincialis. *Aquatic Toxicology* 32(2-3), 227-237.
- Wonnacott, S., & Gallagher, T. (2006). The chemistry and pharmacology of anatoxin-a and related
 homotropanes with respect to nicotinic acetylcholine receptors. *Marine Drugs* 4(3), 228-254.
- 483 Wood, S. A., Selwood, A. I., Rueckert, A., Holland, P. T., Milne, J. R., Smith, K. F., ... & Cary, C. S. (2007).
- 484 First report of homoanatoxin-a and associated dog neurotoxicosis in New Zealand. *Toxicon* 50(2), 292-
- 485 301.





488 489 Figure 1. Anatoxin-a detection and quantification on HR qTOF mass spectrometer coupled to an UHPLC. 490 The screening of (+)-anatoxin-a was performed according to the specific detection of analytes presenting 491 accurate mass of the parent ion (M+H⁺= 166.1232 \pm 0.001 Da), accurate retention time (1.55 \pm 0.2 min, 492 accurate isotopic pattern (mSigma < 50) and co-detection of characteristic fragment ions (149.096 \pm 0.001 493 and 131.086 ± 0.001 Da) (A-B). The similar analyses of the (±)-anatoxin-a can discriminate the (-) and the 494 (+)-anatoxin-a that exhibits distinct retention times (1.1 and 1.6 min, respectively) (C). No signal was 495 detected with phenylalanine (D). Calibration curve performed with 12 serial dilutions of certified standard 496 of (+)-anatoxin-a (NRC, Canada) presenting applicable correlation factor (E). Example of screening 497 diagnostic table provided by TASQ® software for (+)-anatoxin-a quantification summarizing all qualitative 498 and quantitative parameters for each sample (F).





501 502 Figure 2. Anatoxin-a toxicity and toxico-kinetics on adult female medaka fish when administrated by single-503 dose gavage. Estimation of the LD_{50} of (±)-anatoxin-a by linear regression after log transformation (µg of 504 (±)-anatoxin-a.g⁻¹ of fish mass) (A). Toxico-kinetics of (±)-anatoxin-a administrated in a single NOAEL dose 505 (6.67 µg.g⁻¹ fish weight) in the gut (B), the liver (C) and the muscle (D) after 1h, 3h, 6h, 12h, 24h, 3j, 6j or 506 10j of depuration (n-5). On the 30 investigated individuals, only 6 of them does not present detectable amount 507 of anatoxin-a, overall testifying for the global efficiency of the gavage experiments.





510 Figure 3. Metabolomics investigation of the liver metabolite variations after the administrated of a single NOAEL dose of (±)-anatoxin-a (6.67 μ g.g⁻¹ fish weight) by gavage of adult female medaka fish in the liver 511 512 after 1h, 3h, 6h, 12h and 24h. Individual score plots generated from a principal component analysis performed 513 with the intensity count of 591 variables according to components 1-2 (A) and 1-3 (B). Heatmap 514 representation with hierarchical classification (Ward clustering according to Euclidian distances) performed 515 from relative intensities of the 29 significantly dysregulated analytes (P <0.05 ANOVA) (D) and 6 examples 516 of representative box-plots (E). Individual score plot generated from PLS-DA analysis performed with 591 517 extracted variables according to components 1-2 (C), and quality descriptors of the statistical significance 518 and predictive ability of the discriminant model according to corresponding permutation and cross-validation 519 tests (F), respectively.

521 Tables

577	
7//	
222	

Dhamidium stuain	Extraction solvant			
Phormialum strain	Acetone	Water	Methanol	
PMC 1001.17	6436 ± 139^{a}	5864 ± 85^{b}	6689 ± 158^{a}	
PMC 1007.17	2949 ± 148^{b}	$1467 \pm 88^{\circ}$	3655 ± 147^{a}	
PMC 1008.17	1773 ± 95	1420 ± 103	1913 ± 192	

524 Table 1. Efficiency of different protocols for anatoxin-a extraction from different cyanobacterial strains. ^{a-b-}

525 ^c: indicate group results in a decreasing order of Dunn's post-hoc test performed after Krustal-Wallis non-526 parametric tests.

527

528

Recovery rate		Limit of detection		Limit of quantification		
Organ			(LOD)		(LOQ)	
	Method A	Method B	μg.L ⁻¹	$\mu g.g^{-1}$	μ g.L ⁻¹	$\mu g.g^{-1}$
Intestin ^a	78 ± 13 %	126 ± 22%	0.412	20.6	1.25	62.5
Liver ^a	65 ± 7 %	86 ± 15%	0.506	2.53	1.25	62.5
Muscle ^b	25 ± 4 %	51 ± 3%	0.546	14.52	1.25	285

529

530 Table 2. Recovery rate, limit of detection (LOD), and limit of quantification (LOQ) of the two tested 531 extraction procedure (A and B, corresponding to pre-extraction or pre-analysis doping, respectively) 532 measured on intestine, liver and muscle of the medaka fish. ^a: determined on fresh weight; ^b: determined on 533 dry weight.

534

Model organism	Toxicological parameter	Toxicological values (µg.g ⁻¹)	Formulation of the toxicant	Reference
Medaka fish	LD ₅₀ by gavage	11.5	(±)-anatoxin-a	This study
Medaka fish	LD ₅₀ by gavage	5.75*	(+)-anatoxin-a	This study
Medaka fish	LD ₁₀₀ by gavage	20	(±)-anatoxin-a	This study
Medaka fish	LD ₁₀₀ by gavage	10*	(+)-anatoxin-a	This study
Medaka fish	NOAEL by gavage	6.67	(±)-anatoxin-a	This study
Medaka fish	NOAEL by gavage	3.33*	(+)-anatoxin-a	This study
Zebrafish	LD ₁₀₀ by <i>i.p</i> .	0.8	(±)-anatoxin-a	Carneiro et al., 2015
Mouse	LD ₅₀ by <i>i.v.</i>	0.1	(+)-anatoxin-a	Fawell et al., 1999
Mouse	LD ₅₀ by gavage	[1-10]	(+)-anatoxin-a	Fawell et al., 1999
Mouse	NOAEL by gavage	3	(+)-anatoxin-a	Fawell et al., 1999
Mouse	LD ₅₀ by gavage	6.7	A. flos aquae NRC 44-1	Stevens et al., 1991
Mouse	LD ₅₀ by gavage	16.2	(+)-anatoxin-a	Stevens et al., 1991

* determined according to the theoritical 1/1 (±)-racemic ratio, experimentally confirmed by LC-MS,
 considering as negligible the toxicity of (-)-anatoxin-a comparing to the 150-times higher toxicity of (+) anatoxin-a (Osswald et al., 2007).

538

539 Table 3. Summary of lethal concentration estimators determined for anatoxin-a by different administration

540 pathways on different vertebrate models.

MW(Da)	RT (min)	m/z	Annotation	Formula (neutral)	VIP Com. 1	VIP Com. 2	Pattern (C-1-3-6-12-24h)
541.3166	11.8	542.3224	1-icosapentaenoyl-sn-glycero-3-phosphocholine	$C_{28}H_{48}NO_7P$	8.40	6.37	\/
569.3474	12.8	570.3537	lysophosphatidylcholine 22:5	C ₃₀ H ₅₂ NO ₇ P	7.14	5.92	\/
525.2854	12.2	526.2908	lysophosphatidylethanolamine 22:6	C ₂₇ H ₄₄ NO ₇ P	6.62	7.32	\/
499.2704	12.7	500.3048	tauroursodeoxycholic acid	C ₂₆ H ₄₅ NO ₆ S	4.52	3.88	\\
519.3319	12.6	520.3384	1-linoleoyl-sn-glycero-3-phosphocholine	C ₂₆ H ₅₀ NO ₇ P	4.12	3.39	\
1248.619	5.5	625.3167	-	$C_{30}H_{88}N_{24}O_{29}$	3.20	2.32	\
1134.671	12.4	1135.6760	-	$C_{58}H_{90}N_{10}O_{13}$	3.18	3.06	\
1248.117	5.5	625.0657	-	$C_{70}H_{137}N_{17}O_2$	2.85	2.03	\\-//
1082.640	11.8	1083.6469	-	$C_{58}H_{90}N_4O_{15}$	2.80	2.03	\/
267.0969	1.6	268.1040	adenosine	$C_{10}H_{13}N_5O_4$	2.74	1.89	//-\\
131.0939	1.6	132.1010	leucine	$C_6H_{13}NO_2$	2.66	1.92	\
612.1471	1.6	307.0808	-	$C_{24}H_{37}O_{12}PS_2$	2.59	1.72	\/
307.0837	1.3	308.0908	glutathione	$C_{10}H_{17}N_3O_6S$	2.51	2.13	\//
467.3016	12.0	468.3088	1-myristoyl-sn-glycero-3-phosphocholine	$C_{22}H_{46}NO_7P$	2.46	3.88	\/-
639.4269	4.6	320.7208	-	C ₃₁ H ₅₇ N ₇ O ₇	2.45	1.61	\
523.2561	9.3	524.2634	-	$C_{23}H_{42}NO_{10}P$	2.44	1.55	\
204.0911	3.5	205.0984	tryptophan	$C_{11}H_{12}N_2O_2$	2.41	1.68	\
165.0794	6.4	166.0868	tricaine	$C_9H_{11}NO_2$	2.40	3.56	∕
547.3283	13.6	548.3355	-	C ₂₆ H ₃₇ N ₁₃ O	2.16	1.45	\
1064.600	6.1	533.3073	-	$C_{30}H_{84}N_{18}O_{23}$	1.99	2.15	//-\\
543.3335	12.7	544.3400	1-arachidonoyl-sn-glycero-3-phosphocholine	C ₂₈ H ₅₀ NO ₇ P	1.91	2.66	\-/
1050.577	12.2	1051.5843	-	$C_{56}H_{82}N_4O_{15}$	1.88	2.00	\
515.3015	11.3	516.3085	lysophosphatidylcholine 18:4	$C_{26}H_{46}NO_7P$	1.87	3.46	\/
521.3476	13.6	522.3531	1-elaidoyl-sn-glycero-3-phosphocholine	C ₂₆ H ₅₂ NO ₇ P	1.80	7.50	\\-//
517.3184	11.9	518.3246	1-alpha-linolenoyl-sn-glycero-3-phosphocholine	C ₂₆ H ₄₈ NO ₇ P	1.58	2.18	\-/

Table 4. List of the 25 VIP analytes that present score > 2 on either component 1 or component 2 of the PLS-DA analysis performed with all treatment groups, and their putative annotation according to their respective high resolution mass, isotopic and MS/MS fragmentation patterns, searched against ChEBI, PubChem, HMDB

544 and GNPS databases. "\", "/" and "-" indicate when the metabolites globally decreases, increases or maintains their relative quantity between the time series, respectively.